

## Charged Higgs Boson Search via $e^+e^- \rightarrow H^+H^- \rightarrow c\bar{c}b$ at Linear Colliders

Tuesday 16 May 2023 16:40 (25 minutes)

We study charged Higgs boson search via  $e^+e^- \rightarrow H^+H^- \rightarrow c\bar{c}b$  at the 500 GeV ILC. In a general two Higgs doublet model without  $Z_2$  symmetry, extra Yukawa couplings  $\rho_{tt}$  and  $\rho_{tc}$  can drive baryogenesis, but searches at the HL-LHC may still go empty-handed if the couplings are relatively weak. Taking  $m_{H^\pm} \simeq m_H \simeq m_A \sim 200$  GeV, with  $\rho_{tt}$ ,  $\rho_{tc} \sim 0.1$  and no  $h(125)$ - $H$  mixing,  $H^+ \rightarrow c\bar{b}$  decay is dominant, and the  $c\bar{c}b$  final state is likely overwhelmed by QCD background at the LHC. We show that the electroweak production of  $H^+H^-$  at the ILC is discoverable with integrated luminosity of  $1 \text{ ab}^{-1}$ . We show further that  $m_{H^\pm}$  can be extracted by requiring the two pairs of  $b$  and light jets be roughly equal in mass, without assuming the mass value. Thus, ILC can probe low mass Higgs bosons in multijet final states to complement HL-LHC in the future.

**Author:** HOU, George (National Taiwan University (TW))

**Presenter:** HOU, George (National Taiwan University (TW))

**Session Classification:** Collider