


# Search for exotic decays of the Higgs boson into long-lived particles with jet pairs in the final state at CLIC

---

**Mateusz Goncerz**  
Marcin Kucharczyk

# motivation

- additional long-lived particles emerge in many BSM models
- Hidden Valley (HV) framework introduces them via *hidden* gauge sector
  - motivated by, among others, String Theory
  - communication either by overcoming the barrier in high-energy collisions or via massive messenger particles (ex.  $Z'$ , Higgs)
  - decays of HV particles produce displaced vertices (DV)
  - ideal search for detectors like CLIC



- JHEP 03 (2023) 131
- focus on sensitivity of *CLIC\_ILD* detector model to  $H \rightarrow \pi_V^0 \pi_V^0 \rightarrow b\bar{b}b\bar{b}$  decays
  - $m_H = 126$  GeV
  - $m_{\pi_V^0} \in (25, 35, 50)$  GeV,  $\tau_{\pi_V^0} \in (1, 10, 100, 300)$  ps
  - $BR(\pi_V^0 \rightarrow b\bar{b}) = 100\%$
- dominant production mechanism assumed
  - $e^+e^- \rightarrow HZ(\rightarrow q\bar{q})$ ,  $\sigma = 0.93$  pb at  $\sqrt{s} = 350$  GeV
  - $e^+e^- \rightarrow H\nu_e\bar{\nu}_e$ ,  $\sigma = 0.42$  pb at  $\sqrt{s} = 3$  TeV
- assumed integrated luminosities
  - $1 \text{ ab}^{-1}$  at  $\sqrt{s} = 350$  GeV
  - $3 \text{ ab}^{-1}$  at  $\sqrt{s} = 3$  TeV
- custom seeding algorithm to improve signal-background separation combined with multivariate analysis

$m_{\pi^0}$ [GeV]	$\tau_{\pi^0}$ [ps]	$\sqrt{s} = 350$ GeV		$\sqrt{s} = 3$ TeV	
		$\sigma$ [pb]	sample size	$\sigma$ [pb]	sample size
25	1	0.93	$\sim 240$ K	0.42	$\sim 200$ K
25	10	0.93	$\sim 240$ K	0.42	$\sim 200$ K
25	100	0.93	$\sim 240$ K	0.42	$\sim 200$ K
25	300	0.93	$\sim 240$ K	0.42	$\sim 200$ K
35	1	0.93	$\sim 240$ K	0.42	$\sim 200$ K
35	10	0.93	$\sim 240$ K	0.42	$\sim 200$ K
35	100	0.93	$\sim 240$ K	0.42	$\sim 200$ K
35	300	0.93	$\sim 240$ K	0.42	$\sim 200$ K
50	1	0.93	$\sim 240$ K	0.42	$\sim 200$ K
50	10	0.93	$\sim 240$ K	0.42	$\sim 200$ K
50	100	0.93	$\sim 240$ K	0.42	$\sim 200$ K
50	300	0.93	$\sim 240$ K	0.42	$\sim 200$ K



- beam induced  $\gamma\gamma \rightarrow$  hadrons overlaid for each event

background process	$\sqrt{s} = 350 \text{ GeV}$		$\sqrt{s} = 3 \text{ TeV}$	
	$\sigma[\text{pb}]$	sample size	$\sigma[\text{pb}]$	sample size
$q\bar{q}$	24.41	$\sim 2\text{M}$	2.95	$\sim 200\text{K}$
$q\bar{q}\nu\bar{\nu}$	0.32	$\sim 306\text{K}$	1.32	$\sim 200\text{K}$
$q\bar{q}q\bar{q}$	5.85	$\sim 1.44\text{M}$	0.55	$\sim 750\text{K}$
$q\bar{q}q\bar{q}\nu\bar{\nu}$		-	0.07	$\sim 300\text{K}$
$t\bar{t}$	0.45	$\sim 241\text{K}$		-
WWZ	0.01	$\sim 40\text{K}$		-

- more than 4 jets omitted due to low cross-section
- beam induced  $\gamma\gamma \rightarrow$  hadrons overlaid for each event

# reconstruction of DVs

- vertices of  $\pi_V^0$  decays are displaced from the beam axis





$m_{\pi_V^0} = 50$  GeV,  $\tau_{\pi_V^0} = 1$  (red), 10 (blue), 100 (green), 300 (yellow) ps

- LCFI+ (Linear Collider Flavour Identification) algorithms for SV reconstruction found to be inefficient
  - designed primarily for B and D hadron decays
  - not enough charged tracks assigned to displaced vertices
  - hindering signal-background separation, especially in 1 ps samples
  - not possible to solve using available settings

# reconstruction of DVs – modified approach

- tracks and jets reconstructed using standard methods
  - 6 for  $\sqrt{s} = 350$  GeV and 4 for  $\sqrt{s} = 3$  TeV
  - longitudinally invariant  $k_t$  (FastJet), tagging based on BDT
  - optimized for HV particles (jet radius etc.)
  - see CLICdp-Note-2018-001 for details
- manual seeding to find  $\pi_V^0 \rightarrow b\bar{b}$  vertex candidates
  - good quality charged tracks not coming from PV
    - $IP/\sigma_{IP} > 16$
  - find base track with at least 4 tracks within 1 mm
  - find point of closest approach of each pair
  - assign additional tracks within 1 mm
  - look for new base track and repeat until no tracks left
- perform a standard SV reconstruction based on the impact parameter wrt. the seed positions determined in previous step



$$m_{\pi_V^0} = 35 \text{ GeV}, \tau_{\pi_V^0} = 10 \text{ ps}$$

- number of charged tracks assigned to SVs using standard algorithms (left) and our modified approach (right)
- clear signal-background separation introduced

- to match signal signature, events are required to have
  - at least two DVs
  - at least 4 jets, b-tag probability  $\geq 0.95$
- jets matched with vertices in a way that maximizes number of common charged tracks
- assigned jets used to reconstruct di-jets ( $\pi_\nu^0$ ) and four-jet (Higgs)
- at  $\sqrt{s} = 350$  GeV, the two remaining jets are used to reconstruct Z boson candidate



# signal-background separation

- pre-selection requirements very effective at suppressing backgrounds

		$\sqrt{s} = 350 \text{ GeV}$	$\sqrt{s} = 3 \text{ TeV}$
$m_{\pi_V^0} [\text{GeV}]$	$\tau_{\pi_V^0} [\text{ps}]$	Eff. <i>preSEL.</i> [%]	Eff. <i>preSEL.</i> [%]
25	1	78	68
25	10	94	86
25	100	99	93
25	300	97	80
35	1	76	70
35	10	93	86
35	100	99	94
35	300	98	82
50	1	72	72
50	10	89	89
50	100	99	90
50	300	99	86



		$\sqrt{s} = 350 \text{ GeV}$	$\sqrt{s} = 3 \text{ TeV}$
background process		Eff. <i>preSEL.</i> [%]	Eff. <i>preSEL.</i> [%]
$q\bar{q}$		12	6
$q\bar{q}\nu\bar{\nu}$		12	8
$q\bar{q}q\bar{q}$		8	9
$q\bar{q}q\bar{q}\nu\bar{\nu}$		-	11
$t\bar{t}$		12	-
WWZ		14	-

- further separation achieved via multivariate analysis
- decision tree (BDTG) trained using physical variables with good separation:
  - DV, di-jet and four-jet mass
  - DV and their assigned tracks multiplicity
  - distance at which the transitions from 3 to 2-jet and from 4 to 3-jet event takes place
    - effective against backgrounds with different jet number
  - Z candidate mass (at  $\sqrt{s} = 350$  GeV)



$m_{\pi^0} = 35$  GeV,  $\tau_{\pi^0} = 10$  ps

- each signal sample treated independently
- backgrounds combined with weights based on cross-section and pre-selection efficiency




$$m_{\pi_V^0} = 35 \text{ GeV}, \tau_{\pi_V^0} = 10 \text{ ps}$$


- number of events normalized to assumed total luminosity

# sensitivity

- threshold value chosen to maximize sensitivity  $S/\sqrt{S+B}$





$$m_{\pi^0} = 35 \text{ GeV}$$



- sensitivity roughly 4-6 times larger at  $\sqrt{s} = 3 \text{ TeV}$ 
  - bear in mind different luminosities
  - consistent across all  $m_{\pi^0}$  masses



# upper limits

- derived using CL(s) at 95% CL
- assuming absence of signal observation



- slightly higher at  $\sqrt{s} = 350 \text{ GeV}$ , but same order of magnitude
- much better than existing experiments (orders of  $10^0 - 10^2$ )

- normalized to Higgs production cross-section



- an order of magnitude higher at  $\sqrt{s} = 350 \text{ GeV}$

- sensitivity of CLIC\_ILD detector to long-lived HV particles studied for the first ( $\sqrt{s} = 350$  GeV) and last ( $\sqrt{s} = 3$  TeV) stage of CLIC operation
- the standard algorithms are not efficient enough in assigning charged tracks to vertices displaced from the beam axis
  - not possible to tweak with settings available at the time
- modified approach proposed to mitigate these problems
  - good signal-background separation achievable
- resulting upper limits are many orders of magnitude better than achievable in current experiments