

Experimental Physics at Hadron Colliders

CERN Summer Students Lectures, July 17-21, 2023 - Lecture 1/4

Markus KLUTE (markus.klute@kit.edu)
Institute of Experimental Particle Physics (ETP)



CERN Summer Student Lecture Program 2023

- Particle World - David Tong
- Detectors - Wernes Ziegler
- From Raw Data to Physics Results - Paul Laycock
- Accelerator Technology Part I - Susana Bermudez
- The Standard Model - Christophe Grojean
- Foundation of Statistics - Glen Cowan
- Particle Accelerators - Foteini Avesta
- Nuclear Physics at CERN - Magdalena Kowalska
- Theoretical Concepts in Particle Physics - Tim Cohen
- Future High Energy Collider Projects - Barbara Dalena
- Cosmology - Valerie Domcke
- **Heavy Ion Physics** - Francesca Bellini
- Accelerator Technology Part 2 - Walter Delsolaro
- **Experimental Physics at Hadron Colliders - Markus Klute**
- **Flavour Physics** - Mark Williams
- Physics and Medical Applications - Manuela Cirilli
- Accelerator Technology Part 3 - Francesc Pujol
- Astroparticle Physics - Bradley Kavanagh
- Predictions at Hadron Colliders - Alexander Huss
- Lepton Colliders - Frank Simon
- Antimatter in the Laboratory - Jack Devlin
- Physics Beyond the Standard Model - Tevong You
- Electronics, DAQ and Trigger - Tommaso Colombo
- What is String Theory? - Timo Weigand



<https://indico.cern.ch/event/1254879/timetable/>

Learning Objectives

- Understanding basics concepts of experimental particle physics at hadron colliders
- Knowledge of the broad and diverse LHC Physics program, including the vast number of opportunities at the LHC



What will I learn ?



How do I learn?


Particle Physics at Colliders

- The objective of particle physics is to uncover the fundamental laws of nature and gain a comprehensive understanding of the universe at its most fundamental level.

- High-energy particle collisions enable

- Discovery of new and massive particles
- Probing the structure of matter
- Exploring fundamental forces of nature
- Recreating the early universe

Standard Model of Particle Physics



- **Lecture 1: Introduction, fundamentals, cross sections**
- Lecture 2: Standard model measurements
- Lecture 3: Higgs physics
- Lecture 4: Searches for new physics

Following the Protons @ CERN

<https://videos.cern.ch/record/2020780>


<https://home.cern/science/accelerators/accelerator-complex>



Following the Protons @ CERN

<https://videos.cern.ch/record/2020780>


<https://home.cern/science/accelerators/accelerator-complex>



Following the Protons @ CERN

- Linear accelerator 4 (Linac 4) accelerates negative hydrogen ions (H^-) to 160 MeV


<https://videos.cern.ch/record/2020780>
<https://home.cern/science/accelerators/accelerator-complex>



Following the Protons @ CERN

- Linear accelerator 4 (Linac 4) accelerates negative hydrogen ions (H^-) to 160 MeV
- Ions are stripped of their two electrons during injection into the Proton Synchrotron Booster
- Booster is using four superimposed synchrotron rings to accelerate to 2 GeV

<https://videos.cern.ch/record/2020780>
<https://home.cern/science/accelerators/accelerator-complex>




Following the Protons @ CERN

- Linear accelerator 4 (Linac 4) accelerates negative hydrogen ions (H-) to 160 MeV
- Ions are stripped of their two electrons during injection into the Proton Synchrotron Booster
- Booster is using four superimposed synchrotron rings to accelerate to 2 GeV
- Proton Synchrotron (PS) uses conventional electromagnets and accelerates to 26 GeV

<https://videos.cern.ch/record/2020780>


<https://home.cern/science/accelerators/accelerator-complex>



Following the Protons @ CERN

- Super Proton Synchrotron (SPS) uses conventional electromagnets and accelerates protons to 450 GeV
- Discovery of the W and Z boson in 1983.



<https://videos.cern.ch/record/2020780>
<https://home.cern/science/accelerators/accelerator-complex>



Following the Protons @ CERN

<https://videos.cern.ch/record/2020780>



<https://home.cern/science/accelerators/accelerator-complex>



Following the Protons @ CERN

- LHC: 8 arcs and 8 straight sections
- Proton-proton collisions at 13.6 TeV, also pA and AA collisions
- ARCs (2.45km)
 - 23 arc 'cells' with main dipoles, quadrupoles and other multipoles
- Straight sections (528m)
 - Experiments, beam injection, RF acceleration, beam dump

<https://videos.cern.ch/record/2020780>
<https://home.cern/science/accelerators/accelerator-complex>



Following the Protons @ CERN

CMS DETECTOR

Total weight : 14,000 tonnes
 Overall diameter : 15.0 m
 Overall length : 28.7 m
 Magnetic field : 3.8 T

STEEL RETURN YOKE
 12,500 tonnes

SILICON TRACKERS
 Pixel (100x150 μm) $\sim 1\text{m}^2 \sim 66\text{M}$ channels
 Microstrips (80x180 μm) $\sim 200\text{m}^2 \sim 9.6\text{M}$ channels

SUPERCONDUCTING SOLENOID
 Niobium titanium coil carrying $\sim 18,000\text{A}$



MUON CHAMBERS
 Barrel: 250 Drift Tube, 480 Resistive Plate Chambers
 Endcaps: 540 Cathode Strip, 576 Resistive Plate Chambers

PRESHOWER
 Silicon strips $\sim 16\text{m}^2 \sim 137,000$ channels

FORWARD CALORIMETER
 Steel + Quartz fibres $\sim 2,000$ Channels



CRYSTAL ELECTROMAGNETIC CALORIMETER (ECAL)
 $\sim 76,000$ scintillating PbWO_4 crystals

HADRON CALORIMETER (HCAL)
 Brass + Plastic scintillator $\sim 7,000$ channels



<https://videos.cern.ch/record/2020780>

<https://home.cern/science/accelerators/accelerator-complex>




Tile calorimeters
 LAr hadronic end-cap and forward calorimeters
 Pixel detector
 LAr electromagnetic calorimeters
 Transition radiation tracker
 Semiconductor tracker
 Solenoid magnet
 Toroid magnets
 Muon chambers

Following the Protons @ CERN

<https://videos.cern.ch/record/2020780>


<https://home.cern/science/accelerators/accelerator-complex>



Following the Protons @ CERN

<https://videos.cern.ch/record/2020780>

<https://home.cern/science/accelerators/accelerator-complex>



Why Protons and not Electrons?


- Energy loss by Bremsstrahlung ultimately limits the center-of-mass energy of circular lepton colliders
 - LEP reached 209 GeV with an energy loss per of 3.5 GeV
 - Energy loss per turn for the LHC is 7 keV
- Hadron collider limitation is the bending power
 - LHC effective radius is 2.7 km requiring a field of 8.5 T
 - At LEP (209 GeV) only about 0.1T was required.

Energy loss: $\Delta E \propto \left(\frac{E}{m}\right)^4 \frac{1}{R}$

$$\frac{\Delta E(e)}{\Delta E(p)} = \frac{P(e)}{P(p)} = \left(\frac{m_p}{m_e}\right)^4 \simeq 10^{13}$$





LHC DIPOLE CROSS SECTION



CERN AC/DI/MM — 06-2001

Why not Protons on Anti-Protons?

- Producing an anti-protons beam is challenging
- Studying collisions of proton constituents, i.e. quarks and gluons
- The probability to find a Parton with momentum fraction x is captured in parton distribution functions (PDFs)
 - PDFs are measured in experiments
 - Q^2 evolution is calculated with Altarelli-Parisi equations
 - Example: top pair production



Momentum sum rule
$$\sum_i \int_0^1 dx x f_i(x, Q^2) = 1$$

Flavour conservation sum rules

$$\int_0^1 (f_u(x, Q^2) - f_{\bar{u}}(x, Q^2)) dx = 2$$

$$\int_0^1 (f_d(x, Q^2) - f_{\bar{d}}(x, Q^2)) dx = 1$$


$$\int_0^1 (f_s(x, Q^2) - f_{\bar{s}}(x, Q^2)) dx = 0$$

Hadron Collider History



Collider	Location	Operating	Dimensions	Beam Energy	Type	Luminosity	Legacy
ISR	CERN	1971-84	~1km	31 GeV	pp	$2 \cdot 10^{31} \text{cm}^{-2} \text{s}^{-1}$	First of its kind
SPS	CERN	1981-93	~6.9km	450 GeV	p-anti-p	$5 \cdot 10^{30} \text{cm}^{-2} \text{s}^{-1}$	W and Z boson
HERA	DESY	1992-2007	~6.3km	920 GeV	ep	$5 \cdot 10^{31} \text{cm}^{-2} \text{s}^{-1}$	PDFs
Tevatron	FERMILAB	1992-2011	~6.2km	980 GeV	p-anti-p	$4 \cdot 10^{32} \text{cm}^{-2} \text{s}^{-1}$	Top quark discovery
SSC	Texas	Never completed	~87km	20 TeV	pp	$10^{33} \text{cm}^{-2} \text{s}^{-1}$	-
RHIC	Brookhaven	Since 2000	~3.8km	100 GeV	pp, pA, AA	$10^{32} \text{cm}^{-2} \text{s}^{-1}$	Quark-gluon plasma
LHC	CERN	Since 2008	LEP tunnel ~27km	6.8 TeV	pp, pA, AA	$2 \cdot 10^{34} \text{cm}^{-2} \text{s}^{-1}$	Higgs boson discovery
EIC	Brookhaven	~2032	RHIC infrastructure	275 GeV	ep, eA	$10^{34} \text{cm}^{-2} \text{s}^{-1}$	-
FCC-hh	CERN	~2060	FCC-ee tunnel	50 TeV	pp, pA, AA	$3 \cdot 10^{35} \text{cm}^{-2} \text{s}^{-1}$	-

Hadron-Hadron Collision

Broadband beam of various parton types with various energies




Proton



Proton

Challenge: reliable calculations of observables

Hadron-Hadron Collision




$R = 0.8\text{fm}$

Prototype Process: Drell-Yan Production

$$pp \rightarrow l^+ l^- + X$$


- Hadron production of lepton pairs
- Factorising “hard” and “soft” components
 - Calculate hard partonic subprocess
 - Weight cross section with probability to find partons with momenta x_1 and x_2
 - Integrate over all possible parton momenta
 - Sum over all possible parton flavors




$$\sigma_{\text{DY}} = \sum_{i,j} \int dx_i dx_j f_i(x_i) f_j(x_j) \cdot \hat{\sigma}(q_i q_j \rightarrow l^+ l^-)$$

Prototype Process: Drell-Yan Production

$$pp \rightarrow l^+ l^- + X$$



higher orders



Where does this belong to?

The PDF?

Or the parton process?

- Factorisation scale μ_f
- Attribution ambiguous


$$\hat{\sigma}_{ij}(x_i, x_j, \mu_r^2, \alpha_s(\mu_r^2)) \rightarrow \hat{\sigma}_{ij}(x_i, x_j, \mu_f^2, \mu_r^2, \alpha_s(\mu_r^2))$$

$$f_i(x_i) \rightarrow f_i(x_i, \mu_f^2)$$

- Leads to soft and / or collinear divergences (long-distance effects!)
- Solution: introduce a new scale to separate short- and long-distance effects
- Effects are absorbed into PDFs and determined experimentally

Hadron-Hadron Cross Section

- Factorisation valid for more general final states, e.g. jet production



Luminosity

$$\frac{dN}{dt} = \mathcal{L}(t)\sigma$$

Events per second for a given cross section


Cross section in barn (10^{-24} cm^2)
Luminosity in $\text{cm}^{-2}\text{s}^{-1}$

$$N = L\sigma \quad \text{with} \quad L = \int \mathcal{L}(t)dt$$

Luminosity depends on

- Number of protons per bunch
- Beam size at interaction point
- Number of bunches


$$\mathcal{L} = \frac{N_p^2 f N_b}{4\pi \sigma_x \sigma_y}$$



Luminosity measurements

- Crucial for absolute cross section measurements or to set limits on physics beyond the SM
- Calibration performed using “Van der Meer scans”
- Uncertainty of order 1-2%

Luminosity monitoring





Luminosity

$$L = \frac{N_p^2 f N_b}{4\pi \sigma_x \sigma_y} = \frac{N_p^2 f N_b \gamma^4}{4\pi \beta^* \epsilon_N}$$

Luminosity (beam size) expressed using beam parameters


Parameter	2010	2011	2012	2016	2017	2018	Nominal	HL-LHC
CoM Energy	7 TeV	7 TeV	8 TeV	13 TeV	13 TeV	13 TeV	14 TeV	14 TeV
N_p	$1.1 \cdot 10^{11}$	$1.4 \cdot 10^{11}$	$1.6 \cdot 10^{11}$	$1.2 \cdot 10^{11}$	$1.2 \cdot 10^{11}$	$1.2 \cdot 10^{11}$	$1.15 \cdot 10^{11}$	$2.2 \cdot 10^{11}$
Bunches k	368	1380	1380	2300	2450	2500	2808	2760
Spacing	150 ns	50 ns	50 ns	25 ns	25 ns	25 ns	25 ns	25ns
ϵ (mm rad)	2.4-4	1.9-2.3	2.5	2.6	2.3	2.6	3.75	2.5
β^* (m)	3.5	1.5-1	0.6	0.4	0.3-0.4	0.4	0.55	0.15
L ($\text{cm}^{-2}\text{s}^{-1}$)	$2 \cdot 10^{32}$	$3.3 \cdot 10^{33}$	$\sim 7 \cdot 10^{33}$	$1.5 \cdot 10^{33}$	$2.0 \cdot 10^{34}$	$2 \cdot 10^{34}$	10^{34}	$8 \cdot 10^{34}$
PU	~ 2	~ 10	~ 30	~ 30	~ 50	~ 50	~ 25	~ 130

Integrated Luminosity and Pile-Up



Event rates at the LHC

- Total cross sections
 - $\sim 1.6 \cdot 10^9$ /s (80mb, $2 \cdot 10^{34} \text{cm}^{-2} \text{s}^{-1}$)
 - Bunch crossing rate of 40MHz
- Jets ($E_T^{\text{jet}} > 100$ GeV)
 - ~ 40000 Hz
- W & Z bosons
 - ~ 4000 Hz, ~ 1000 Hz
- Top Quarks
 - ~ 20 Hz
- Jets ($E_T^{\text{jet}} > 650$ GeV)
 - ~ 6 Hz
- Higgs bosons
 - ~ 1 Hz (50pb , $2 \cdot 10^{34} \text{cm}^{-2} \text{s}^{-1}$)



$$\sigma = \frac{N_{\text{sel}} - N_{\text{bkg}}}{\int L dt \cdot A \cdot \epsilon}$$

- Luminosity
- Selection efficiency (objects, kinematics, binning)
- Acceptance (extrapolation from fiducial volume)
- Background estimation

- **Lecture 1: Introduction, fundamentals, cross sections**
- Lecture 2: Standard model measurements
- Lecture 3: Higgs physics
- Lecture 4: Searches for new physics

Quiz

- Why don't we collide electrons and positrons or protons and anti-protons at the LHC?
- What is the minimal energy of a proton hitting a proton target at rest to produce an anti-proton?
- How many proton-proton collisions does the LHC produce per second?
- How can we estimate the expected number of pile-up events?
- How many Higgs Bosons have been produced in Run 2?
- If the hunt for the Higgs Boson can be compared to the search of a needle in the haystack, how big is the haystack?



References and further reading

■ Textbooks

- Modern Particle Physics by Mark Thomson
- QCD at Colliders by Ellis, Stirling, and Weber

■ Pictures

- CERN Document Server
- Wikipedia
- Or reference on page

■ References

- Previous CERN Summer Lectures - <https://indico.cern.ch/category/97/>
- MIT's OCW 8.701 and 8.811
- KIT's Particle Physics master courses (you can contact me)
- Or reference on page