

FAIROS-HEP + AGC

Analysis Preservation & Reinterpretation

Kyle Stuart Cranmer (Wisconsin)
Mike Hildreth (Notre Dame)
Peter Elmer (Princeton)



FAIR and Open Science in High Energy Physics
OAC-2226378, OAC-2226379 and OAC-2226380



FAIROS-HEP

Recently, the US National Science Foundation funded a new Research Coordination Network project titled “FAIROS-HEP”.

Findable

Accessible

Interoperable

Reusable

Open **S**cience

The NSF's FAIROS Research Coordination Networks

Findable, Accessible, Interoperable, Reusable, Open Science Research Coordination Networks (FAIROS RCN) program represent a pooled investment of over \$12.5 million in open science from all directorates comprising NSF.

FAIROS RCN supports groups of investigators to communicate, innovate, coordinate, and standardize research practices, training, and educational activities across disciplinary, organizational, geographic, and international boundaries to achieve the goals of FAIR and other open-science guiding principles.

Research coordination networks are a form of awards that NSF makes to advance scientific practices and standards broadly across multiple research fields. These RCN awards will be for **three-year projects**.

FAIROS-HEP Continues a Legacy of Contributions

DASPOS (2012-2016)

- <https://daspos.crc.nd.edu/>
- Contributions to RECAST led to REANA as a spinoff project now led by CERN
- Supported REANA Common Workflow Language

DIANA-HEP (2015-2021)

- <https://diana-hep.org/>
- Contributions to REANA, RECAST, launched pyhf likelihood publishing, Active Learning for reinterpretation
- Supported GitHub -> Zenodo DOI minting

IRIS-HEP (2018-?)

- <https://iris-hep.org/>
- Major contributions to likelihood publishing, HEPData integration,

SCAILFIN (2018-2021)

- <https://scaifin.github.io/>
- Contributions to REANA (Slurm and HPC backends, applications built on top of REANA, etc.), Active Learning for reinterpretation

FAIROS-HEP (2022-2025)

- <https://fairos-hep.org/> (under construction)
- Continue the legacy of contributions, help coordinate the ecosystem

What is FAIROS-HEP?

The FAIROS-HEP project aims to connect groups of researchers thinking about FAIR data in HEP and other experts in this field to envision a more cohesive infrastructure around data and publications in HEP.

- By focusing on FAIR data practices and **how data and software can be linked to physics results**, we hope to **build a network of researchers** thinking about how we can create a “**living publication**” to preserve and extend physics results.
- The project includes **some funding for building infrastructure** as well as **future workshops connecting groups**.

The NSF's FAIROS Research Coordination Networks

We aim to leverage the CNI to broaden HEP's interaction with Libraries and IT groups in the US. Essentially all HEP universities in the US are members of CNI.



The Coalition for Networked Information (CNI) is dedicated to supporting the transformative promise of digital information technology for the advancement of scholarly communication and the enrichment of intellectual productivity. Over 200 institutions representing higher education, publishing, information technology, scholarly and professional organizations, foundations, and libraries and library organizations make up CNI's members.

Semi-annual membership meetings bring together representatives of CNI's constituencies to discuss ongoing and new projects and to plan for future initiatives.

ARL Applauds NSF Open Science Investment

Jessica Aiwuyor | 202-296-2296 | jaiwuyor@arl.org | August 15, 2022

Share

Tweet

Email



image © iStock.com/kentoh

The Association of Research Libraries (ARL) commends the ongoing commitment of the US National Science Foundation (NSF) to open science. NSF today announced awards for 10 new projects focused on building and enhancing coordination among researchers and other stakeholders to advance FAIR (findable, accessible, interoperable, reusable) data principles and open-science practices.


The inaugural awards in NSF's Findable, Accessible, Interoperable, Reusable, Open Science Research Coordination Networks (FAIROS RCN) program represent a pooled investment of over \$12.5 million in open science from all directorates comprising NSF. This program is particularly unique given that the 10 projects are composed of 28 distinct NSF awards (detailed below) representing many organizations and institutions in the United States seeking to advance open-science efforts.

More info here: <https://www.arl.org/news/arl-applauds-nsf-open-science-investment/>

Kickoff workshop

The goals of the workshop were to:

1. Assess progress by each experiment in producing reusable data,
2. Establish updated ideas regarding the use cases for data access, interoperability, and reuse across the different experiments and experimental fields,
3. Define what data and associated information supports the use cases, and
4. Identify a preliminary set of access methods and infrastructure that would support these use cases.



FAIROS-HEP Kick-Off Workshop

8-10 Feb 2023
CERN
Europe/Zurich timezone

Enter your search term

Overview

Scientific Programme

Timetable

Contribution List

My Conference

My Contributions

Recently, the US National Science Foundation funded a new Research Coordination Network project titled "FAIROS-HEP". The project aims to connect groups of researchers thinking about FAIR data in HEP and other experts in this field to envision a more cohesive infrastructure around data and publications in HEP. By focusing on FAIR data practices and how data and software can be linked to physics results, we hope to build a network of researchers thinking about how we can create a "living publication" to preserve and extend physics results. The project includes some funding for building infrastructure as well as future workshops connecting groups.

This is our kick-off workshop, located at CERN.

The goals of the workshop will be to:

1. Assess progress by each experiment in producing reusable data,
2. Establish updated ideas regarding the use cases for data access, interoperability, and reuse across the different experiments and experimental fields,
3. Define what data and associated information supports the use cases, and
4. Identify a preliminary set of access methods and infrastructure that would support these use cases.

We look forward to engaging with you in this project.

<https://indico.cern.ch/event/1234612/>

Technical Recommendations: Provide initial direction for which elements of the cyber ecosystem will be most relevant for the first round of technical improvements. Initiate investigations.

How can academic publishing support FAIR and open science?

01 Feb 2023 Sponsored by [Machine Learning: Science and Technology](#)


Join us for a live webinar at 1 p.m. GMT on 10 February 2023, sponsored by the IOP Publishing journal, ***Machine Learning: Science and Technology***. Hear from data science and physical science experts as they share experiences and explore solutions to common barriers in FAIR and open science

- **Keith Butler** (Chair), Queen Mary University of London, Programme Director - Sustainable Engineering
- **Kyle Cranmer**, Director of the University of Wisconsin-Madison Data Institute & MLST Editor in Chief
- **Sünje Dallmeier-Tiessen**, Data coordinator in the Scientific Information Service, CERN
- **James Warren**, NIST Materials Genome
- **Megan Ansdell**, Program Officer, Planetary Science Division at NASA
- **Harry Enke**, head of Section EScience & SuperComputing at Leibniz Institute for Astrophysics Potsdam (AIP) & PUNCH4NFDI
- **Daniel Keirs**, Head of Journal Strategy and Performance, IOP Publishing



IRIS-HEP

The IRIS-HEP Analysis Systems focus area extends to reuse and preservation, but it has not been a major area of activity. FAIROS-HEP will coordinate closely.



Some Recent Developments in HEP

Open Science & Beyond

The field is at a tipping point. CERN has publicly embraced Open Science and the experiments are adopting new policies.

But we also realize Open Data is not the end of the story.
Reuse is key!

nature
physics

PERSPECTIVE

<https://doi.org/10.1038/s41567-018-0342-2>

Corrected: Publisher Correction

OPEN

Open is not enough

Xiaoli Chen^{1,2}, Sünje Dallmeier-Tiessen^{1*}, Robin Dasler^{1,11}, Sebastian Feger^{1,3}, Pamfilos Fokianos¹, Jose Benito Gonzalez¹, Harri Hirvonsalo^{1,4,12}, Dinos Kousidis¹, Artemis Lavasa¹, Salvatore Mele¹, Diego Rodriguez Rodriguez¹, Tibor Šimko^{1*}, Tim Smith¹, Ana Trisovic^{1,5*}, Anna Trzcinska¹, Ioannis Tsanaksidis¹, Markus Zimmermann¹, Kyle Cranmer⁶, Lukas Heinrich⁶, Gordon Watts⁷, Michael Hildreth⁸, Lara Lloret Iglesias⁹, Kati Lassila-Perini⁴ and Sebastian Neubert¹⁰

The solutions adopted by the high-energy physics community to foster reproducible research are examples of best practices that could be embraced more widely. This first experience suggests that reproducibility requires going beyond openness.



Highlight: 2020 CERN homepage

HEPData
@HEPData

Thanks @KyleCranmer for your support and promotion of @HEPData over several years. Looking forward to future collaboration with @iris_hep on #pyhf likelihoods and more.

Kyle Cranmer @KyleCranmer · Jan 29

I would like to applaud @STFC_Matters for funding @HEPData, a vital piece of cyberinfrastructure for HEP. The @NSF has been supporting HEP software and cyberinfrastructure with DASPOS, @diana_hep and @iris_hep. @iris_hep looks forward to collaborating with you! twitter.com/HEPData/status...

1:15 PM · Jan 30, 2020 · Twitter Web App

The screenshot shows the CERN homepage with a navigation bar at the top containing 'ABOUT', 'NEWS', 'SCIENCE', 'RESOURCES', 'SEARCH', and 'EN'. The main banner features a large image of the ATLAS detector with a text box on the left that reads: 'CERN Council appoints Fabiola Gianotti for second term of office as CERN Director General' and 'Press release | 6 November, 2019'. Below the banner is a 'LATEST NEWS' section with three featured articles: 'New open release allows theorists to explore ...' (Knowledge sharing | News | 9 January, 2020), 'Relive 2019 at CERN' (Knowledge sharing | News | 30 December, 2019), and 'Dive into the world of accelerators' (Accelerators | News | 19 December, 2019). A vertical sidebar on the right lists 'Events' and 'Key Achievements'.



The screenshot shows a CERN news article page. The header includes 'CERN Accelerating science' and navigation links for 'ABOUT' and 'NEWS'. The article title is 'New open release allows theorists to explore LHC data in a new way', with a subtitle 'The ATLAS collaboration releases full analysis likelihoods, a first for an LHC experiment'. The byline is '9 JANUARY, 2020 | By Katarina Anthony'. The main image shows a laptop displaying a data visualization next to a printed document titled 'ATLAS 2019 Data Release Summary'. A small 'menu' icon is visible in the bottom left corner of the article page.





News › News › Topic: Knowledge sharing

Voir en [français](#)

New open release allows theorists to explore LHC data in a new way

The ATLAS collaboration releases full analysis likelihoods, a first for an LHC experiment

9 JANUARY, 2020 | By Katarina Anthony



Courtesy of CERN

ATLAS releases 'full orchestra' of analysis instruments

01/14/21 | By Stephanie Melchor

The ATLAS collaboration has begun to publish likelihood functions, information that will allow researchers to better understand and use their experiment's data in future analyses.

Published Probability Models

Published Probability Models

Updated list of HEPData entries for publications using `histFactory` JSON statistical models:

- Search for charginos and neutralinos in final states with two boosted hadronically decaying bosons and missing transverse momentum in pp collisions at $\sqrt{s}=13$ TeV with the ATLAS detector. 2021. doi:10.17182/hepdata.104458
- Measurement of the $t\bar{t}$ production cross section in pp collisions at $\sqrt{s}=13$ TeV with the ATLAS detector. 2021. doi:10.17182/hepdata.105039
- Search for R-parity violating supersymmetry in a final state containing leptons and many jets with the ATLAS experiment using $\sqrt{s}=13$ TeV proton-proton collision data. 2021. doi:10.17182/hepdata.104860
- Search for chargino-neutralino pair production in final states with three leptons and missing transverse momentum in $\sqrt{s}=13$ TeV pp collisions with the ATLAS detector. 2021. doi:10.17182/hepdata.95751
- Measurements of the inclusive and differential production cross sections of a top-quark-antiquark pair in association with a Z boson at $\sqrt{s}=13$ TeV with the ATLAS detector. 2021. doi:10.17182/hepdata.100351
- Search for pair production of third-generation scalar leptoquarks decaying into a top quark and a τ -lepton in pp collisions at $\sqrt{s}=13$ TeV with the ATLAS detector. 2021. doi:10.17182/hepdata.100174.
- Search for squarks and gluinos in final states with one isolated lepton, jets, and missing transverse momentum at $\sqrt{s}=13$ TeV with the ATLAS detector. 2021. doi:10.17182/hepdata.97041
- Search for trilepton resonances from chargino and neutralino pair production in $\sqrt{s}=13$ TeV pp collisions with the ATLAS detector. 2020. doi:10.17182/hepdata.99806.
- Search for displaced leptons in $\sqrt{s}=13$ TeV pp collisions with the ATLAS detector. 2020. doi:10.17182/hepdata.98796.
- Search for squarks and gluinos in final states with jets and missing transverse momentum using 139 fb⁻¹ of $\sqrt{s}=13$ TeV pp collision data with the ATLAS detector. 2020. doi:10.17182/hepdata.95664.
- Measurement of the $t\bar{t}$ production cross-section in the lepton+jets channel at $\sqrt{s}=13$ TeV with the ATLAS experiment. 2020. doi:10.17182/hepdata.95748.
- Search for long-lived, massive particles in events with a displaced vertex and a muon with large impact parameter in pp collisions at $\sqrt{s}=13$ TeV with the ATLAS detector. 2020. doi:10.17182/hepdata.91760
- Search for chargino-neutralino production with mass splittings near the electroweak scale in three-lepton final states in $\sqrt{s}=13$ TeV pp collisions with the ATLAS detector. 2019. doi:10.17182/hepdata.91127.
- Searches for electroweak production of supersymmetric particles with compressed mass spectra in $\sqrt{s}=13$ TeV pp collisions with the ATLAS detector. 2019. doi:10.17182/hepdata.91374
- Search for direct stau production in events with two hadronic τ -leptons in $\sqrt{s}=13$ TeV pp collisions with the ATLAS detector. 2019. doi:10.17182/hepdata.92006.
- Search for direct production of electroweakinos in final states with one lepton, missing transverse momentum and a Higgs boson decaying into two b-jets in (pp) collisions at $\sqrt{s}=13$ TeV with the ATLAS detector. 2019. doi:10.17182/hepdata.90607.
- Search for squarks and gluinos in final states with same-sign leptons and jets using 139 fb⁻¹ of data collected with the ATLAS detector. 2019. doi:10.17182/hepdata.91214.
- Search for bottom-squark pair production with the ATLAS detector in final states containing Higgs bosons, b-jets and missing transverse momentum. 2019. doi:10.17182/hepdata.89408.

Search for bottom-squark pair production with the ATLAS detector in final states containing Higgs bosons, b -jets and missing transverse momentum

The ATLAS collaboration

Aad, Georges , Abbott, Brad , Abbott, Dale Charles , Abdinov, Ovsat , Abed Abud, Adam , Abeling, Kira , Abhayasinghe, Deshan Kavishka , Abidi, Syed Haider , Abouzeid, Ossama , Abraham, Nicola

JHEP 12 (2019) 060, 2019.

<https://doi.org/10.17182/hepdata.89408.v3>

Journal

INSPIRE

Resources

 HistFactory

Reuse in <30 lines of code

```
1 import json
2 import cabinetry
3 import pyhf
4 from cabinetry.model_utils import prediction
5 from pyhf.contrib.utils import download
6
7 # download the ATLAS bottom-squarks analysis probability models from HEPData
8 download("https://www.hepdata.net/record/resource/1935437?view=true", "bottom-squarks")
9
10 # construct a workspace from a background-only model and a signal hypothesis
11 bkg_only_workspace = pyhf.Workspace(json.load(open("bottom-squarks/RegionC/BkgOnly.json")))
12 patchset = pyhf.PatchSet(json.load(open("bottom-squarks/RegionC/patchset.json")))
13 workspace = patchset.apply(bkg_only_workspace, "sbottom_600_280_150")
14
15 # construct the probability model and observations
16 model, data = cabinetry.model_utils.model_and_data(workspace)
17
18 # produce visualizations of the pre-fit model and observed data
19 prefit_model = prediction(model)
20 cabinetry.visualize.data_mc(prefit_model, data)
21
22 # fit the model to the observed data
23 fit_results = cabinetry.fit.fit(model, data)
24
25 # produce visualizations of the post-fit model and observed data
26 postfit_model = prediction(model, fit_results=fit_results)
27 cabinetry.visualize.data_mc(postfit_model, data)
```






Figure 3: Pre-fit (left) and post-fit (right) visualizations of a selected signal hypothesis for four signal regions of the ATLAS search [41] of a bottom-squark of mass 600 GeV with a second-lightest neutralino of mass 280 GeV and lightest supersymmetric particle of mass 150 GeV generated from the full statistical models published in Ref. [20] using code from Ref. [40].



A SModelS interface for pyhf likelihoods

#6

Gaël Alguero (LPSC, Grenoble), Sabine Kraml (LPSC, Grenoble), Wolfgang Waltenberger (Vienna, OAW and Vienna U.) (Sep 3, 2020)

Published in: *Comput.Phys.Commun.* 264 (2021) 107909 • e-Print: [2009.01809](https://arxiv.org/abs/2009.01809) [hep-ph]

[pdf](#) [DOI](#) [cite](#) [claim](#)

[reference search](#)

[20 citations](#)

Community Contributions

Reinterpretation and Long-Term Preservation of Data and Code #1

Stephen Bailey (LBL, Berkeley), K.S. Cranmer (Wisconsin U., Madison), Matthew Feickert (Wisconsin U., Madison), Rob Fine (Los Alamos), Sabine Kraml (LPSC, Grenoble) et al. (Sep 16, 2022)

Contribution to: [2022 Snowmass Summer Study](#) · e-Print: [2209.08054](#) [physics.comp-ph]

[pdf](#) [cite](#) [claim](#) [reference search](#) [2 citations](#)

Data and Analysis Preservation, Recasting, and Reinterpretation #2

Stephen Bailey (LBL, Berkeley), Christian Bierlich (Lund U. (main)), Andy Buckley (Glasgow U.), Jon Butterworth (University Coll. London), Kyle Cranmer (New York U.) et al. (Mar 18, 2022)

Contribution to: [2022 Snowmass Summer Study](#) · e-Print: [2203.10057](#) [hep-ph]

[pdf](#) [cite](#) [claim](#) [reference search](#) [4 citations](#)

Signal region combination with full and simplified likelihoods in MadAnalysis 5 #2

Gaël Alguero (LPSC, Grenoble and Annecy, LAPTH), Jack Y. Araz (Durham U., IPPP), Benjamin Fuks (Paris, LPTHE), Sabine Kraml (LPSC, Grenoble) (Jun 29, 2022)

Published in: *SciPost Phys.* 14 (2023) 009 · e-Print: [2206.14870](#) [hep-ph]

[pdf](#) [DOI](#) [cite](#) [claim](#) [reference search](#) [1 citation](#)

Publishing statistical models: Getting the most out of particle physics experiments #4

Kyle Cranmer (New York U.), Sabine Kraml (LPSC, Grenoble), Harrison B. Prosper (Florida State U.), Philip Bechtle (Bonn U.), Florian U. Bernlochner (Bonn U.) et al. (Sep 10, 2021)

Published in: *SciPost Phys.* 12 (2022) 1, 037, *SciPost Phys.* 12 (2022) 037 · e-Print: [2109.04981](#) [hep-ph]

[pdf](#) [DOI](#) [cite](#) [claim](#) [reference search](#) [23 citations](#)

Reinterpretation of LHC Results for New Physics: Status and Recommendations after Run 2 #8

LHC Reinterpretation Forum Collaboration · Waleed Abdallah (Harish-Chandra Res. Inst. and Cairo U.) et al. (Mar 19, 2020)

Published in: *SciPost Phys.* 9 (2020) 2, 022 · e-Print: [2003.07868](#) [hep-ph]

[pdf](#) [links](#) [DOI](#) [cite](#) [claim](#) [reference search](#) [57 citations](#)

SciPost Physics

Submission

Publishing statistical models: Getting the most out of particle physics experiments

Kyle Cranmer^{1*}, Sabine Kraml^{2†}, Harrison B. Prosper^{3§} (editors), Philip Bechtle⁴, Florian U. Bernlochner⁴, Itay M. Bloch⁵, Enzo Canonero⁶, Marcin Chrzaszcz⁷, Andrea Coccaro⁸, Jan Conrad⁹, Glen Cowan¹⁰, Matthew Feickert¹¹, Nahuel Ferreira Iachellini^{12,13}, Andrew Fowlie¹⁴, Lukas Heinrich¹⁵, Alexander Held¹, Thomas Kuhr^{3,16}, Anders Kveltestad¹⁷, Maeve Madigan¹⁸, Farvah Mahmoudi^{15,19}, Knut Dundas Morá²⁰, Mark S. Neubauer¹¹, Maurizio Pierini¹⁵, Juan Rojo⁸, Sezen Sekmen²², Luca Silvestrini²³, Veronica Sanz^{24,25}, Giordon Stark²⁶, Riccardo Torre⁸, Robert Thorne²⁷, Wolfgang Waltenberger²⁸, Nicholas Wardle²⁹, Jonas Wittbrodt³⁰

1 New York University, USA **2** LPSC Grenoble, France **3** Florida State University, USA **4** University of Bonn, Germany **5** School of Physics and Astronomy, Tel-Aviv University, Israel **6** University of Genova, Italy **7** Institute of Nuclear Physics, Polish Academy of Sciences, Krakow, Poland **8** INFN, Sezione di Genova, Italy **9** Oskar Klein Centre, Stockholm University, Sweden **10** Royal Holloway, University of London, UK **11** University of Illinois at Urbana-Champaign, USA **12** Max Planck Institute for Physics, Munich, Germany **13** Exzellenzcluster ORIGINS, Garching, Germany **14** Nanjing Normal University, Nanjing, PRC **15** CERN, Switzerland **16** Ludwig-Maximilians-Universität München, Germany **17** University of Oslo, Norway **18** DAMTP, University of Cambridge, UK **19** Lyon University, France **20** Columbia University 10027, USA **21** VU Amsterdam and Nikhef, The Netherlands **22** Kyungpook National University, Daegu, Korea **23** INFN, Sezione di Roma, Italy **24** University of Sussex, UK **25** IFIC, Universidad de Valencia-CSIC, Spain **26** SCIPP, UC Santa Cruz, CA, USA **27** University College London, UK **28** HEPHY and University of Vienna, Austria **29** Imperial College London, UK **30** Lund University, Sweden

* kyle.cranmer@nyu.edu, † sabine.kraml@lpsc.in2p3.fr, § harry@hep.fsu.edu
(mailing list: [open-likelihoods@cern.ch](#))

September 9, 2021

Abstract

The statistical models used to derive the results of experimental analyses are of incredible scientific value and are essential information for analysis preservation and reuse. In this paper, we make the scientific case for systematically publishing the full statistical models and discuss the technical developments that make this practical. By means of a variety of physics cases — including parton distribution functions, Higgs boson measurements, effective field theory interpretations, direct searches for new physics, heavy flavor physics, direct dark matter detection, world averages, and beyond the Standard Model global fits — we illustrate how detailed information on the statistical modelling can enhance the short- and long-term impact of experimental results.

1


arXiv:2109.04981v1 [hep-ph] 10 Sep 2021

CMS+ATLAS Open Data

- CMS about to release more LHC Run II data
 - Will add to more than 2PB of open data and MC available
- Not necessarily easy to use?
 - Frequent workshops for the curious
- Yet, lots of publications, reuse for ML

Physics results obtained using Open Data (partial list)

- Search for dark matter production in association with the Z boson at the LHC in pp collisions at $\sqrt{s} = 8$ TeV using Monte Carlo simulations, S. Elgammal, M. Louka, A. Ellithi and M. Hussein, Chin.Phys.C 45 083001 (2021) [07 Mar 2021].
- Analysis of Prospective Super-Symmetry Inherent in the pp Collision Data at 7 TeV from CMS Collaboration Using Novel Two-Dimensional Multifractal-Detrended Fluctuation Analysis Method with Rectangular Scale, S. Bhaduri and A. Bhaduri, arXiv 2012.00442 (01 Dec 2020) [1 citation].
- Probing resonance states in high-energy interaction: a novel approach using complex network technique based on symmetry scaling, S. Bhaduri, A. Bhaduri and D. Ghosh, Eur.Phys.J.A 56 244 (2020) (30 Sep 2020) [1 citation].
- Searching for Supersymmetry at LHC Using the Complex-Network-Based Method of the Three-Dimensional Visibility-Graph, S. Bhaduri and A. Bhaduri, MDPI Physics 2 (2020) (10 Sep 2020).
- Lorentz and permutation invariants of particles III: Constraining nonstandard sources of parity violation, C. Lester, W. Haddadin and B. Griपाios, Int.J.Mod.Phys.A 37 2250093 (2022) (12 Aug 2020) [4 citations].
- The Hidden Geometry of Particle Collisions, P. Komiske, E. Metodiev and J. Thaler, JHEP 07 006 (2020) (21 Apr 2020) [26 citations]. *Uses CMS Open Data*
- Reinterpretation of LHC Results for New Physics: Status and Recommendations after Run 2, W. Abdallah et al., SciPost Phys. 9 022 (2020) (19 Mar 2020) [56 citations].
- Dataset of tau neutrino interactions recorded by the OPERA experiment, G. De Lellis, S. Dmitrievsky, G. Galati, A. Lavasa, T. Šimko, I. Tsanaktsidis and A. Ustyuzhanin, EPJ Web Conf. 245 08013 (2020) (01 Jan 2020).
- Study of Di-muon Production Process in pp Collision in CMS Data from Symmetry Scaling Perspective, S. Bhaduri, A. Bhaduri and D. Ghosh, Adv.High Energy Phys. 2020 4510897 (2020) (22 Nov 2019) [6 citations].
- Exploring the Space of Jets with CMS Open Data, P. Komiske, R. Mastandrea, E. Metodiev, P. Naik and J. Thaler, Phys.Rev.D 101 034009 (2020) (22 Aug 2019) [32 citations].
- Testing non-standard sources of parity violation in jets at the LHC, trialled with CMS Open Data, C. Lester and M. Schott, JHEP 12 120 (2019) (25 Apr 2019) [14 citations].
- Searching in CMS Open Data for Dimuon Resonances with Substantial Transverse Momentum, C. Cesarotti, Y. Soreq, M. Strassler, J. Thaler and W. Xue, Phys.Rev.D 100 015021 (2019) (11 Feb 2019) [12 citations].
- Symmetry-Scaling Based Complex Network Approach to Explore Exotic Hadronic States in High Energy Collision, S. Bhaduri, A. Bhaduri and D.



Hosted by the Data Preservation and Open Access Group of the CMS Collaboration

opendata CMS

Nein, Professor Bohr. First, you apply the trigger mask to your Monte Carlo, then...

CMS Open Data Workshop


Aug 1st - 4th, 2022 CERN, Geneva, CH

Use of Open Data to support software, machine learning and algorithms development

- Analysis-Specific Fast Simulation at the LHC with Deep Learning, C. Chen, O. Cerri, T. Nguyen, J. Vilimant and M. Pierini, Comput.Softw.Big Sci. 5 15 (2021) (09 Jun 2021) [5 citations].
- Evaluating Query Languages and Systems for High-Energy Physics Data [Extended Version], D. Graur, I. Müller, M. Proffitt, G. Fourny, G. Watts and G. Alonso, arXiv 2104.12615 (26 Apr 2021) [1 citation]. *Uses CMS Open Data*
- End-to-end jet classification of boosted top quarks with the CMS open data, M. Andrews et al., EPJ Web Conf. 251 04030 (2021) (19 Apr 2021) [7 citations].
- Porting CMS Heterogeneous Pixel Reconstruction to Kokkos, M. Kortelainen, M. Kwok, T. Childers, A. Strelchenko and Y. Wang, EPJ Web Conf. 251 03034 (2021) (14 Apr 2021). *Uses CMS Open Data*
- Data Analysis with GPU-Accelerated Kernels, J. Pata, I. Dutta, N. Lu, J. Vilimant, H. Newman, M. Spiropulu, C. Reissel and D. Ruini, PoS ICHEP2020 908 (2021) (02 Mar 2021) [2 citations]. *Uses ATLAS and CMS Open Data*
- Performance of CUDA Unified Memory in CMS Heterogeneous Pixel Reconstruction, M. Kortelainen and M. Kwok, EPJ Web Conf. 251 03035 (2021) (25 Feb 2021). *Uses CMS Open Data*
- Machine Learning scientific competitions and datasets, D. Rousseau and A. Ustyuzhanin, arXiv 2012.08520 (15 Dec 2020) [3 citations].
- Quantum-inspired machine learning on high-energy physics data, T. Felsner, M. Trenti, L. Sestini, A. Gianelle, D. Zuliani, D. Lucchesi and S. Montanero, npj Quantum Inf. 7 111 (2021) (28 Apr 2020) [20 citations].
- A meta-algorithm for classification using random recursive tree ensembles: A high energy physics application, V. Lalchand and A. Faul, arXiv 2001.06880 (19 Jan 2020). *Thesis, Uses ATLAS Open Data*
- End-to-end particle and event identification at the Large Hadron Collider with CMS Open Data, J. Alison, S. An, P. Bryant, B. Burkle, S. Gleyzer, M. Narain, M. Paulini, B. Poczvos and E. Usai, arXiv 1910.07029 (15 Oct 2019) [3 citations].
- Interaction networks for the identification of boosted H \rightarrow b \bar{b} decays, E. Moreno, T. Nguyen, J. Vilimant, O. Cerri, H. Newman, A. Perival, M. Spiropulu, J. Duarte and M. Pierini, Phys.Rev.D 102 012010 (2020) (26 Sep 2019) [48 citations]. *Uses CMS Open Data*
- Processing Columnar Collider Data with GPU-Accelerated Kernels, J. Pata and M. Spiropulu, arXiv 1906.06242 (14 Jun 2019) [1 citation]. *Uses CMS Open Data*
- End-to-end jet classification of quarks and gluons with the CMS Open Data, M. Andrews, J. Alison, S. An, P. Bryant, B. Burkle, S. Gleyzer, M. Narain, M. Paulini, B. Poczvos and E. Usai, Nucl.Instrum.Meth.A 977 164304 (2020) (21 Feb 2019) [30 citations].
- Exploring End-to-end Deep Learning Applications for Event Classification at CMS, M. Andrews, M. Paulini, S. Gleyzer and B. Poczvos, EPJ Web Conf.

AGC analysis task & Open Data

Community benchmark built on
CMS Open Data



- Analysis task: **ttbar cross-section measurement** in single lepton channel
 - Includes simple top reconstruction
 - Captures relevant workflow aspects and can easily be extended
 - E.g. conversion into a BSM search
 - Analysis task prominently features handling of systematic uncertainties
- Analysis is based on **Run-2 CMS Open Data** (~400 TB of MiniAOD available)
 - Open Data is crucial: everyone can participate
 - Currently using 4 TB of ntuple inputs (pre-converted, ~1B events before cuts)
- Goal of setup is showing **functionality**, not discovering new physics
 - Want to capture workflow; use made-up tools for calibrations & systematic uncertainties

Preservation & Reinterpretation

RECAST: Extending the Impact of Existing Analyses

#3

Kyle Cranmer (New York U., CCPP), Itay Yavin (New York U., CCPP) (Oct, 2010)

Published in: *JHEP* 04 (2011) 038 • e-Print: [1010.2506](https://arxiv.org/abs/1010.2506) [hep-ex]


pdf DOI cite claim reference search 76 citations

Archiving Real ATLAS Analyses

Using **Industry Standard** Software Packaging to archive analysis:

- Linux Containers ("Docker")
- Integrated into existing analysis infrastructure (revision control, continuous integration, grid computing)

Plain-text JSON formats to capture commands and workflows
Close coordination with CERN Analysis Preservation / Reuse Projects



Scalable Declarative HEP Analysis Workflows for Containerised Compute Clouds

#1

Tibor Šimko (CERN), Lukas Alexander Heinrich (CERN), Clemens Lange (CERN), Adelina Eleonora Lintuluoto (CERN and Helsinki U.), Danika Marina MacDonell (Victoria U.) et al. (May 7, 2021)

Published in: *Front. Big Data* 4 (2021) 661501, *Frontiers in Big Data* 4 (2021)

pdf DOI cite claim reference search 1 citation

REANA: A System for Reusable Research Data Analyses

#2

Tibor Šimko (CERN), Lukas Heinrich (New York U. (main)), Harri Hirvonsalo (Espoo, Sci. Computing Ctr.), Dinos Kousidis (CERN), Diego Rodríguez (CERN) (Dec 3, 2018)

Published in: *EPJ Web Conf.* 214 (2019) 06034 • Contribution to: CHEP 2018

pdf DOI cite claim reference search 11 citations

Analysis Preservation and Systematic Reinterpretation within the ATLAS experiment

#3

ATLAS Collaboration • Kyle Cranmer (New York U.) et al. (Oct 18, 2018)

Published in: *J.Phys.Conf.Ser.* 1085 (2018) 4, 042011 • Contribution to: ACAT 2017

pdf DOI cite claim reference search 6 citations

HEP Software Foundation Community White Paper Working Group - Data and Software Preservation to Enable Reuse

#4

M.D. Hildreth (Notre Dame U.), A. Boehlein (Jefferson Lab), K. Cranmer (New York U.), S. Dallmeier-Tiessen (CERN), R. Gardner (Chicago U.) et al. (Oct 2, 2018)

e-Print: [1810.01191](https://arxiv.org/abs/1810.01191) [physics.comp-ph]

pdf links cite claim reference search 2 citations

Yadage and Packtivity - analysis preservation using parametrized workflows

#5

Kyle Cranmer (New York U.), Lukas Heinrich (New York U.) (Jun 6, 2017)


Published in: *J.Phys.Conf.Ser.* 898 (2017) 10, 102019 • Contribution to: CHEP 2016 • e-Print: [1706.01878](https://arxiv.org/abs/1706.01878) [physics.data-an]

pdf DOI cite claim reference search 19 citations


Preservation & Reinterpretation

First results using the RECAST reinterpretation framework and publishing full statistical likelihoods (using pyhf)





ATLAS PUB Note
ATL-PHYS-PUB-2019-029
5th August 2019




Reproducing searches for new physics with the ATLAS experiment through publication of full statistical likelihoods

The ATLAS Collaboration


The ATLAS Collaboration is starting to publicly provide likelihoods associated with statistical fits used in searches for new physics on HEPData. These likelihoods adhere to a specification first defined by the BiStFactory public template. This note introduces a JSON schema that fully describes the BiStFactory statistical model and is sufficient to reproduce key results from published ATLAS analyses. This is peer independent of its implementation in ROOT and it can be used to run statistical analysis outside of the ROOT and RooStats/ROOT framework. The first of these likelihoods published on HEPData is from a search for bottom-squark pair production. Using two independent implementations of the model, one in ROOT and one in pure Python, the limits on the bottom-squark mass are reproduced, underscoring the implementation independence and long-term viability of the archived data.

ATL-PHYS-PUB-2019-029
5 August 2019

© 2019 CERN for the benefit of the ATLAS Collaboration.
Reproduction of this article or parts of it is allowed as specified in the CC-BY-4.0 license.



ATLAS PUB Note
ATL-PHYS-PUB-2019-032
11th August 2019



RECAST framework reinterpretation of an ATLAS Dark Matter Search constraining a model of a dark Higgs boson decaying to two *b*-quarks

The ATLAS Collaboration

The reinterpretation of a search for dark matter produced in association with a Higgs boson decaying to *b*-quarks performed with RECAST, a software framework designed to facilitate the reinterpretation of existing searches for new physics, is presented. Reinterpretation using RECAST is enabled through the sustainable preservation of the original data analysis as re-executable declarative workflows using modern cloud technologies and integrated with the wider CERN Analysis Preservation efforts. The reinterpretation targets a model predicting dark matter production in association with a hypothetical dark Higgs boson decaying into *b*-quarks where the mass of the dark Higgs boson m_h is a free parameter, necessitating a faithful reinterpretation of the analysis. The dataset has an integrated luminosity of 79.8 fb^{-1} and was recorded with the ATLAS detector at the Large Hadron Collider at a centre-of-mass energy of $\sqrt{s} = 13 \text{ TeV}$. Constraints on the parameter space of the dark Higgs model for a fixed choice of dark matter mass $m_\chi = 200 \text{ GeV}$ exclude model configurations with a mediator mass up to 3.2 TeV .

ATL-PHYS-PUB-2019-032
12 August 2019

© 2019 CERN for the benefit of the ATLAS Collaboration.
Reproduction of this article or parts of it is allowed as specified in the CC-BY-4.0 license.


Scalable ATLAS pMSSM computational workflows using containerised REANA reusable analysis platform

M. Donadoni¹, L. Heinrich², Y. Liu³, A. Meclonis¹, V. Moiseichenkov¹, G. Stark⁴, T. Simko¹, M. Vidal Garcia¹

¹ CERN, Geneva, Switzerland
² Max-Planck-Institut für Physik, München, Germany
³ Sun Yat-sen University, Guangzhou, China
⁴ University of California, Santa Cruz, United States

ATLAS pMSSM searches


We have developed a streamlined framework for large-scale pMSSM reinterpretations of ATLAS analyses of LHC Run 2 using containerised computational workflows. The project is looking to assess the global coverage of BSM physics and requires running numerous computational workflows representing pMSSM model points.



ATLAS pMSSM studies from LHC Run 1. [arXiv:1508.06602](https://arxiv.org/abs/1508.06602)

REANA reusable analysis platform

The computational workflows were run at scale using the REANA reusable analysis platform. The workflows typically run on Kubernetes clusters.



The architecture of the REANA cloud platform. Users can use a command-line client and a web interface to submit containerised workflows that are then orchestrated on supported compute backends.

Selection of analyses


Following ATLAS analysis preservation policies, many ATLAS analyses have been preserved as containerised Yastage workflows. After validation they are added to a curated selection of analyses suitable for the pMSSM study.



ATLAS SUSY analyses preserved on ODLB.

Scaling up to O(5k) workflows

We have improved the REANA platform scheduling in order to maximise the scheduling throughput of incoming workflows.




A scalability test submitting 250 workflows every 10 minutes. A cluster with 448 cores (top) cannot keep up with the load. A cluster with 1072 cores (bottom) can comfortably host the incoming workload.




The benchmark tests were running in the CERN Computer Centre and on the Google Cloud Platform public cloud. The REANA scheduling parameters were optimised to maximise CPU utilisation for the pMSSM workflows taking into account the three time-consuming coupling jobs per one workflow run.

Computational workflow

One typical containerised workflow consists of three time-consuming coupling steps that run in parallel and the fitting steps that run afterwards. The workflow dependency graph is simple; the complexity relies in having to run O(5k) of these workflows in order to cover sufficient number of pMSSM model points.



A typical pMSSM workflow. The computational runtime is about 10 minutes without systematics (fast payload) and about 10 hours with all systematics (real payload).



The workload burstdown throughput rate is sustainable over a long period of time.

Conclusions

- ATLAS searches for supersymmetry are effectively preserved with computational workflow recipes enabling their future reuse and reinterpretation.
- We have launched several ATLAS pMSSM workflows on REANA and studied the workflow burstdown throughput rate as a function of increasing Kubernetes cluster size.
- REANA platform has been extended to support the workload of many concurrent workflows. The solution was benchmarked on medium to large clusters (from 500 to 5000 cores).
- It is essential to adapt cluster parameters to the type of workflows in order to ensure best throughput and cluster resource utilisation (CPU per node, RAM per node).
- The developed system is ready to run large-scale ATLAS pMSSM reinterpretations of LHC Run 2 analyses.



This project was carried out in a close collaboration between ATLAS and CERN IT.

reana
www.reana.io

Docs
docs.reana.io

Email
info@reana.io

GitHub
@reanahub

Twitter
@reanahub


Preservation & Training

Analysis Preservation Technology and Infrastructure are going mainstream

- GitLab, Docker, REANA workflows, etc.
- Training is essential



Josh McFayden @JoshMcFayden · Feb 19, 2020
SUCCESS! ✓ 🎉



1 1 7




Participants in Analysis Preservation Bootcamp showing off their ability to reproduce an LHC analysis. Photo Credit: Samuel Meehan

Bootcamp: <https://indico.cern.ch/event/854880/>
Blog post: <https://iris-hep.org/2020/02/17/analysis-preservation.html>

Building systems on top of reusable, preserved analyses

Once we have a library of reusable analyses, we can build applications on top of them.

Here, ATLAS uses an active learning technique to efficiently reinterpret an analysis over a 4-D parameter space.



ATLAS PUB Note
ATL-PHYS-PUB-2022-045
November 3, 2022



Active Learning reinterpretation of an ATLAS Dark Matter search constraining a model of a dark Higgs boson decaying to two b -quarks


The ATLAS Collaboration

A reinterpretation of a search for dark matter produced in association with a Higgs boson decaying to b -quarks using Active Learning, a technique to facilitate efficient and comprehensive inference in multi-dimensional new physics parameter spaces, is presented. The dataset has an integrated luminosity of 139 fb^{-1} and was recorded with the ATLAS detector at the Large Hadron Collider at a centre-of-mass energy of $\sqrt{s}=13 \text{ TeV}$. The reinterpretation refers to a model predicting dark matter production in association with a dark sector Higgs boson decaying to b -quarks. The Active Learning approach makes use of a Gaussian Process to determine the exclusion limit contour and a corresponding uncertainty in a four-dimensional new physics parameter space. Each exclusion limit is determined accurately by means of the RECAST protocol. The combined approach of RECAST and Active Learning provides a blueprint for accurate, efficient and comprehensive interpretations of new physics searches at the Large Hadron Collider.

FAIROS-HEP + AGC

AGC analysis task & Open Data

Community benchmark built on
CMS Open Data




- Analysis task: **ttbar cross-section measurement** in single lepton channel
 - Includes simple top reconstruction
 - Captures relevant workflow aspects and can easily be extended
 - E.g. conversion into a BSM search
 - Analysis task prominently features handling of systematic uncertainties
- Analysis is based on **Run-2 CMS Open Data** (~400 TB of MiniAOD available)
 - Open Data is crucial: everyone can participate
 - Currently using 4 TB of ntuple inputs (pre-converted, ~1B events before cuts)
- Goal of setup is showing **functionality**, not discovering new physics
 - Want to capture workflow; use made-up tools for calibrations & systematic uncertainties

Analysis Facilities impact on Analysis Preservation & Reinterpretation

Analysis preservation, reuse / reinterpretation / RECAST will look significantly different when coupled to analysis facilities.

Coffea-casa Analysis Facility is glueing together different areas of IRIS-HEP (AS, SSL & DOMA), providing **AGC execution environment** to explore analysis workflows at scale



Social scientists interested in our field

MENU ≡

Home / Nicole C. Nelson



Nicole C. Nelson

Associate Professor in the Department of Medical History and Bioethics

nicole.nelson@wisc.edu
608.265.3525

Office: 1426 Medical Sciences Center
[Curriculum Vitae \(pdf\)](#) | [Website](#)
Office Hours: Mondays 2:00-3:00pm and Wednesdays 3:30-4:30pm in 1426 Medical Sciences Center

Joint Appointment: [Medical History & Bioethics](#)

Biography

I am an Associate Professor in the Department of Medical History and Bioethics at the University of Wisconsin–Madison. My other affiliations are with the Holtz Center for Science and Technology Studies and the History department. Previously I was a postdoctoral research fellow in Social Studies of Medicine at McGill University.

My research examines scientists’ assumptions about the natural world and how these assumptions shape scientific practice. In my book, *Model Behavior*, I explore how animal behavior geneticists’ beliefs about the complexity of gene action and of psychiatric disorders are reflected in their research with mouse models. For this work I won a First Book Award from the UW–Madison Center for the Humanities

[Programs](#) [Graduate](#) [Undergraduate](#) [Continuing Education](#) [iSchool Research](#) [Alumni](#) [Faculty & Staff Directory](#)



Naomi Mine

PhD Student

nmine@wisc.edu
pronouns
she | her | hers

Education: University of Chicago, BA 2015 (Interdisciplinary Studies in the Humanities, Visual Arts)

Areas of interest: Technology-centered social movements; sociology of quantification; science and technology studies (STS)

Classes taught:
LIS 201: The Information Society (Teaching Assistant)
LIS 461: Data and Algorithms: Ethics and Policy (Teaching Assistant)

Website: <http://naomimine.com>

FAIROS-HEP: Future Workshop Trajectory

Workshop 2: Broader Community Engagement and Theory Reinterpretation

Attendees: Particle Physics Experimentalists and Theorists

Location: TBD

Workshop 3: Specific Reuse Case: Deriving EFT Results from Future LHC Data

Attendees: Participants from the LHC experiments and theorists working on EFT interpretations

Location: US University, TBD

Workshop 4: Broader Engagement: CNI and External Science Partners

Attendees: Core RCN members, CNI membership, External Science Partners

Location: Semi-Annual CNI Membership Meeting

Workshop 5: Reuse Case: Kinematic RECASTing for New Physics Discovery

Attendees: Participants from the LHC experiments and theorists working on new physics searches

Location: US University

Thank you!

Questions?

FAIROS-HEP: “The Network”

GO FAIR US: US office of GO FAIR

DANCE Network: “Dark matter and Neutrino Computation Explored”

- Dark matter experiments (XENONnT, DarkSide, SuperCDMS, LZ, DEAP, ARGO), neutrino-oriented experiments (incl. μ BooNe, NOvA, COHERENT, HyperK), and double- beta-decay oriented experiments (incl. EXO, nEXO)

DOE Labs

Electron-Ion collider

Zenodo

HEPData

InspireHEP

CERN Open Data Portal

CERN Analysis Preservation Portal

REANA

RECAST

arXiv

FAIR4HEP

PUNCH4NFDI

IRIS-HEP

The NSF's FAIROS Research Coordination Networks

We are one of 10 funded projects:

DBER+ Commons

- FAIR in Education research

MaRCN

- Open science in materials

FAIR facilities and instruments

- PIDs for research instruments

FARR

- Best practices for ML/AI

Geospatial Big Data Infrastructure

- Environmental research

Paleobio/zooarchaeology databases

- Community-coordinated resources

SEEKCommons project

- Bridge social and environ. sciences

NoCTURN

- Non-clinical tomography

REPETO

- Reproducibility in CS Education