


Focus topic MC needs: Bhabha luminometry

S. Jadach, W. Płaczek, M. Skrzypek, B.F.L. Ward, S.A. Yost

Partly supported by the CERN FCC Design Study Programme.

ECFA Higgs Factories: 2nd Topical Meeting on Generators, 21-22 June 2023

Dedicated to the memory of Staszek Jadach





XXX Cracow EPIPHANY Conference

on Precision Physics at High Energy Colliders
dedicated to the memory of Staszek Jadach

8-12 January 2024

[Home](#) [General Information](#) [Registration](#) [Programme](#) [Proceedings](#) [Participants](#) [Previous Conferences](#)

Intro – lumi basics



Bhabha cross sect. depends on detector acceptance angles

$$\sigma_{Bh} \simeq 4\pi\alpha^2 \left(\frac{1}{t_{\min}} - \frac{1}{t_{\max}} \right) = 4\pi\alpha^2 \left(\frac{t_{\max} - t_{\min}}{\bar{t}^2} \right), \quad \bar{t} = \sqrt{t_{\min} t_{\max}}$$

\bar{t} is the characteristic scale of the process

\bar{t}/s is the suppression factor between s - and t -channel contributions

Machine	$\theta_{\min} \div \theta_{\max}$ [mrad]	\sqrt{s} [GeV]	$\bar{t}/s \simeq \bar{t}^2/4$	$\sqrt{\bar{t}}$ [GeV]
LEP	28 \div 50	M_Z	3.5×10^{-4}	1.70
FCCee	64 \div 86	M_Z	13.7×10^{-4}	3.37
FCCee	64 \div 86	240	13.7×10^{-4}	8.9
FCCee	64 \div 86	350	13.7×10^{-4}	13.0
ILC	31 \div 77	500	6.0×10^{-4}	12.2
ILC	31 \div 77	1000	6.0×10^{-4}	24.4
CLIC	39 \div 134	3000	13.0×10^{-4}	108

Luminosity today – BHLUMI status



The 2019 update comes from P. Janot & S. Jadach Phys.Lett.B 803 (2020) 135319

Type of correction / Error	1999	Update 2019
(a) Photonic $\mathcal{O}(L_e \alpha^2)$	0.027%	0.027%
(b) Photonic $\mathcal{O}(L_e^3 \alpha^3)$	0.015%	0.015%
(c) Vacuum polariz.	0.040%	0.009%
(d) Light pairs	0.030%	0.010%
(e) Z and s -channel γ exchange	0.015%	0.015%
(f) Up-down interference	0.0014%	0.0014%
(f) Technical Precision	–	(0.027)%
Total	6.1×10^{-4}	3.7×10^{-4}

Table: Summary of the total (physical+technical) theoretical uncertainty for a typical calorimetric LEP luminosity detector within the generic angular range of 18–52 mrad. Total error is summed in quadrature.

- ▶ Hadronic vacuum polarisation from F. Jegerlehner (fortran code `hadr5x.f`) 2019
- ▶ Light pairs: real – FERMISV MC by J. Hilgart et.al. 1993 and KoralW by S. Jadach et.al.; virtual – S. Actis et.al. 2008

Current BHLUMI precision forecast for FCCee



Type of correction / Error	M_Z (2019) [1]	240 GeV	350 GeV [2]
(a) Photonic $\mathcal{O}(L_e \alpha^2)$	0.027%	0.032%	0.033%
(b) Photonic $\mathcal{O}(L_e^3 \alpha^3)$	0.015%	0.026%	0.028%
(c) Vacuum polariz.	0.009%	0.020%	0.022%
(d) Light pairs	0.010%	0.015%	0.015%
(e) Z and s -channel γ exchange	0.09%	0.25% (0.034%)	0.5% (0.07%)
(f) Up-down interference	0.009%	0.010%	0.010%
(g) Technical Precision	[0.027%]		
Total	10×10^{-4}	25×10^{-4} (6×10^{-4})	50×10^{-4} (8.7×10^{-4})

Table: Entries in curly brackets represent hypothetic situation with all Born-level interferences included in BHLUMI

Entry (c) for M_Z optimistic, 0.015% more realistic

Few times worse than at LEP !!

[1] S. Jadach *et.al.* Phys. Lett B790 (2019) 314

[2] S. Jadach *et.al.* Eur. Phys. J. C (2021) 81:1047

Photonic corrections

- ▶ Included in BHLUMI: $\mathcal{O}(\alpha + \alpha^2 L^2)$ -YFS exponentiated
- ▶ To be added: to BHLUMI $\mathcal{O}(\alpha^3 L^3)$ and $\mathcal{O}(\alpha^2 L^1)$ – known
- ▶ Errors: $\mathcal{O}(\alpha^4 L^4)$ and $\mathcal{O}(\alpha^3 L^2)$
 - ▶ reference points – LEP:
 $\mathcal{O}(\alpha^3 L^3) \simeq 1.5 \times 10^{-4}$ and $\mathcal{O}(\alpha^2 L^1) \simeq 2.7 \times 10^{-4}$
 - ▶ estimated based on LEP analysis and scale $(\alpha/\pi)^n L^m$
 - ▶ scale with energy/angles as $\ln^m(\bar{t}_{xx}/m_e^2)$
- ▶ Likely not needed: $\mathcal{O}(\alpha^2 L^0)$ – known
 $\sim \mathcal{O}(\alpha^2 L^1)/L \simeq 2.7 \times 10^{-4}/16.3 \simeq 0.17 \times 10^{-4}$

$(\gamma_s + Z_s + \gamma_t + Z_t)^{\otimes 2}$ EW interferences



- ▶ Included in BHLUMI: $(\gamma_s + Z_s) \otimes \gamma_t$
- ▶ To be added:
 - ▶ complete Born – trivial
 - ▶ complete $\mathcal{O}(\alpha_{EW})$ – known, e.g. BHWIDE
- ▶ Error: $\mathcal{O}(\alpha_{EW}^2)$
 - ▶ estimated at FCCee(M_Z) based on analysis of S. Jadach *et.al.* Phys. Lett B790 (2019) 314 – from BHWIDE
 - ▶ estimated at other energies/angles based on analysis done with $\mathcal{O}(\alpha_{EW})$ DIZET/ZFITTER (by changing switch NPAR(2) from 2 to 3) M. Battaglia, S. Jadach, D. Bardin, *eConf C010630* (2001) E3015, <http://www.slac.stanford.edu/econf/C010630/papers/E3015.PDF> for the energies of 800 GeV and 3 TeV.
Extrapolation from 800 to 350/240 GeV not done \Rightarrow error likely overestimated (factor of 2-3 ???)
 - ▶ Error at higher \bar{t}/M_Z^2 almost entirely from $\gamma_t \otimes Z_t$ interference
- ▶ Amplitude-level exponentiation (KKMC-style) needed to account for leading $\mathcal{O}(\alpha_{EW}^2)$ corrs.

QED photonic up-down interference

- ▶ Missing in BHLUMI

size at $\mathcal{O}(\alpha)$: $0.07 \times \bar{t}_{xx}/s$ – easy to include,
 \bar{t}_{xx}/s depends only on angles

LEP → FCCee: t/s grows 4 times (LEP → ILC: 2 times)

- ▶ Error: h.o.t. – suppressed by $(\alpha/\pi) \ln(\bar{t}_{xx}/m_e^2)$ times safety factor of 2 ($\mathcal{O}(\alpha_{QED}^2)$ calculations exist) – almost negligible

Vacuum polarisation



- ▶ Uncertainty due to vacuum polarisation:

$$\delta_{VP}\sigma/\sigma = 2\delta\alpha_{\text{eff}}(\bar{t})/\alpha_{\text{eff}}(\bar{t})$$

- ▶ $\delta\alpha_{\text{eff}}(\bar{t})$ from

F. Jegerlehner, *CERN Yellow Reports: Monographs 3* (2020) 9–37




Fig. B.1.15: Hadronic uncertainty $\delta\Delta\alpha_{\text{had}}(\sqrt{t})$. The progress since LEP times, from 1996 (left) to now (right) is remarkable. A great deal of much more precise low-energy data, $\pi\pi$, etc., are now available.

- ▶ $\alpha_{\text{eff}}(\bar{t})$ from
F. Jegerlehner, *Nucl. Phys. Proc. Suppl.* **162** (2006) 22–32
- ▶ By FCCee operation time factor of 2 improvement expected
(F. Jegerlehner)

Light pairs



- ▶ Current state of the art: BHLUMI + external four-fermion code + virtual semianalytical corrections
 - P. Janot and S. Jadach, *Phys. Lett. B* **803** (2020) 135319
- ▶ included components:
 - ▶ ee-pair, $\mu\mu$ -pair, $\tau\tau$ -pair, qq -pair with s-channel photonic emissions (FERMISV, KORALW)
 - ▶ result for LEP: $4 \times 10^{-4} \pm 1 \times 10^{-4}$
- ▶ future prospects for external *4fermion* code scenario
 - error components:
 - ▶ $4f + \gamma$ (25% of $4f$) – s vs. t mismatch $\sim 30\%$
 - ▶ $\mathcal{O}(\alpha)$ *4fermion* calculations exist for selected final states
 - ▶ $4f + 2\gamma$, $6f$
- ▶ future prospects for BHLUMI upgrade scenario
 - error components:
 - ▶ $4f + \gamma$ – absent – correct t -channel behavior (LL+soft),
 $\mathcal{O}(\alpha)$ *4fermion* likely not needed
 - ▶ $4f + 2\gamma$ – included via exponentiation + LL,
 - ▶ $6f$

Light pairs



Extrapolation to other energies/angles

- ▶ use LEP result for ff : $4 \times 10^{-4} \pm 1 \times 10^{-4}$ and scale with $\ln^2(\bar{t}_{xx}/m_{yy}^2)/\ln^2(\bar{t}_{LEP}/m_{yy}^2)$ (pairs)
- ▶ use LEP result for $ff\gamma$ terms: $20\% \times 4 \times 10^{-4}$
(G. Montagna, M. Moretti, O. Nicrosini, A. Pallavicini, and F. Piccinini, *Nucl. Phys.* **B547** (1999) 39–59),
and scale with
 $\ln(\bar{t}_{xx}/m_e^2)/\ln(\bar{t}_{LEP}/m_e^2)$ (photons)
- ▶ τ -pair (negligible at LEP) estimated relative to muon-pair as $\ln^2(\bar{t}_{xx}/m_\tau^2)/\ln^2(\bar{t}_{xx}/m_\mu^2)$
- ▶ hadron-pair estimated relative to muon-pair as $R_{had} \times \ln^2(\bar{t}_{xx}/(0.5\text{GeV})^2)/\ln^2(\bar{t}_{xx}/m_\mu^2)$

Lumi at FCCee: Technical precision



- ▶ At LEP BHLUMI technical prec. was tested in two ways:
 - ▶ Comparison with semian. integration of $\mathcal{O}(\alpha^2)_{exp}$ matrix el. of BHLUMI: agreement 2.7×10^{-4}
 - ▶ Comparison with LUMLOG+OLDBIS hybrid MC and with SABSPV MC. All of these MCs have incomplete soft resummation: agreement 2.7×10^{-4} (for sharp photon energy cut-offs 1.7×10^{-3})
- ▶ Now another MC code BabaYaga [Balossini et.al.] with complete soft-photon resummation is available. After upgrade to NNLO in hard process it could be ideal for technical comparison with BHLUMI

Lumi at FCCee – Forecast



Forecast			
Type of correction / Error	FCCee _{M_Z} [1]	FCCee ₂₄₀	FCCee ₃₅₀
(a) Photonic $\mathcal{O}(L_e^2 \alpha^3)$	0.10×10^{-4}	0.10×10^{-4}	0.13×10^{-4}
(b) Photonic $\mathcal{O}(L_e^4 \alpha^4)$	0.06×10^{-4}	$0.26 \times 10^{-4}(a)$	$0.27 \times 10^{-4}(a)$
(c) Vacuum polariz.	0.6×10^{-4}	1.0×10^{-4}	1.1×10^{-4}
(d) Light pairs	0.5×10^{-4}	0.4×10^{-4}	0.4×10^{-4}
(e) Z and s-channel γ exch.	0.1×10^{-4}	$1.0 \times 10^{-4}(*)$	$1.0 \times 10^{-4}(*)$
(f) Up-down interference	0.1×10^{-4}	0.09×10^{-4}	0.1×10^{-4}
Total	1.0×10^{-4}	1.5×10^{-4}	1.6×10^{-4}

Numbers: (*) likely overestimated, (a) include safety factor 2. Technical error is not included

Precision dominated by:

- ▶ Vacuum polarisation (c) – seems irreducible.
- ▶ The EW $\mathcal{O}(\alpha^2)$ hard process uncertainty (e). Numbers (*) are likely overestimated (taken from 800 GeV estimate)
– factor 2 too big ???.

Precision loss at higher energies reasonable (?)
factor of 2 loss w.r.t. M_Z

[1] S. Jadach, W. Płaczek, M. Skrzypek, B. F. L. Ward, S. A. Yost, *Phys. Lett. B* **790** (2019) 314

Forecast study for FCCee M_Z		
Type of correction / Error	Published [1]	Redone
(a) Photonic $\mathcal{O}(L_e^2 \alpha^3)$	0.10×10^{-4}	0.10×10^{-4}
(b) Photonic $\mathcal{O}(L_e^4 \alpha^4)$	0.06×10^{-4}	0.06×10^{-4}
(b') Photonic $\mathcal{O}(\alpha^2 L_e^0)$		0.17×10^{-4}
(c) Vacuum polariz.	0.6×10^{-4}	0.6×10^{-4}
(d) Light pairs	0.5×10^{-4}	0.27×10^{-4}
(e) Z and s -channel γ exch.	0.1×10^{-4}	0.1×10^{-4}
(f) Up-down interference	0.1×10^{-4}	0.08×10^{-4}
Total	1.0×10^{-4}	0.70×10^{-4}

Lumi at FCCee M_Z – Forecast study

- ▶ (d) light pairs are re-analysed w.r.t. [1] (safety factor 1.25 is removed; $ff\gamma$ non-leading contrib. less conservative: $z_{cut} \leq .5$ can help; *hadr*-pair uncertainty is set to few % as in [2])
- ▶ (f) value not rounded up is used as compared to Ref. [1]
- ▶ "Total" value not rounded up is used as compared to Ref. [1] (above three entries corrected at 240 and 350 GeV as well)
- ▶ (b') missing non-logarithmic $\mathcal{O}(\alpha^2 L_e^0)$ correction added for completeness
- ▶ (e): size of $\mathcal{O}(\alpha^2)_{EW}$ corrs. to be revisited – available BHWIDE
CEEX amplitude level exponentiation instrumental (KKMC style) ?

[1] S. Jadach, W. Płaczek, M. Skrzypek, B. F. L. Ward, S. A. Yost, *Phys. Lett. B* **790** (2019) 314

[2] ALEPH Collaboration, D. Buskulic *et al.*, *Z. Phys. C* **66** (1995) 3–18

Possible precision $\sim 0.7 \times 10^{-4}$ within the reach ??



Vacuum polarisation

From BFL Ward's presentation at RADCOR 2023:

Note: Lattice methods with jegerlehner's results allow, in principle, (c) -> (c)/6

$$\Delta\alpha_{had}(t) = \Delta\alpha_{had}(-Q_0^2)|_{lat} + [\Delta\alpha_{had}(t) - \Delta\alpha_{had}(-Q_0^2)]|_{pQCD Adler}$$



Lattice results are mainly limited now by statistics (?), so if enough computing resources are available, the 0.1×10^{-4} precision at $-few \text{ GeV}^2$ may be feasible.

The above is more optimistic than the 3.5σ tension with estimates based on exp. data of R-ratio reported in arXiv: 2203.08676, 2211.11401 [hep-lat] for $\Delta\alpha_{had}^{(5)}(-Q^2)$, $Q^2 = 3 \div 7 \text{ GeV}^2$.

The precision of lattice results given in the above papers is $\Delta\alpha_{had}(-5 \text{ GeV}^2) = 0.00716 \pm 0.9 \times 10^{-4}$ – on par with R-ratio method.

EW corrections

In *S. Jadach, W. Płaczek, M. Skrzypek, B. F. L. Ward, and S. A. Yost, Phys. Lett. B 790 (2019) 314–321* we estimated

- ▶ $\mathcal{O}(\alpha_{EW}^2)$ uncertainties in BHWIDE at Z-peak:

Conservatively estimated as $\frac{\alpha}{\pi} \ln \frac{\bar{t}}{m_e^2} \times \mathcal{O}(\alpha_{\text{from exponentiation}}^2)$
times safety factor of 2.

This gives 0.7×10^{-4} for QED part and 0.3×10^{-4} for EW part.
Added linearly one obtains 1×10^{-4} .

- ▶ More aggressive estimate (no safety factor, added in quadratures)
would give 0.46×10^{-4}

(Not so) science-fiction: 0.1×10^{-4} at Z-peak?




DIZET analysis of EW corrs. done above Z-peak. At the peak different graphs contribute ($\gamma_t \otimes Z_s$ vs $\gamma_t \otimes Z_t$), but rough idea could be valid?

M. Battaglia, S. Jadach, and D. Bardin, eConf C010630 (2001) E3015

S. Jadach, "MC tools for extracting luminosity spectra. What do we need?".

<https://jadach.web.cern.ch/jadach/public/LumLCslac.pdf>, 2002

How big is, therefore, uncertainty of due to EW corrections?



DIZET: Varied $M_H = 120 \rightarrow 500\text{GeV}$, $M_t = 165 \rightarrow 185\text{GeV}$ and NPAR(2)=3 \rightarrow NPAR(2)=4

which manipulates non-leading 2-loop EW corrections $\mathcal{O}(G_F^2 M_t^2 M_Z^2)$, Degrassi et al.,

keeping 2-loop EW corrections $\mathcal{O}(G_F^2 M_t^4)$.

At 800 GeV $\mathcal{O}(\alpha_{EW}^2)$ contributes below 0.4×10^{-4} and decreases with energy decrease ($15. \times 10^{-4}$ at 3 TeV).

Bottom line: leading α_{EW}^2 contribs may be needed



Fermion pairs

One will probably need $\mathcal{O}(\alpha)$ corrections to four fermion final state.

- ▶ Calculations of Denner et.al. (*PLB 612(2005) 223*) exist for charged current final states. Claimed physical precision (due to higher orders) at WW threshold is few $\times 0.1\%$ of the 4f Born.
- ▶ The whole pair contribution to Bhabha is $\sim 4 \times 10^{-4}$. Assuming precision of 1% for NC final states we are **well below** 0.1×10^{-4} target, provided *t*-channel multiphotons are properly resummed.

Note, that above ~ 500 GeV Sudakov logs must be resummed.

Bottom line

0.1×10^{-4} precision *a priori* not excluded

Lumi forecast at ILC and CLIC GeV



Type of correction / Error	Forecast		
	ILC ₅₀₀	ILC ₁₀₀₀	CLIC ₃₀₀₀
(a) Photonic $\mathcal{O}(L_e^2 \alpha^3)$	0.13×10^{-4}	0.15×10^{-4}	0.20×10^{-4}
(b) Photonic $\mathcal{O}(L_e^4 \alpha^4)$	0.27×10^{-4}	0.37×10^{-4}	0.63×10^{-4}
(c) Vacuum polariz.	1.1×10^{-4}	1.1×10^{-4}	1.2×10^{-4}
(d) Light pairs	0.4×10^{-4}	0.5×10^{-4}	0.7×10^{-4}
(e) Z and s-channel γ exch.	$1.0 \times 10^{-4} (*)$	2.4×10^{-4}	16×10^{-4}
(f) Up-down interference	$< 0.1 \times 10^{-4}$	$< 0.1 \times 10^{-4}$	0.1×10^{-4}
Total	1.6×10^{-4}	2.7×10^{-4}	16×10^{-4}

Number (*) is somewhat overestimated (taken from 800 GeV estimate)

- ▶ Precision at high energies totally due to the EW $\mathcal{O}(\alpha^2)$ hard process uncertainty (e).
- ▶ EW interferences are dominated by $\gamma_t \otimes Z_t$ (15% of $\gamma_t \otimes \gamma_t$ at CLIC) and $Z_t \otimes Z_t$ (2% of $\gamma_t \otimes \gamma_t$ at CLIC)
 - usefull for $\mathcal{O}(\alpha_{EW}^2)$ calculation ?
- CEEX amplitude level exponentiation mandatory ?

**At 3 TeV loss of precision is dramatic,
dominant $\mathcal{O}(\alpha_{EW}^2)$ and CEEX are a must!**

Summary



- ▶ Our starting point is BHLUMI 4.04 with the inherited from LEP precision of 0.06%
- ▶ 2019 development of Janot&Jadach reduced this error to 0.037%
- ▶ The precision of BHLUMI for FCCee_{240} as of now is 25×10^{-4} and forecasted one is 1.5×10^{-4} , factor of 2 worse than at FCCee_{M_Z}
- ▶ At high energies forecasted precision deteriorates drastically, up to 16×10^{-4} for CLIC at 3 TeV, due to missing $\mathcal{O}(\alpha^2)_{EW}$ corrections
- ▶ Forecasted in Jadach et.al. (2019) precision 1×10^{-4} at FCCee_{M_Z} seems to be reducible to 0.7×10^{-4} by reducing error on pair emission and loosening conservative approach to safety factors; $\mathcal{O}(\alpha_{EW}^2)$ corrs must be revisited. Further precision improvement seems to be blocked by the error on vacuum polarisation contrib.
- ▶ Precision 0.1×10^{-4} at Z-peak could be discussed provided lattice QCD delivers vacuum polarisation with precision 0.1×10^{-4} (matter of CPU?), dominant $\mathcal{O}(\alpha_{EW}^2)$ corrs to Bhabha and $\mathcal{O}(\alpha_{EW})$ corrs to 4-fermions are available.
- ▶ Technical precision requires second MC code, e.g. BABAYAGA