

Future Accelerator Projects

Bernhard Holzer
CERN

A Short Introduction ... LOL

*In the end and after all ... : We try to explain the structure of the “hadronic matter” in the Universe.
In other words: What is going on up there ???*

1869

PERIODENSYSTEM DER ELEMENTE

<http://www.kfz-split.hr/periodni/de/>

Legend:

- Metalle (Metals)
- Alkalimetalle (Alkali metals)
- Erdalkalimetalle (Alkaline earth metals)
- Ubergangselemente (Transition elements)
- Lanthaniden (Lanthanides)
- Actiniden (Actinides)
- Halbmetalle (Metalloids)
- Nichtmetalle (Non-metals)
- Chalkogene (Chalcogens)
- Halogene (Halogens)
- Edelgase (Noble gases)

ZUSTAND (100 °C, 101 kPa):

- Ne - gasförmig (gaseous)
- Fe - fest (solid)
- Ga - flüssig (liquid)
- To - künstliche (synthetic)

PERIODENSYSTEM DER ELEMENTE (1869)

1	2	13	14	15	16	17	18											
1	H						He											
2	Li	Be					B	C	N	O	F	Ne						
3	Na	Mg								Al	Si	P	S	Cl	Ar			
4	K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
5	Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe
6	Cs	Ba	La-Lu	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn
7	Fr	Ra	Ac-Lr	Rf	Db	Sg	Bh	Hs	Mt	Uub	Uuq	Uub	Uuq	Uub	Uuq	Uub	Uuq	Uub

LANTHANIDEN

57	58	59	60	61	62	63	64	65	66	67	68	69	70	71
La	Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu

ACTINIDEN

89	90	91	92	93	94	95	96	97	98	99	100	101	102	103
Ac	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr

Redaktion: Marc Hens (mhens@gmx.de)

$$E=mc^2, \lambda=h/p$$




1.) Where are we ?

- * Standard Model of HEP***
- * Higgs discovery***

... and why all that ??

High Light of the HEP-Year 2012 / 13 naturally the HIGGS



ATLAS event display: Higgs => two electrons & two muons

$$E = m_0c^2 = m_{e1} + m_{e2} + m_{\mu1} + m_{\mu2} = 125.4 \text{ GeV}$$

2.) Where do we go ?

- * Physics beyond the Standard Model***
- * Dark Matter / Dark Energy***

Physics *Beyond the Standard Model (BSM)*

Example: *Dark Matter*

The outer region of galaxies rotate faster than expected from visible matter


$$\frac{mv^2}{r} = \frac{mMg}{r^2} \longrightarrow v_{\text{cric}} = \sqrt{\frac{Mg}{r}}$$

Dark matter would explain this

*Other observations exist ... (grav. lens effects)
but all through gravity*

What is it?

Budget: *Dark Matter: 26 %
Dark Energy: 70 %
Anything else (including us) 4 %*



European Strategy Group

Future High Energy Frontier Colliders

Luminosity Upgrade of LHC:
HL-LHC

Circular colliders:

FCC (Future Circular Collider)

***FCC-hh**: 100 TeV proton-proton cm energy*

***FCC-ee**: 90-350 GeV lepton collider*

Linear colliders

ILC (International Linear Collider): e^+e^- , 500 GeV cms energy,

***CLIC** (Compact Linear Collider): e^+e^- , 380GeV - 3TeV cms energy,*

Others

ERLs

Muon collider,

Plasma acceleration

A Bit of Theory

The big storage rings

„Synchrotrons“

$$D_n = \beta_C \sin n\phi_C * \delta_{\text{supr}} * \sum_{i=1}^n \cos\left(i\phi_C - \frac{1}{2}\phi_C \pm \varphi_m\right) * \sqrt{\frac{\beta_m}{\beta_C}} -$$

$$- \cos n\phi_C * \delta_{\text{supr}} * \sum_{i=1}^n \sqrt{\beta_m \beta_C} * \sin\left(i\phi_C - \frac{1}{2}\phi_C \pm \varphi_m\right)$$

$$D_n = \sqrt{\beta_m \beta_C} * \sin n\phi_C * \delta_{\text{supr}} * \sum_{i=1}^n \cos\left((2i-1)\frac{\phi_C}{2} \pm \varphi_m\right) -$$

$$- \sqrt{\beta_m \beta_C} * \delta_{\text{supr}} * \cos n\phi_C \sum_{i=1}^n \sin\left((2i-1)\frac{\phi_C}{2} \pm \varphi_m\right)$$

Remembering the trigonometric gymnastics shown above we get

$$D_n = \delta_{\text{supr}} * \sqrt{\beta_m \beta_C} * \sin n\phi_C * \sum_{i=1}^n \cos\left((2i-1)\frac{\phi_C}{2}\right) * 2 \cos\varphi_m$$

$$- \delta_{\text{supr}} * \sqrt{\beta_m \beta_C} * \cos n\phi_C * \sum_{i=1}^n \sin\left((2i-1)\frac{\phi_C}{2}\right) * 2 \cos\varphi_m$$

$$D_n = 2\delta_{\text{supr}} * \sqrt{\beta_m \beta_C} * \cos\varphi_m \left\{ \sin(n\phi_C) - \sum_{i=1}^n \sin\left((2i-1)\frac{\phi_C}{2}\right) * \cos(n\phi_C) \right\}$$

$$\delta_{\text{supr}} * \sqrt{\beta_m \beta_C} * \cos\varphi_m \sin(n\phi_C) \frac{\sin \frac{n\phi_C}{2} * \cos \frac{n\phi_C}{2}}{\sin \frac{\phi_C}{2}} -$$

$$- 2\delta_{\text{supr}} * \sqrt{\beta_m \beta_C} * \cos\varphi_m * \cos(n\phi_C) * \frac{\sin \frac{n\phi_C}{2} * \sin \frac{n\phi_C}{2}}{\sin \frac{\phi_C}{2}}$$

What

we

will

not

do

replace by ...

“after some TLC Transformations”

... or ... “after some beer”

$$- (\cos^2 \frac{n\phi_C}{2} - \sin^2 \frac{n\phi_C}{2}) \sin^2 \frac{n\phi_C}{2}$$


1.) Geometry of a Storage Ring:

A charged particle in a magnetic dipole field feels a transverse deflecting force, The „Lorentz force“

$$\cancel{E = mc^2} \quad E^2 = (pc)^2 + m^2c^4$$

We have to calculate relativistically, which is not so difficult and leads to $E \leftrightarrow p$

Condition for an ideal circular orbit:



Lorentz force

$$F_L = e v B$$

centrifugal force

$$F_{centr} = \frac{\gamma m_0 v^2}{\rho}$$

$$p = \gamma \cdot m_0 v$$

$$\frac{\cancel{\gamma m_0 v^2}}{\rho} = \cancel{e v B}$$

$$\frac{p}{e} = B \rho$$

$B \rho =$ "beam rigidity"

The overall integral of all dipole fields around the ring has to give 2π bending angle

2.) Focusing Forces: Hook's law


*... keeping the flocs together:
In addition to the pure bending of the beam
we have to keep 10^{11} particles close together*



focusing force



*And here we borrow the idea from classical mechanics:
The pendulum*



*there is a **restoring force**, proportional
to the elongation x :*

$$F = m * a = - \text{const} * x$$

$$F = m * \frac{d^2x}{dt^2} = - \text{const} * x$$

*general solution:
free harmonic oscillation*

$$x(t) = A * \cos(\omega t + \varphi)$$


...this is how grandma's Kuckuck's clock is working!!!

Dipoles:

Create a constant field $B_y = \text{const}$

Quadrupoles:

Create a linear increasing magnetic field $B_y = g \cdot x, \quad B_x = g \cdot y$



A linear increasing restoring force leads always (!) to a harmonic oscillation.

==> quadrupoles do that for us



And dipoles define the particle momentum

$$B_y = g \cdot x$$

$$B_y = \text{const}$$


The movement of a charged particle in the “Lattice” of external magnetic fields can be described analytically.

... and corresponds - in linear fields - to a harmonic transverse oscillation.



Sesame Light Source

We can calculate the single particle trajectories for an arbitrary number of turns.





*typical values
in a strong
foc. machine:
 $x \approx \text{mm}$, $x' \leq \text{mrad}$*

LHC Operation: Beam Commissioning

First turn steering "by sector:"


- One beam at the time
- Beam through 1 sector (1/8 ring), correct trajectory, open collimator and move on.



Many particles: The Beam

$$x(s) = \sqrt{\varepsilon} \sqrt{\beta(s)} \cdot \cos(\Psi(s) + \phi)$$

$$\hat{x}(s) = \sqrt{\varepsilon} \sqrt{\beta(s)}$$



single particle trajectories, $N \approx 10^{11}$ per bunch


Beta-Function describing the size of the Particle Ensemble

Gauß Particle Distribution:
$$\rho(x) = \frac{N \cdot e}{\sqrt{2\pi}\sigma_x} \cdot e^{-\frac{1}{2}\frac{x^2}{\sigma_x^2}}$$


particle at distance 1σ from centre
 \leftrightarrow 68.3 % of all beam particles

LHC:

$$\sigma = \sqrt{\varepsilon * \beta} = \sqrt{5 * 10^{-10} m * 180 m} = 0.3 mm$$



aperture requirements: $r_0 = 17 * \sigma$




ATTENTION: its classical mechanics

Beam Dynamics in a Storage Ring


The particle movement described in

phase space, x, x'


—> plot x, x' as a function of „s“



X



X'





Theorem of Liouville

... and now the ellipse:

note for each turn x , x' at a given position „s“ and plot in the phase space diagram

under the influence of conservative forces, the particle kinematics will always follow an ellipse in phase space x , x' phase space volume = constant

We use the area of that beam-ellipse as quality attribute for the particle ensemble: $A = \varepsilon \pi$



Time for a blue Slide ...

Why do we do that ?

—> *the beam size is given by two parameters:
β function - focusing properties
ε as intrinsic beam quality*

—> *beam size:* $\sigma = \sqrt{\varepsilon \cdot \beta}$


—> *the stability of the phase space ellipse, ε,
tells us about the stability
of the particle oscillation, which is ...
... “the lifetime” of the beam.*

—> *the size of the ellipse tells us about the particle density,
... which is the beam quality in collision.*



Particle Density in matter

Hydrogen molecule




Atomic Distance in Hydrogen Molecule


$$R_B \approx 0.5 \text{ \AA}$$

in solids / fluids $\lambda \approx 2.1 \dots 2.9 \text{ \AA}$

in gases $\lambda \approx 33 \text{ \AA} = 3.3 \text{ nm}$





Particle Distance in Accelerators: $\lambda \approx 6000 \text{ \AA} = 600 \text{ nm}$ (Arc LHC)



Problem: Our particles are *VERY* small !!

Overall cross section of the Higgs:




$$1 \text{ b} = 10^{-24} \text{ cm}^2$$

$$1 \text{ pb} = 10^{-12} \cdot 10^{-24} \text{ cm}^2 = 1/\text{mio} \cdot 1/\text{mio} \cdot 1/\text{mio} \cdot 1/\text{mio} \cdot 1/\text{mio} \cdot 1/10000 \text{ mm}^2$$

*The only chance we have:
compress the transverse beam size ... at the IP*

The particles are “very small”




LHC typical:

$$\sigma = 0.1 \text{ mm} \rightarrow 16 \mu\text{m}$$

Luminosity

$$R = L * \Sigma_{react}$$



Example: Luminosity run at LHC

$$\beta_{x,y} = 0.55 \text{ m}$$

$$f_0 = 11.245 \text{ kHz}$$

$$\varepsilon_{x,y} = 5 * 10^{-10} \text{ rad m}$$

$$n_b = 2808$$

$$\sigma_{x,y} = 16 \text{ }\mu\text{m}$$

$$I_p = 584 \text{ mA}$$


$$L = \frac{1}{4\pi e^2 f_0 n_b} * \frac{I_{p1} I_{p2}}{\sigma_x \sigma_y}$$

$$L = 1.0 * 10^{34} \text{ } 1/\text{cm}^2 \text{ s}$$

*Every future collider has to push for highest possible luminosity
... and energy.*

3.) *The HL-LHC*

- * increasing the luminosity of LHC*
- * higher bunch intensities*
- * smaller β^**




3.) The HL-LHC



- * increasing the luminosity of LHC
- * higher bunch intensities
- * smaller β^*

	<i>LHC</i>	<i>HL-LHC</i>
<i>Energy</i>	7 TeV	7 TeV
<i>Particles / bunch</i>	$1.2 \cdot 10^{11}$	$2.2 \cdot 10^{11}$
<i>number of bunches</i>	2808	2748
β^*	55 cm	15 cm
ε	$5.0 \cdot 10^{-10} \text{ m rad}$	$3.3 \cdot 10^{-10} \text{ m rad}$
σ	16 μm	7 μm
<i>Luminosity</i>	$1.0 \cdot 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$	$7.0 \cdot 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$

LHC & HL-LHC



At one (or a very few) points in the accelerator, we make the beams as small as possible, to push for highest particle density.





Mini-Beta Insertion

B. J. Holzer, CERN

Theorem of Liouville

... area of phase space ellipse is constant !!!

kleine Strahldimension < = = > grosse Divergenz




Challenge: High Field Nb₃Sn Quad

Stronger focusing needs stronger magnets

We need a material that can withstand this higher field in its super conducting phase !!! Nb₃Sn

LQS01b Quench History



reminder: LHC standard inner triplet NbTi: G=215 T/m, $\Phi=66$ mm



The Luminosity defines the number of "hits". It depends on the particle density at the collision point.

The Beta function at the IP " β^* " should be made as small as possible to increase the particle density. In a drift β is growing quadratically and proportional to $1/\beta^*$, which sets the ultimate limit to the achievable luminosity.

The distance L^* of the focusing magnets from the IP should be as small as possible.

... try to avoid detectors like ATLAS or CMS whenever possible. LOL.

The beam dimensions at the IP are typically a few μm .

What is a μm ?



beam sizes in the order of your cat's hair !!

4.) Push for higher energy: FCC

- * increasing the ring size***
- * stronger magnets***

Future Colliders: Hadrons or Leptons?

Hadron collisions: compound particles



Proton = $u+u+d$ + gluons + sea-quarks

Mix of quarks, anti-quarks and gluons

→ variety of processes

Parton energy spread

Hadron collisions ⇒ **large discovery range**



LHC Pb-Pb collision (Atlas)

Lepton collisions: Elementary particles / Anti-particles


Collision process known

Well defined energy

Other physics background limited

Lepton collisions ⇒ precision measurements


in $e^+ e^-$ collisions **quantum numbers disappear**



PETRA: gluon discovery



The Next Generation Ring Collider



FCC

LHC

PS


SPS

Maximum Beam Energy in a Storage Ring:

For a given magnet technology it is the size of the machine that defines the maximum particle momentum ... and so the energy

~~$$E = mc^2$$~~

$$E^2 = (pc)^2 + m^2c^4$$



Condition for an ideal circular orbit:

Lorentz force

$$F_L = e v B$$

centrifugal force

$$F_{centr} = \frac{\gamma m_0 v^2}{\rho}$$

$$p = \gamma \cdot m_0 v$$


~~$$\frac{\gamma m_0 v^2}{\rho} = e v B$$~~

$$\frac{p}{e} = B \rho$$

$B \rho =$ "beam rigidity"


The maximum particle momentum is given by the field strength B and the storage ring size $2\pi\rho$

The Push for Higher Beam Energy



NbTi LHC standard dipoles,
8.3 T

Nb₃Sn FCC type dipole coils,
11 T – 16 T



FCC energy reach:

it is a simple scaling wrt LHC:

circumference 100km /27km

→ Factor 3.7

dipole field: 16 T / 8.3 T

→ Factor 1.93

LHC: $E_{cm} = 2 * 7 \text{ TeV} = 14 \text{ TeV}$

FCC: $E_{cm} = 100 \text{ TeV}$ centre of mass

FCC-hh Parameter List

Pushing the limit (Dipole Fill Factor):


12 dipoles per cell, $l_{dipole} = 14.2m$

34 cells per arc

12 arcs

dipole field = 16T \leftrightarrow 50TeV

} 5016 dipoles




	LHC		HL-LHC		FCC-hh	
					Initial	Nominal
Main parameters and geometrical aspects						
c.m. Energy (TeV)		14			100	
Circumference C (km)		26.7			97.75	
Dipole field (T)		8.33			<16	
Physics performance and beam parameters						
Peak luminosity ¹ ($10^{34} \text{ cm}^{-2} \text{ s}^{-1}$)	1.0		5.0		5.0	<30.0
Beam parameters						
Number of bunches n			2808			10 400
Bunch spacing (ns)	25		25			25
Bunch population $N(10^{11})$	1.15		2.2			1.0
RMS bunch length ² (cm)			7.55			8
IP beta function (m)	0.55		0.15 (min)		1.1	0.3
RMS IP spot size (μm)	16.7		7.1 (min)		6.8	3.5
Full crossing angle (μrad)	285		590		104	200 ³
Other beam and machine parameters						
Stored energy per beam (GJ)	0.362		0.694			8.3
SR power per ring (MW)	0.0036		0.0073			2.4

LHC Operation:

Machine Protection & Safety

Energy stored in magnet system	10	GJ
Energy stored in one main dipole circuit	1.1	GJ
Energy stored in one beam	362	MJ

Enough to melt 500 kg of copper



$2 \cdot 10^{12}$ $4 \cdot 10^{12}$ $8 \cdot 10^{12}$ $6 \cdot 10^{12}$

450 GeV p Beam

5.) High Energy Lepton Colliders

- * Limited by Synchrotron Radiation***
- * and RF Power***



The next Generation e^+/e^- Ring Collider



Synchrotron Radiation



ca 400 000 v. Chr.: Mankind discovers the Fire

Synchrotron Radiation

In a circular accelerator *charged particles loose energy via emission of intense light.*

$$P_s = \frac{2}{3} \alpha \hbar c^2 \frac{\gamma^4}{\rho^2} \quad \text{radiation power}$$



$$\Delta E = \frac{4}{3} \pi \alpha \hbar c \frac{\gamma^4}{\rho} \quad \text{energy loss}$$

$$\omega_c = \frac{3}{2} \frac{c \gamma^3}{\rho} \quad \text{critical frequency}$$

$$\alpha \approx \frac{1}{137}$$

$$\hbar c \approx 197 \text{ MeV fm}$$

1946 observed for the first time in the General Electric Synchrotron



court. K. Wille

FCC-ee: a collider that is dominated

by synchrotron light losses.

→ Planning the next generation e^+ / e^- Ring Colliders means build it LARGE.



$$\Delta U_0 (keV) \approx \frac{89 * E^4 (GeV)}{\rho}$$

$$\Delta U_0 \approx 8.62 GeV$$

$$\Delta P_{sy} \approx \frac{\Delta U_0}{T_0} * N_p = \frac{10.4 * 10^6 eV * 1.6 * 10^{-19} Cb}{263 * 10^{-6} s} * 9 * 10^{12}$$

$\Delta P_{sy} \approx 47 MW$... per beam

Circular e^+ / e^- colliders are severely limited by synchrotron radiation losses and have to be replaced for higher energies by linear accelerators



220 A × 64 turns

FCC-ee Parameters

	Z	WW	ZH	tt	
Circumference [km]			97.756		
Bending radius [km]			10.760		
Free length to IP l^* [m]			2.2		
Solenoid field at IP [T]			2.0		
Full crossing angle at IP θ [mrad]			30		
SR power / beam [MW]			50		
Beam energy [GeV]	45.6	80	120	175	182.5
Beam current [mA]	1390	147	29	6.4	5.4
Bunches / beam	16640	2000	328	59	48
Bunch population [10^{11}]	1.7	1.5	1.8	2.2	2.3
Horizontal emittance ϵ_x [nm]	0.27	0.84	0.63	1.34	1.46
Vertical emittance ϵ_y [pm]	1.0	1.7	1.3	2.7	2.9
Horizontal β_x^* [m]	0.15	0.2	0.3	1.0	
Vertical β_y^* [mm]	0.8	1.0	1.0	1.6	
Luminosity / IP [$10^{34}/\text{cm}^2 \text{ s}$]	230	28	8.5	1.8	1.55

*For a given particle energy
the beam intensity will be
limited by the maximum
tolerable Synchrotron
radiation power loss*

*RF Voltage applied depends on the
beam energy as $U \propto \gamma^4$*

$$\Delta U(\text{keV}) = 89 * \frac{E^4 / (mc^2)^4}{\rho}$$


6.) Push for higher lepton energy

**** go linear***

**** higher acceleration gradients***

CLIC ... a future Linear e^+ / e^- Accelerator

„C“-LIC ... = CERN ... or „compact“



← 50 km →

Description [units]	500 GeV	3 TeV
Total (peak 1%) luminosity	$2.3 (1.4) \times 10^{34}$	$5.9 (2.0) \times 10^{34}$
Total site length [km]	13.0	48.4
Loaded accel. gradient [MV/m]	80	100
Main Linac RF frequency [GHz]		12
Beam power/beam [MW]	4.9	14
Bunch charge [$10^9 e^+ / e^-$]	6.8	3.72
Bunch separation [ns]		0.5
Bunch length [μm]	72	44
Beam pulse duration [ns]	177	156
Repetition rate [Hz]		50
Hor./vert. norm. emitt. [$10^{-6} / 10^{-9} \text{m}$]	2.4/25	0.66/20
Hor./vert. IP beam size [nm]	202/2.3	40/1

CLIC parameter list

CLIC: Normal conducting RF system



challenge: running at the break down limit

Acceleration Gradient **100MV/m** studied & optimised since years




“ how far can we go and how much can we optimise such a future accelerator before we reach technical limits and how can we push these limits ? ”

they have impact on

- => the accelerator performance (luminosity)
- => beam quality
- => and the accelerating structure itself



Plasma Wake Acceleration




*Excite a plasma wave
with an intense LASER pulse*

*—> create acc. fields of
 $E > \text{GV/m}$*


AWAKE:

Proton driven Wake Acceleration Experiment at CERN

Prototype: 1m long Rb Plasma Cell



*Eupraxia: European PWA accelerator
INFN, Frascati*



Study of High Gradient Acceleration Techniques

Plasma Wake Acceleration


particle beam driven / LASER driven

Incoming laser pulse (or pulse of particles) **creates a travelling plasma wave** in a low-pressure gas

Plasma wake **field gradient accelerates electrons** that 'surf' on the plasma wave

S. Corde et al

Field Gradients up to 100 GeV/m observed




Plasma cell Univ. Texas, Austin

$E_e = 2 \text{ GeV}$

B. J. Holzer, CERN

$$\begin{aligned} \Delta E / \Delta s &= 200 \text{ MeV} / 4 \text{ mm} \\ &= 50 \text{ GeV} / \text{m} \end{aligned}$$

Netzwerk Teilchenwelt '23



Open questions in particle physics

Dark matter & Energy

... on which energy scale to look for it ?

Physics beyond the standard model

... Lepton or Proton colliders ?

Beam dynamics aspects

... Circular or linear ?


Technical aspects

... Traditional, sc / nc or PWA ?




Merci

ATLAS detector in LHC




Energy stored in the magnet

in case of problems ...
react fast and safe !!



(A. Gómez)




During magnet

Quench in a LHC magnet

failure of nc. dipole D1: $\tau_{\text{damage}} = 6.4 \text{ ms}$

damage level after 20 turns !!






Challenge: HL-LHC Crab Cavities

$$L = L_{ideal} \cdot F$$

$$F = \frac{1}{\sqrt{1 + 2 \frac{\sigma_s^2}{\sigma_{1x}^2 + \sigma_{2x}^2} \tan^2 \frac{\phi}{2}}}$$


F = Spaghetti-Loss Factor



Transverse deflecting cavity at 800 MHz

Prototype test in SPS ... at the moment technical challenge:

*fast, precise, compact,
Fail SAFE !!*



Circular vs. Linear Colliders

... the light problem

F. Gianotti

