

Mueller Tang jets in next-to-leading BFKL

D. Colferai^{1,2}, F. Deganutti³, T. Raben³, C. Royon³



STRONG-2020
GA 824093

¹University of Florence, Italy



²INFN Florence, Italy



³Kansas University, Kansas



JHEP 06 (2023) 091 [[hep-ph 2304.09073](https://arxiv.org/abs/2304.09073)]

MPI@LHC

Manchester

26.11.2023

Outline


- Introduction:
 - Definition and motivation of Mueller-Tang jets
 - PT description in QCD: LL BFKL factorization formula
- Previous phenomenological studies
 - MTJ in LL approx (and NLL GGF)
 - → need of a full NLL calculation
- NLL impact factors
 - Structure of NLL impact factor
 - Implementation of NLL impact factors: numerical and conceptual issues
 - (Small) breaking of BFKL factorization at NLL level
- Numerical results
- Conclusions and outlook

Mueller-Tang jets

An important process for studying PT high-energy QCD and the Pomeron at hadron colliders [Mueller, Tang '87]

Final state:

- two jets with similar p_T
- large rapidity distance $Y \simeq \log(s/p_T^2)$;
- absence of any additional emission in central rapidity region (gap)





Mueller-Tang jets

An important process for studying PT high-energy QCD and the Pomeron at hadron colliders [Mueller, Tang '87]


Final state:

- two jets with similar p_T
 - large rapidity distance $Y \simeq \log(s/p_T^2)$;
 - absence of any additional emission in central rapidity region (gap)
- Gap \implies mostly colour-singlet exchanges contribute to cross section
 - $Y \gg 1 \implies$ enhanced PT series $(\alpha_S Y)^n$ resummed into singlet BFKL GGF
 - In LLA factorization formula holds





Mueller-Tang jets at LO and LL

- LO amplitude: box + crossed diagrams
projected onto colour-singlet
 $\Pi^{ab,a'b'} = \delta^{ab}\delta^{a'b'} / (N_c^2 - 1)$






Mueller-Tang jets at LO and LL

- LO amplitude: box + crossed diagrams projected onto colour-singlet
 $\Pi^{ab,a'b'} = \delta^{ab}\delta^{a'b'} / (N_c^2 - 1)$
- Elastic amplitude at higher orders: affected by large $\log^n s$ due to gluon-ladder diagrams (UV and IR finite)
- All LL resummed in (colour-singlet) gluon Green function (GGF)




Mueller-Tang jets at LO and LL

- LO amplitude: box + crossed diagrams projected onto colour-singlet
 $\Pi^{ab,a'b'} = \delta^{ab}\delta^{a'b'} / (N_c^2 - 1)$
- Elastic amplitude at higher orders: affected by large $\log^n s$ due to gluon-ladder diagrams (UV and IR finite)
- All LL resummed in (colour-singlet) gluon Green function (GGF)
- LL partonic cross section: 2 GGF * 2 (trivial) impact factors
- Two outgoing partons to be identified with the (back-to-back) jets




CMS analysis at 7 TeV




- Charged-particle multiplicity in the gap region between the tagged jets compared to PYTHIA and HERWIG predictions.
- HERWIG 6: include contributions from color singlet exchange (CSE), based on **BFKL at LL**.
- PYTHIA 6: inclusive dijets (tune Z2*), **no-CSE**.

D0 and CMS analysis at 7 TeV



Left: LL & NLL BFKL at Tevatron [hep-ph/1012.3849].


- Ratio $R = \frac{NLL^* BFKL}{NLO QCD}$ of jet-gap-jet events to inclusive dijet events as a function of p_t and the rapidity gap Y .



NLL* BFKL calculations different implementations of the soft rescattering processes (EEI models), describe many features of the data, but none of the implementations is able to simultaneously describe all the features of the measurement. [Ekstedt, Enberg, Ingelman, \[1703.10919\]](https://arxiv.org/abs/1703.10919)


NL impact factors

- Compelling to include all NLL corrections into the game




NL impact factors

- Compelling to include all NLL corrections into the game




- Idea: generalize MT factorization formula at NLL
- BFKL GGF at NLL known since long [*Fadin, Fiore et al.*]





NL impact factors

- Compelling to include all NLL corrections into the game



- Idea: generalize MT factorization formula at NLL
- BFKL GGF at NLL known since long [*Fadin, Fiore et al.*]
- NL impact factors determined by NLO calculation, with IR (soft and collinear) divergencies



Not a trivial statement:

- all $\log(s)$ terms must reproduce LL kernel (GGF at 1st order)
- all IR singularities (taken away collinear ones proportional to splitting functions) must cancel

NL impact factors

$$\begin{aligned}
 \Phi(l_1, l_2, \mathbf{q}) &= \frac{\alpha_S^3}{2\pi(N_C^2 - 1)} \int_0^1 dz \int d^2\mathbf{k} \\
 &\times \mathbf{S}_J(\mathbf{k}, \mathbf{q}, z) C_F \frac{1 + (1-z)^2}{z} \\
 &\times \left\{ C_F^2 \frac{z^2 \mathbf{q}^2}{\mathbf{k}^2 (\mathbf{k} - z\mathbf{q})^2} + C_F C_A f_1(l_{1,2}, \mathbf{k}, \mathbf{q}, z) + C_A^2 f_2(l_{1,2}, \mathbf{k}, \mathbf{q}) \right\}
 \end{aligned}$$

- The calculation of NL impact factors for Mueller-Tang jets was performed by [\[Hentchinski, Madrigal Martinez, Murdaca, Sabio Vera, '14\]](#) using Lipatov's effective action (and confirmed by F.Deganutti and myself)


NL impact factors

$$\begin{aligned}
 \Phi(l_1, l_2, \mathbf{q}) &= \frac{\alpha_S^3}{2\pi(N_c^2 - 1)} \int_0^1 dz \int d^2\mathbf{k} \\
 &\times \mathbf{S}_J(\mathbf{k}, \mathbf{q}, z) C_F \frac{1 + (1-z)^2}{z} \\
 &\times \left\{ C_F^2 \frac{z^2 \mathbf{q}^2}{\mathbf{k}^2(\mathbf{k} - z\mathbf{q})^2} + C_F C_A f_1(l_{1,2}, \mathbf{k}, \mathbf{q}, z) + C_A^2 f_2(l_{1,2}, \mathbf{k}, \mathbf{q}) \right\}
 \end{aligned}$$

- The calculation of NL impact factors for Mueller-Tang jets was performed by [\[Hentchinski, Madrigal Martinez, Murdaca, Sabio Vera, '14\]](#) using Lipatov's effective action (and confirmed by F.Deganutti and myself)
- Phase-space integration restricted by **IR-safe jet algorithm** (e.g., $kt \simeq \text{cone}$)
- The two partons in the same hemisphere form (at least) one jet:
 - $\Delta\Omega \equiv \sqrt{\Delta y^2 + \Delta\phi^2} < R \implies J = \{qg\}$ composite jet
 - $\Delta\Omega > R \implies J = \{g\}$ and q outside jet cone or $J = \{q\}$ and g outside

Problem with NL impact factor


$$\Phi(l_1, l_2, \mathbf{q}) = \frac{\alpha_S^3}{2\pi(N_C^2 - 1)} \int_0^1 dz \int d^2 \mathbf{k} \\ \times \mathbf{S}_J(\mathbf{k}, \mathbf{q}, z) C_F \frac{1 + (1-z)^2}{z} \\ \times \left\{ C_F^2 \frac{z^2 \mathbf{q}^2}{k^2 (\mathbf{k} - z\mathbf{q})^2} + C_F C_A f_1(l_{1,2}, \mathbf{k}, \mathbf{q}, z) + C_A^2 f_2(l_{1,2}, \mathbf{k}, \mathbf{q}) \right\}$$



- There is a **problem** in the C_A^2 term, due to $\int_0^1 dz/z$ integration
- If integration is not constrained, we have a divergence

Problem with NL impact factor


$$\Phi(I_1, I_2, \mathbf{q}) = \frac{\alpha_S^3}{2\pi(N_C^2 - 1)} \int_0^1 dz \int d^2\mathbf{k} \\ \times \mathbf{S}_J(\mathbf{k}, \mathbf{q}, z) C_F \frac{1 + (1-z)^2}{z} \\ \times \left\{ C_F^2 \frac{z^2 \mathbf{q}^2}{\mathbf{k}^2 (\mathbf{k} - z\mathbf{q})^2} + C_F C_A f_1(I_{1,2}, \mathbf{k}, \mathbf{q}, z) + C_A^2 f_2(I_{1,2}, \mathbf{k}, \mathbf{q}) \right\}$$



- There is a **problem** in the C_A^2 term, due to $\int_0^1 dz/z$ integration
- If integration is not constrained, we have a divergence
- Such region $z \rightarrow 0$ corresponds to gluon in central (and backward) region, where the emission probability of the gluon turns out to be flat in rapidity: $\int_0^1 dz/z = \int_{-\infty}^{\log \sqrt{s}/k} dy = \infty$!
- If we believe the IF calculation to be reliable at least in the forward hemisphere ($y > 0$) $\implies \int_0^{\log \sqrt{s}/k} dy = \int_{k/\sqrt{s}}^1 dz/z = \frac{1}{2} \log(s/k^2)$
- But a $\log(s)$ in IFs is **not acceptable** within the spirit of BFKL factorization


Violation of BFKL factorization

- What happens for MT jets? The theoretical argument:
"colour-singlet momentum transfer \implies no $\log s$
is **wrong**"
- Here colour-singlet either below or above
 $\implies \log s$ unavoidable without constraints



Violation of BFKL factorization

- What happens for MT jets? The theoretical argument: "colour-singlet momentum transfer \Rightarrow no log s is **wrong**



- Here colour-singlet either below or above



\Rightarrow log s unavoidable without constraints


- MT event selection constrains particles not to be emitted within the gap provided they are above some energy threshold E_{th} (cal resolution)
- Only particles below threshold can be emitted at any rapidity
- This prescription is IR safe because inclusive for $E_g < E_{th}$

But gluons below threshold can have **any rapidity** $\Rightarrow \sigma \ni C_A^2 \frac{E_{th}^2}{E_J^2} \log \frac{s}{E_J^2}$

With such "minimal" experimental prescription, **BFKL factorization is violated** (impact factors depend on s). However **violation** is expected to be **small**.

Results

- We go on and use NL IF and GGF for $d\sigma/dY$
- Numerical implementation requires 10-dimensional integrals of singular functions



Leading VS Next-to-leading cross section


We follow CMS setup:

- $\sqrt{s} = 14$ TeV
- $E_J \geq 40$ GeV
- $1.5 \leq |y_J| \leq 5$
- $3 \leq Y \equiv y_{J1} - y_{J2} \leq 9$
- $y_{\text{gap}} \in [-1, 1] \rightarrow \Delta Y_{\text{gap}} = 2$
- $E_{\text{thresh}} = 1.0$ GeV

Leading VS Next-to-leading cross section

We follow CMS setup:


- $\sqrt{s} = 14$ TeV
- $E_J \geq 40$ GeV
- $1.5 \leq |y_J| \leq 5$
- $3 \leq Y \equiv y_{J1} - y_{J2} \leq 9$
- $y_{\text{gap}} \in [-1, 1] \rightarrow \Delta Y_{\text{gap}} = 2$
- $E_{\text{thresh}} = 1.0$ GeV
- NLL corrections of the impact factors are negative
- $\sigma_{\text{full}} \lesssim \sigma_{LL}$
with a slightly steeper decrease



Factorization violation and dependence on gap width

Contribution of the term $C_A^2 \log \frac{s}{E^2}$
that violates factorization:

- Violation of factorization is small, $\sim 10\%$ (with $Y_{\text{gap}} = 2$)
- Resummation of such logarithms not necessary for phenomenology
- Cross section slightly increases while decreasing ΔY_{gap} and saturates with no gap
- Emission from singlet exchange in central region is dynamically suppressed




Renormalization scale dependence

Running coupling $\alpha_S(Q^2)$ at
physical scale $Q = \lambda(E_{J1} + E_{J2})$

Renormalization scale dependence

Running coupling $\alpha_S(Q^2)$ at
physical scale $Q = \lambda(E_{J1} + E_{J2})$


- Renorm scale dependence still large



Renormalization scale dependence

Running coupling $\alpha_S(Q^2)$ at
physical scale $Q = \lambda(E_{J1} + E_{J2})$


- Renorm scale dependence still large
- Need for scale fixing procedure (BLM, PMS, ...)



Renormalization scale dependence


Running coupling $\alpha_S(Q^2)$ at
physical scale $Q = \lambda(E_{J1} + E_{J2})$

- Renorm scale dependence still large
- Need for scale fixing procedure (BLM, PMS, ...)
- Minimum sensitivity reached at $\lambda = 4$



Final theoretical prediction

- Central value with PMS renorm scale fixing
- Total error from all sources μ_R, μ_F, s_0, MC combined in quadrature
- At NL level the theoretical uncertainty is much reduced
- Results are compatible with those of the LL approx.



Conclusions and outlook

- Complete numerical implementation of MT jets at LHC in NLLA with collinear resummation of BFKL kernel; cross section slightly lower and steeper than in LLA
- Gap survival probability and showering still to be taken into account
- Strictly speaking jet-gap-jet observable violates BFKL factorization in NLLA
- Nevertheless the violation is small and factorization formula is expected to work well for LHC (non-asymptotic) kinematics.
- Good stability w.r.t. gap/threshold parameters
- Better description expected with proper renorm scale fixing ($\simeq 4$ times larger than natural scale)

Conclusions and outlook

- Complete numerical implementation of MT jets at LHC in NLLA with collinear resummation of BFKL kernel; cross section slightly lower and steeper than in LLA
- Gap survival probability and showering still to be taken into account
- Strictly speaking jet-gap-jet observable violates BFKL factorization in NLLA
- Nevertheless the violation is small and factorization formula is expected to work well for LHC (non-asymptotic) kinematics.
- Good stability w.r.t. gap/threshold parameters
- Better description expected with proper renorm scale fixing ($\simeq 4$ times larger than natural scale)
- Improvements could include hadronization, resummation of $\log s$ term in IFs, and inclusion of gap survival probability

Additional slides

Description with Pomeron loop

- The Pomeron loop correspond to the gap region
- The triple Pomeron vertex is not known at NLO

