

# Constraining neutrino models predictions

Szymon Zięba

University of Silesia in Katowice

Based on arXiv:2310.20681, submitted to PPNP

## Phenomenology of Lepton Masses and Mixing with Discrete Flavor Symmetries

authors: Garv Chauhan, P. S. Bhupal Dev, Ievgen Dubovyk, Bartosz Dziewit, Wojciech Flieger, Krzysztof Grzanka, Janusz Gluza, Biswajit Karmakar, Szymon Zięba

XXX Cracow EPIPHANY Conference on Precision Physics at High Energy Colliders,  
Kraków, January 11, 2024



# Neutrino mixing, $3\nu$ .


Neutrino flavor and mass eigenstates are related by

$$|\nu_\alpha\rangle = U_{\alpha i} |\nu_i\rangle$$


Pontecorvo–Maki–Nakagawa–Sakata parametrization of mixing matrix

$$U_{\text{PMNS}} = \begin{bmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{i\delta} \\ s_{12}c_{23} - c_{12}s_{13}s_{23}e^{i\delta} & c_{12}c_{33} - s_{12}s_{13}s_{23}e^{i\delta} & c_{13}s_{23} \\ s_{12}s_{23} - c_{12}s_{13}c_{23}e^{i\delta} & -c_{12}s_{23} - s_{12}s_{13}c_{23}e^{i\delta} & c_{13}c_{23} \end{bmatrix} \begin{bmatrix} e^{i\alpha_1} & 0 & 0 \\ 0 & e^{i\alpha_2} & 0 \\ 0 & 0 & 1 \end{bmatrix},$$

where  $c_{ij} \equiv \cos \theta_{ij}$ ,  $s_{ij} \equiv \sin \theta_{ij}$ ,  $\delta \equiv \delta_{\text{CP}}$ .



(a) CKM quark flavor mixing



(b) PMNS lepton flavor mixing

Taken from [arXiv:2210.11922](https://arxiv.org/abs/2210.11922), Figure 2.

Mass ordering,  $m_0 = m_{\text{lightest}}$ .

Normal mass ordering (NO)

$$m_1 = m_0,$$

$$m_2 = \sqrt{m_0^2 + \Delta m_{21}^2},$$


$$m_3 = \sqrt{m_0^2 + \Delta m_{31}^2},$$

Inverted mass ordering (IO)

$$m_1 = \sqrt{m_0^2 - \Delta m_{21}^2 - \Delta m_{32}^2},$$

$$m_2 = \sqrt{m_0^2 - \Delta m_{32}^2},$$

$$m_3 = m_0,$$



Taken from <https://globalfit.astroparticles.es>, updated arXiv:1806.11051, Figure 1

# Oscillation data, with SK.

Parameter	Ordering	NuFIT 5.2 (2022)		de Salas et al. (2021)		Capozzi et al. (2021)	
		bf $\pm 1\sigma$	3 $\sigma$ range	bf $\pm 1\sigma$	3 $\sigma$ range	bf $\pm 1\sigma$	3 $\sigma$ range
$\sin^2 \theta_{12} / 10^{-1}$	NO, IO	$3.03^{+0.12}_{-0.12}$	2.70 — 3.41	$3.18^{+0.16}_{-0.16}$	2.71 — 3.69	$3.03^{+0.13}_{-0.13}$	2.63 — 3.45
$\sin^2 \theta_{23} / 10^{-1}$	NO	$4.51^{+0.19}_{-0.16}$	4.08 — 6.03	$5.74^{+0.14}_{-0.14}$	4.34 — 6.10	$4.55^{+0.18}_{-0.15}$	4.16 — 5.99
$\theta_{23}$ octant	IO	$5.69^{+0.16}_{-0.21}$	4.12 — 6.13	$5.78^{+0.10}_{-0.17}$	4.33 — 6.08	$5.69^{+0.12}_{-0.21}$	4.17 — 6.06
$\sin^2 \theta_{13} / 10^{-2}$	NO	$2.225^{+0.056}_{-0.059}$	2.052 — 2.398	$2.200^{+0.069}_{-0.062}$	2.000 — 2.405	$2.23^{+0.07}_{-0.06}$	2.04 — 2.44
$\neq 0$	IO	$2.223^{+0.058}_{-0.058}$	2.048 — 2.416	$2.225^{+0.064}_{-0.070}$	2.018 — 2.424	$2.23^{+0.06}_{-0.06}$	2.03 — 2.45
$\delta_{CP} / \pi$	NO	$1.29^{+0.20}_{-0.14}$	0.80 — 1.94	$1.08^{+0.13}_{-0.12}$	0.71 — 1.99	$1.24^{+0.18}_{-0.13}$	0.77 — 1.97
can be 0?	IO	$1.53^{+0.12}_{-0.16}$	1.08 — 1.91	$1.58^{+0.15}_{-0.16}$	1.11 — 1.96	$1.52^{+0.15}_{-0.11}$	1.07 — 1.90
$\Delta m_{21}^2 / 10^{-5} \text{eV}^2$	NO, IO	$7.41^{+0.21}_{-0.20}$	6.82 — 8.03	$7.50^{+0.22}_{-0.20}$	6.94 — 8.14	$7.36^{+0.16}_{-0.15}$	6.93 — 7.93
$ \Delta m_{atm}^2  / 10^{-3} \text{eV}^2$	NO	$2.507^{+0.026}_{-0.027}$	2.427 — 2.590	$2.55^{+0.02}_{-0.03}$	2.47 — 2.63	$2.485^{+0.023}_{-0.031}$	2.401 — 2.565
	IO	$2.486^{+0.028}_{-0.025}$	2.406 — 2.570	$2.45^{+0.02}_{-0.03}$	2.37 — 2.53	$2.455^{+0.030}_{-0.025}$	2.376 — 2.541
$\Delta \chi^2$	IO - NO	6.4		6.4		6.4	




Figure taken from Peter B. Denton talk, [link here](#).




Figure taken from Biswajit Karmakar talk, [link here](#).

See also [arXiv:2012.12893](#), Figure 1 and [arXiv:2204.08668](#), Figure 2.1

BM, TB, GR, HG,  $\theta_{13} = 0$  (early 2010s).

$$U_{\text{PMNS}} = \begin{bmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{i\delta} \\ s_{12}c_{23} - c_{12}s_{13}s_{23}e^{i\delta} & c_{12}c_{s3} - s_{12}s_{13}s_{23}e^{i\delta} & c_{13}s_{23} \\ s_{12}s_{23} - c_{12}s_{13}c_{23}e^{i\delta} & -c_{12}s_{23} - s_{12}s_{13}c_{23}e^{i\delta} & c_{13}c_{23} \end{bmatrix}$$

$$\theta_{13} = 0^\circ \quad \Downarrow \quad \theta_{23} = 45^\circ$$

$$U_0 = \begin{bmatrix} \frac{c_{12}}{\sqrt{2}} & \frac{s_{12}}{\sqrt{2}} & 0 \\ -\frac{s_{12}}{\sqrt{2}} & \frac{c_{12}}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ -\frac{s_{12}}{\sqrt{2}} & \frac{c_{12}}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

$\Downarrow$

$$\theta_{12} = 45^\circ, s_{12} = 1/\sqrt{2}$$

$$\theta_{12} = 35, 26^\circ, s_{12} = 1/\sqrt{3}$$

$$\theta_{12} = 31, 7^\circ$$

$$\theta_{12} = 30^\circ, s_{12} = 1/2$$

Bimaximal Mixing

$$\begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ -\frac{1}{2} & \frac{1}{2} & \frac{1}{\sqrt{2}} \\ \frac{1}{2} & -\frac{1}{2} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

Tribimaximal Mixing

$$\begin{bmatrix} \sqrt{\frac{2}{3}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

Golden Ratio Mixing

$$\begin{bmatrix} \frac{\varphi}{\sqrt{2+\varphi}} & \frac{1}{\sqrt{2+\varphi}} & 0 \\ \frac{-1}{\sqrt{4+2\varphi}} & \frac{\varphi}{\sqrt{4+2\varphi}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{4+2\varphi}} & \frac{-\varphi}{\sqrt{4+2\varphi}} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

Hexagonal Mixing

$$\begin{bmatrix} \sqrt{\frac{3}{4}} & \frac{1}{2} & 0 \\ -\frac{1}{2\sqrt{2}} & \frac{\sqrt{3}}{2\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ -\frac{1}{2\sqrt{2}} & \frac{\sqrt{3}}{2\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

Golden Ratio Mixing:  $\tan \theta_{12} = 1/\varphi$ ,  $\varphi = (1 + \sqrt{5})/2$  being the golden ratio.

Based on Biswajit Karmakar talk, [link here](#).

# $\theta_{13} \neq 0$ , Daya Bay, RENO (2012).

BM, TB, GR, HG disfavored by non-zero  $\theta_{13}$ .

$$U_{\text{PMNS}} = \begin{bmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{i\delta} \\ s_{12}c_{23} - c_{12}s_{13}s_{23}e^{i\delta} & c_{12}c_{23} - s_{12}s_{13}s_{23}e^{i\delta} & c_{13}s_{23} \\ s_{12}s_{23} - c_{12}s_{13}c_{23}e^{i\delta} & -c_{12}s_{23} - s_{12}s_{13}c_{23}e^{i\delta} & c_{13}c_{23} \end{bmatrix}$$

$$\theta_{13} \neq 0^\circ \quad \Downarrow \quad \theta_{23} = 45^\circ$$

$$U_0 = \begin{bmatrix} c_{12} & s_{12} & 0 \\ -\frac{s_{12}}{\sqrt{2}} & \frac{c_{12}}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ -\frac{s_{12}}{\sqrt{2}} & \frac{c_{12}}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

$\Downarrow$

$$\theta_{12} = 45^\circ, s_{12} = 1/\sqrt{2}$$

$$\theta_{12} = 35, 26^\circ, s_{12} = 1/\sqrt{3}$$

$$\theta_{12} = 31, 7^\circ$$

$$\theta_{12} = 30^\circ, s_{12} = 1/2$$

Bimaximal Mixing

$$\begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ -\frac{1}{2} & \frac{1}{2} & \frac{1}{\sqrt{2}} \\ \frac{1}{2} & -\frac{1}{2} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

Tribimaximal Mixing

$$\begin{bmatrix} \sqrt{\frac{2}{3}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

Golden Ratio Mixing

$$\begin{bmatrix} \frac{\varphi}{\sqrt{2+\varphi}} & \frac{1}{\sqrt{2+\varphi}} & 0 \\ \frac{-1}{\sqrt{4+2\varphi}} & \frac{\varphi}{\sqrt{4+2\varphi}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{4+2\varphi}} & \frac{-\varphi}{\sqrt{4+2\varphi}} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

Hexagonal Mixing

$$\begin{bmatrix} \sqrt{\frac{3}{4}} & \frac{1}{2} & 0 \\ -\frac{1}{2\sqrt{2}} & \frac{\sqrt{3}}{2\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ -\frac{1}{2\sqrt{2}} & \frac{\sqrt{3}}{2\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

Golden Ratio Mixing:  $\tan \theta_{12} = 1/\varphi$ ,  $\varphi = (1 + \sqrt{5})/2$  being the golden ratio.

Based on Biswajit Karmakar talk, [link here](#).

# Non-zero $\theta_{13}$ : Successors of tribimaximal mixing, $\text{TM}_1$ , $\text{TM}_2$ .

$$U_{\text{TBM}} = \begin{bmatrix} \frac{2}{\sqrt{6}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \end{bmatrix}, \quad U_{\text{PMNS}} \simeq \begin{bmatrix} \frac{2}{\sqrt{6}} & \frac{1}{\sqrt{3}} & 0.15 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \end{bmatrix},$$

$$U_{\text{TM}_1} = \begin{bmatrix} \frac{2}{\sqrt{6}} & \frac{c_\theta}{\sqrt{3}} & \frac{s_\theta}{\sqrt{3}} e^{-i\gamma} \\ -\frac{1}{\sqrt{6}} & \frac{c_\theta}{\sqrt{3}} - \frac{s_\theta}{\sqrt{2}} e^{i\gamma} & -\frac{s_\theta}{\sqrt{3}} e^{-i\gamma} - \frac{c_\theta}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{c_\theta}{\sqrt{3}} - \frac{s_\theta}{\sqrt{2}} e^{i\gamma} & -\frac{s_\theta}{\sqrt{3}} e^{-i\gamma} + \frac{c_\theta}{\sqrt{2}} \end{bmatrix},$$

$$U_{\text{TM}_2} = \begin{bmatrix} \frac{2c_\theta}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{2s_\theta}{\sqrt{6}} e^{-i\gamma} \\ -\frac{c_\theta}{\sqrt{6}} + \frac{s}{\sqrt{2}} e^{i\gamma} & \frac{1}{\sqrt{3}} & -\frac{s_\theta}{\sqrt{3}} e^{-i\gamma} - \frac{c_\theta}{\sqrt{2}} \\ -\frac{c_\theta}{\sqrt{6}} + \frac{s}{\sqrt{2}} e^{i\gamma} & \frac{1}{\sqrt{3}} & -\frac{s_\theta}{\sqrt{3}} e^{-i\gamma} + \frac{c_\theta}{\sqrt{2}} \end{bmatrix},$$

$$|U_{\text{TM}_1}| = \begin{bmatrix} \frac{2}{\sqrt{6}} & * & * \\ \frac{1}{\sqrt{6}} & * & * \\ \frac{1}{\sqrt{6}} & * & * \end{bmatrix}, \quad |U_{\text{TM}_2}| = \begin{bmatrix} * & \frac{1}{\sqrt{3}} & * \\ * & \frac{1}{\sqrt{3}} & * \\ * & \frac{1}{\sqrt{3}} & * \end{bmatrix}.$$

Based on Biswajit Karmakar talk, [link here](#).

# TM<sub>1</sub> oscillation parameters predictions.

$$U_{PMNS} = \begin{bmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{i\delta} \\ s_{12}c_{23} - c_{12}s_{13}s_{23}e^{i\delta} & c_{12}c_{s3} - s_{12}s_{13}s_{23}e^{i\delta} & c_{13}s_{23} \\ s_{12}s_{23} - c_{12}s_{13}c_{23}e^{i\delta} & -c_{12}s_{23} - s_{12}s_{13}c_{23}e^{i\delta} & c_{13}c_{23} \end{bmatrix},$$

$$U_{TM_1} = \begin{bmatrix} \frac{2}{\sqrt{6}} & \frac{c_\theta}{\sqrt{3}} & \frac{s_\theta}{\sqrt{3}}e^{-i\gamma} \\ -\frac{c_\theta}{\sqrt{6}} & \frac{c_\theta}{\sqrt{3}} - \frac{s_\theta}{\sqrt{2}}e^{i\gamma} & -\frac{s_\theta}{\sqrt{3}}e^{-i\gamma} - \frac{c_\theta}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{c_\theta}{\sqrt{3}} - \frac{s}{\sqrt{2}}e^{i\gamma} & -\frac{s_\theta}{\sqrt{3}}e^{-i\gamma} + \frac{c_\theta}{\sqrt{2}} \end{bmatrix}.$$

Comparing the corresponding elements of the first column of  $U_{PMNS}$  and  $U_{TM_1}$ .

$$|U_{e1}|^2 = c_{12}^2 c_{13}^2 = 2/3 : s_{12}^2 = \frac{1 - 3s_{13}^2}{3 - 3s_{13}^2},$$

$$|U_{\mu 1}|^2 = |U_{\tau 1}|^2 = 1/6 : \cos \delta_{CP} = \frac{(1 - 5s_{13}^2)(2s_{23}^2 - 1)}{4s_{13}s_{23}\sqrt{2(1 - 3s_{13}^2)(1 - s_{23}^2)}}.$$

Based on [arXiv:1212.3247](https://arxiv.org/abs/1212.3247)

# TM<sub>2</sub> oscillation parameters predictions.

$$U_{PMNS} = \begin{bmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{i\delta} \\ s_{12}c_{23} - c_{12}s_{13}s_{23}e^{i\delta} & c_{12}c_{s_3} - s_{12}s_{13}s_{23}e^{i\delta} & c_{13}s_{23} \\ s_{12}s_{23} - c_{12}s_{13}c_{23}e^{i\delta} & -c_{12}s_{23} - s_{12}s_{13}c_{23}e^{i\delta} & c_{13}c_{23} \end{bmatrix},$$



$$U_{TM_2} = \begin{bmatrix} \frac{2c_\theta}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{2s_\theta}{\sqrt{6}}e^{-i\gamma} \\ -\frac{c_\theta}{\sqrt{6}} + \frac{s}{\sqrt{2}}e^{i\gamma} & \frac{1}{\sqrt{3}} & -\frac{s_\theta}{\sqrt{3}}e^{-i\gamma} - \frac{c_\theta}{\sqrt{2}} \\ -\frac{c_\theta}{\sqrt{6}} + \frac{s}{\sqrt{2}}e^{i\gamma} & \frac{1}{\sqrt{3}} & -\frac{s_\theta}{\sqrt{3}}e^{-i\gamma} + \frac{c_\theta}{\sqrt{2}} \end{bmatrix}.$$

Comparing the corresponding elements of the second column of  $U_{PMNS}$  and  $U_{TM_2}$ .

$$|U_{e2}|^2 = s_{12}^2 c_{13}^2 = 1/3 \quad : \quad s_{12}^2 = \frac{1}{3 - 3s_{13}^2},$$

$$|U_{\mu 2}|^2 = |U_{\tau 2}|^2 = 1/3 \quad : \quad \cos \delta_{CP} = -\frac{(2 - 4s_{13}^2)(2s_{23}^2 - 1)}{4s_{13}s_{23}\sqrt{(2 - 3s_{13}^2)(1 - s_{23}^2)}}.$$



# TM<sub>1</sub> and TM<sub>2</sub>, $\delta_{\text{CP}}$ vs. $\sin^2 \theta_{23}$ .



$$\text{TM}_1 : \cos \delta_{\text{CP}} = \frac{(1 - 5s_{13}^2)(2s_{23}^2 - 1)}{4s_{13}s_{23}\sqrt{2(1 - 3s_{13}^2)(1 - s_{23}^2)}},$$

$$\text{TM}_2 : \cos \delta_{\text{CP}} = -\frac{(2 - 4s_{13}^2)(2s_{23}^2 - 1)}{4s_{13}s_{23}\sqrt{(2 - 3s_{13}^2)(1 - s_{23}^2)}}.$$


# $\text{TM}_1$ and $\text{TM}_2$ , $\sin^2 \theta_{12}$ vs. $\sin^2 \theta_{13}$ .



$$\text{TM}_1 : s_{12}^2 = \frac{1 - 3s_{13}^2}{3 - 3s_{13}^2}, \quad \text{TM}_2 : s_{12}^2 = \frac{1}{3 - 3s_{13}^2}.$$



# NuFIT 5.2 (2022).

NuFIT 5.2 (2022)			
	Normal Ordering (best fit)		Inverted Ordering ( $\Delta\chi^2 = 2.3$ )
	bfp $\pm 1\sigma$	3 $\sigma$ range	bfp $\pm 1\sigma$
$\sin^2 \theta_{12}$	$0.303^{+0.012}_{-0.011}$	$0.270 \rightarrow 0.341$	$0.303^{+0.012}_{-0.011}$
$\theta_{12}/^\circ$	$33.41^{+0.75}_{-0.72}$	$31.31 \rightarrow 35.74$	$33.41^{+0.75}_{-0.72}$
$\sin^2 \theta_{23}$	$0.572^{+0.018}_{-0.023}$	$0.406 \rightarrow 0.620$	$0.578^{+0.016}_{-0.021}$
$\theta_{23}/^\circ$	$49.1^{+1.0}_{-1.3}$	$39.6 \rightarrow 51.9$	$49.5^{+0.9}_{-1.2}$
$\sin^2 \theta_{13}$	$0.02203^{+0.00056}_{-0.00059}$	$0.02029 \rightarrow 0.02391$	$0.02219^{+0.00060}_{-0.00057}$
$\theta_{13}/^\circ$	$8.54^{+0.11}_{-0.12}$	$8.19 \rightarrow 8.89$	$8.57^{+0.12}_{-0.11}$
$\delta_{CP}/^\circ$	$197^{+42}_{-25}$	$108 \rightarrow 404$	$286^{+27}_{-32}$
$\frac{\Delta m^2_{21}}{10^{-5} \text{ eV}^2}$	$7.41^{+0.21}_{-0.20}$	$6.82 \rightarrow 8.03$	$7.41^{+0.21}_{-0.20}$
$\frac{\Delta m^2_{3\ell}}{10^{-3} \text{ eV}^2}$	$+2.511^{+0.028}_{-0.027}$	$+2.428 \rightarrow +2.597$	$-2.498^{+0.032}_{-0.025}$
with SK atmospheric data			
	Normal Ordering (best fit)		Inverted Ordering ( $\Delta\chi^2 = 6.4$ )
	bfp $\pm 1\sigma$	3 $\sigma$ range	bfp $\pm 1\sigma$
$\sin^2 \theta_{12}$	$0.303^{+0.012}_{-0.012}$	$0.270 \rightarrow 0.341$	$0.303^{+0.012}_{-0.011}$
$\theta_{12}/^\circ$	$33.41^{+0.75}_{-0.72}$	$31.31 \rightarrow 35.74$	$33.41^{+0.75}_{-0.72}$
$\sin^2 \theta_{23}$	$0.451^{+0.019}_{-0.016}$	$0.408 \rightarrow 0.603$	$0.569^{+0.016}_{-0.021}$
$\theta_{23}/^\circ$	$42.2^{+1.1}_{-0.9}$	$39.7 \rightarrow 51.0$	$49.0^{+1.0}_{-1.2}$
$\sin^2 \theta_{13}$	$0.02225^{+0.00056}_{-0.00059}$	$0.02052 \rightarrow 0.02398$	$0.02223^{+0.00058}_{-0.00058}$
$\theta_{13}/^\circ$	$8.58^{+0.11}_{-0.11}$	$8.23 \rightarrow 8.91$	$8.57^{+0.11}_{-0.11}$
$\delta_{CP}/^\circ$	$232^{+36}_{-26}$	$144 \rightarrow 350$	$276^{+22}_{-29}$
$\frac{\Delta m^2_{21}}{10^{-5} \text{ eV}^2}$	$7.41^{+0.21}_{-0.20}$	$6.82 \rightarrow 8.03$	$7.41^{+0.21}_{-0.20}$
$\frac{\Delta m^2_{3\ell}}{10^{-3} \text{ eV}^2}$	$+2.507^{+0.026}_{-0.027}$	$+2.427 \rightarrow +2.590$	$-2.486^{+0.025}_{-0.028}$



Taken from <http://www.nu-fit.org>, updated arXiv:2007.14792, Table 3.

# $\text{TM}_1$ and $\text{TM}_2$ , $\delta_{\text{CP}}$ vs. $\sin^2 \theta_{23}$ , constrained, $3\sigma$ .





$$\text{TM}_1 : \quad \cos \delta_{\text{CP}} = \frac{(1-5s_{13}^2)(2s_{23}^2-1)}{4s_{13}s_{23}\sqrt{2(1-3s_{13}^2)(1-s_{23}^2)}},$$

$$\text{TM}_2 : \quad \cos \delta_{\text{CP}} = -\frac{(2-4s_{13}^2)(2s_{23}^2-1)}{4s_{13}s_{23}\sqrt{(2-3s_{13}^2)(1-s_{23}^2)}}.$$

Darker shaded regions = constrained with correlations.

# TM<sub>1</sub> and TM<sub>2</sub>, $\delta_{\text{CP}}$ vs. $\sin^2 \theta_{23}$ , constrained, $2\sigma$ .





$$\text{TM}_1 : \cos \delta_{\text{CP}} = \frac{(1-5s_{13}^2)(2s_{23}^2-1)}{4s_{13}s_{23}\sqrt{2(1-3s_{13}^2)(1-s_{23}^2)}},$$

$$\text{TM}_2 : \cos \delta_{\text{CP}} = -\frac{(2-4s_{13}^2)(2s_{23}^2-1)}{4s_{13}s_{23}\sqrt{(2-3s_{13}^2)(1-s_{23}^2)}}.$$

Darker shaded regions = constrained with correlations.

# TM<sub>1</sub> and TM<sub>2</sub>, $\delta_{\text{CP}}$ vs. $\sin^2 \theta_{13}$ , constrained, $3\sigma$ .





$$\text{TM}_1 : \quad \cos \delta_{\text{CP}} = \frac{(1 - 5s_{13}^2)(2s_{23}^2 - 1)}{4s_{13}s_{23}\sqrt{2(1 - 3s_{13}^2)(1 - s_{23}^2)}},$$

$$\text{TM}_2 : \quad \cos \delta_{\text{CP}} = -\frac{(2 - 4s_{13}^2)(2s_{23}^2 - 1)}{4s_{13}s_{23}\sqrt{(2 - 3s_{13}^2)(1 - s_{23}^2)}}.$$

Darker shaded regions = constrained with correlations.

# TM<sub>1</sub> and TM<sub>2</sub>, $\delta_{\text{CP}}$ vs. $\sin^2 \theta_{13}$ , constrained, $2\sigma$ .



$$\text{TM}_1 : \quad \cos \delta_{\text{CP}} = \frac{(1 - 5s_{13}^2)(2s_{23}^2 - 1)}{4s_{13}s_{23}\sqrt{2(1 - 3s_{13}^2)(1 - s_{23}^2)}},$$

$$\text{TM}_2 : \quad \cos \delta_{\text{CP}} = -\frac{(2 - 4s_{13}^2)(2s_{23}^2 - 1)}{4s_{13}s_{23}\sqrt{(2 - 3s_{13}^2)(1 - s_{23}^2)}}.$$

Darker shaded regions = constrained with correlations.

# Summary


Improvements in precise determination neutrino oscillation parameters triggered construction and tests of discrete symmetry neutrino flavor models.

- ① disfavored by oscillation parameters (history)
  - Bimaximal, Tribimaximal, Golden Ratio, Hexagonal Mixings disfavored by non-zero  $\theta_{13}$ ;
- ② successors of Tribimaximal Mixings (current work)
  - TM<sub>1</sub> and TM<sub>2</sub> predictions are significantly constrained by oscillation parameters correlations;
  - TM<sub>2</sub> is not applicable at  $2\sigma$  or less;
- ③ conclusion
  - taking into account correlations between oscillation parameters is a significant step forward in testing and constraining the predictions of discrete symmetry neutrino flavor models;

Thank you  
for your attention.

# Neutrino oscillation parameters data files.

NuFIT 5.2, [link here](#),



Valencia neutrino global fit, [link here](#).

