



UNIVERSITÀ
DEGLI STUDI
DI MILANO



The inclusion of QED corrections in the NNPDF4.0 fitting framework

Niccolò Laurenti,
on behalf of the NNPDF collaboration
Based on [2401.08749]


ICHEP24, Prague, 18/07/2024




Istituto Nazionale di Fisica Nucleare




Outline




PDFs fitting




How to add QED effects



Results




Impact on phenomenology




Summary and Outlook


PDFs fitting



PDFs fitting



How to add QED effects



Results





Impact on phenomenology



Summary and Outlook


- How do we compute observables in HEP?
- What are the PDFs?

Deep inelastic scattering (DIS)




$$\sigma = \sum_i \hat{\sigma}_i \otimes f_i + \mathcal{O}(\Lambda^2/Q^2)$$


- How are the PDFs fitted?
- We have to define a **theory**
- We have to choose a **dataset**
- We have to choose a fitting **methodology**




How to add QED effects



PDFs fitting



How to add QED effects



Results



Impact on phenomenology




Summary and Outlook

- Why do we want to add QED effects in PDFs?
- Are there cases in which they are not negligible?

$$\alpha \sim \mathcal{O}(\alpha_s^2) \sim \mathcal{O}(1\%)$$


percent correction

quarks can emit photons



we get a photon PDF

For some processes we can't neglect photon induced contributions



starts at $\mathcal{O}(\alpha_s^0)$

- How is the photon PDF determined?
- LuxQED gives a constraint between the photon PDF and the QCD PDFs

LuxQED approach

$$x\gamma(x, \mu^2) = \frac{1}{2\pi\alpha(\mu^2)} \int_x^1 \frac{dz}{z} \left\{ \int_{\frac{m_p^2 x^2}{1-z}}^{\frac{\mu^2}{1-z}} \frac{dQ^2}{Q^2} \alpha^2(Q^2) \left[-z^2 F_L(x/z, Q^2) + \left(zP_{\gamma q}(z) + \frac{2x^2 m_p^2}{Q^2} \right) F_2(x/z, Q^2) \right] - \alpha^2(\mu^2) z^2 F_2(x/z, \mu^2) \right\}$$

$$F_{2,L} = \sum_i C_{2,L,i} \otimes f_i$$

It modifies the sum rules

$$\int_0^1 dx x \left(\Sigma(x, Q^2) + g(x, Q^2) + \gamma(x, Q^2) \right) = 1$$

- How are DGLAP equations in presence of QED corrections?
- The photon PDF mixes with the other PDFs through evolution

$$\mu^2 \frac{d}{d\mu^2} f_i(x, \mu^2) = \sum_{j=q, \bar{q}, g, \gamma} \int_x^1 \frac{dz}{z} P_{ij} \left(\frac{x}{z}, \alpha_s(\mu^2), \alpha(\mu^2) \right) f_j(z, \mu^2)$$

$$i = q, \bar{q}, g, \gamma$$

$$P_{ij}(\alpha_s, \alpha) = P_{ij}^{\text{QCD}}(\alpha_s) + \tilde{P}_{ij}(\alpha_s, \alpha)$$

pure QCD terms


pure QED and QCD \otimes QED terms

$$P_{ij}^{\text{QCD}}(\alpha_s) = \alpha_s P_{ij}^{(0)} + \alpha_s^2 P_{ij}^{(1)} + \alpha_s^3 P_{ij}^{(2)} + \dots$$

$$\tilde{P}_{ij}(\alpha_s, \alpha) = \alpha P_{ij}^{(0,1)} + \alpha_s \alpha P_{ij}^{(1,1)} + \alpha^2 P_{ij}^{(0,2)} + \dots$$


The QED case is more difficult to solve than the pure QCD one (backup)

- What is the fitting methodology?
- LuxQED formula gives a constraint between γ and the other PDFs: **such constraint is implemented iteratively**




Computed at 100 GeV and evolved back to fitting scale with DGLAP [Manohar, Nason, Salam, Zanderighi, 2017]
(backup)


Results



PDFs fitting



How to add QED effects



Results





Impact on phenomenology





Summary and Outlook



- Results at fitting scale
- Very small differences in the quarks and gluon




- Results at 100 GeV
- Difference grows due to the effect of the photon in the evolution




- Photon PDF:
- Difference with NNPDF3.1QED is less than percent
- Percent difference with the other photon PDFs from the latest QED fits




Impact on phenomenology




PDFs fitting



How to add QED effects



Results




Impact on phenomenology




Summary and Outlook

- There are regions in which QED effects are not negligible
- Difference is at the level of few percent
- Photon in subtracting momentum from the other PDFs




- There are regions in which QED effects are not negligible
- Difference is at the level of few percent
- Photon in subtracting momentum from the other PDFs




Summary and Outlook



PDFs fitting




How to add QED effects



Results



Impact on phenomenology




Summary and Outlook

Summary and Outlook


- We can add QED corrections to PDF fitting, getting a photon PDF
- The photon PDF is compatible with the most recent QED fits
- Quarks and gluon are almost unchanged (the photon PDF is small)
- There are processes in which photon initiated contributions are not negligible

Thank you for your attention!


Backup slides



PDFs fitting



How to add QED effects



Results




Impact on phenomenology




Summary and Outlook

- **Theory**
- What defines the theory of a fit?




- **Dataset**
- Which data points are included in the fit?



4618 data points from
different processes

- **Methodology**
- How are the PDFs extracted?



Solving DGLAP

Pure QCD case

$$\mu^2 \frac{d}{d\mu^2} \begin{pmatrix} g \\ \Sigma \end{pmatrix} = \begin{pmatrix} P_{gg} & P_{gq} \\ P_{qg} & P_{qq} \end{pmatrix} \otimes \begin{pmatrix} g \\ \Sigma \end{pmatrix}$$

$$\mu^2 \frac{d}{d\mu^2} V = P_{ns,v} \otimes V$$

$$\mu^2 \frac{d}{d\mu^2} f_{ns,\pm} = P_{ns,\pm} \otimes f_{ns,\pm}$$

$$\Sigma = \sum_{i=1}^{n_f} q_i^+ \quad V = \sum_{i=1}^{n_f} q_i^- \quad f_{ns,\pm} = \begin{cases} u^\pm - d^\pm \\ u^\pm + d^\pm - 2s^\pm \\ u^\pm + d^\pm + s^\pm - 3c^\pm \\ u^\pm + d^\pm + s^\pm + c^\pm - 4b^\pm \\ u^\pm + d^\pm + s^\pm + c^\pm + b^\pm - 5t^\pm \end{cases}$$

$$q^\pm = q \pm \bar{q}$$

QCD \otimes QED case

$$\mu^2 \frac{d}{d\mu^2} \begin{pmatrix} g \\ \gamma \\ \Sigma \\ \Sigma_\Delta \end{pmatrix} = \mathbf{P}_s \otimes \begin{pmatrix} g \\ \gamma \\ \Sigma \\ \Sigma_\Delta \end{pmatrix}$$

$$\mu^2 \frac{d}{d\mu^2} \begin{pmatrix} V \\ V_\Delta \end{pmatrix} = \mathbf{P}_v \otimes \begin{pmatrix} V \\ V_\Delta \end{pmatrix}$$

$$\mu^2 \frac{d}{d\mu^2} f_{ns,\pm}^{ul/d} = \left(P_{ns,\pm} + \tilde{P}_{ns,\pm}^{ul/d} \right) \otimes f_{ns,\pm}^{ul/d}$$

$$f_{ns,\pm}^u = \begin{cases} u^\pm - c^\pm \\ u^\pm + c^\pm - 2t^\pm \end{cases} \quad f_{ns,\pm}^d = \begin{cases} d^\pm - s^\pm \\ d^\pm + s^\pm - 2b^\pm \end{cases}$$

$$\Sigma_\Delta = \frac{n_d}{n_u} \sum_{i=1}^{n_u} u_i^+ - \sum_{i=1}^{n_d} d_i^+ \quad V_\Delta = \frac{n_d}{n_u} \sum_{i=1}^{n_u} u_i^- - \sum_{i=1}^{n_d} d_i^-$$

Solving DGLAP

$$\mathbf{P}_s = \begin{pmatrix} P_{gg} + \tilde{P}_{gg} & \tilde{P}_{g\gamma} & P_{gq} + \langle \tilde{P}_{gq} \rangle & \nu_u \tilde{P}_{g\Delta q} \\ \tilde{P}_{\gamma g} & \tilde{P}_{\gamma\gamma} & \langle \tilde{P}_{\gamma q} \rangle & \nu_u \tilde{P}_{\gamma\Delta q} \\ 2n_f(P_{qg} + \langle \tilde{P}_{qg} \rangle) & 2n_f \langle \tilde{P}_{q\gamma} \rangle & P_{qq} + \langle \tilde{P}_q^{\text{ns},+} \rangle + \langle e_q^2 \rangle^2 \tilde{P}_{ps} & \nu_u \tilde{P}_{\Delta q}^{\text{ns},+} + \nu_u e_{\Delta q}^2 \langle e_q^2 \rangle \tilde{P}_{ps} \\ 2n_f \nu_d \tilde{P}_{\Delta qg} & 2n_f \nu_d \tilde{P}_{\Delta q\gamma} & \nu_d \tilde{P}_{\Delta q}^{\text{ns},+} + \nu_d e_{\Delta q}^2 \langle e_q^2 \rangle \tilde{P}_{ps} & P_{\text{ns},+} + \{ \tilde{P}_q^{\text{ns},+} \} + \nu_u \nu_d (e_{\Delta q}^2)^2 \tilde{P}_{ps} \end{pmatrix}$$

$$\mathbf{P}_v = \begin{pmatrix} P_{\text{ns},V} + \langle \tilde{P}_q^{\text{ns},-} \rangle & \nu_u \tilde{P}_{\Delta q}^{\text{ns},-} \\ \nu_d \tilde{P}_{\Delta q}^{\text{ns},-} & P_{\text{ns},-} + \{ \tilde{P}_q^{\text{ns},-} \} \end{pmatrix}$$

$$\nu_{uld} = \frac{n_{uld}}{n_f}, \quad \langle \tilde{P}_q^{\text{ns},\pm} \rangle = \nu_u \tilde{P}_u^{\text{ns},\pm} + \nu_d \tilde{P}_d^{\text{ns},\pm},$$

$$\{ \tilde{P}_q^{\text{ns},\pm} \} = \nu_d \tilde{P}_u^{\text{ns},\pm} + \nu_u \tilde{P}_d^{\text{ns},\pm}, \quad \tilde{P}_{\Delta q}^{\text{ns},\pm} = \tilde{P}_u^{\text{ns},\pm} - \tilde{P}_d^{\text{ns},\pm}$$

Solution of the non-diagonal sectors

$$\mathbf{E}_S(\mu^2 \leftarrow \mu_0^2) = \mathcal{P} \exp \left(- \int_{\log \mu_0^2}^{\log \mu^2} \gamma_S(\alpha_s(\mu'^2), \alpha(\mu'^2)) d \log \mu'^2 \right) \simeq \prod_{k=0}^{n-1} \mathbf{E}_S(\mu^{2(k+1)} \leftarrow \mu^{2(k)})$$

$$\mu^{2(n)} = \mu^2$$

$$\gamma(N) = - \int_0^1 dz z^{N-1} P(z)$$

$$\mathbf{E}_S(\mu^{2(k+1)} \leftarrow \mu^{2(k)}) = \exp \left(- \gamma_S(\alpha_s(\mu^{2(k+1/2)}), \alpha(\mu^{2(k+1/2)})) \Delta \log \mu^{2(k)} \right)$$

Solved in Mellin space

$$\log \mu^{2(k+1/2)} = \frac{\log \mu^{2(k+1)} + \log \mu^{2(k)}}{2}$$

$$\Delta \log \mu^{2(k)} = \log \mu^{2(k+1)} - \log \mu^{2(k)}$$

Computation of the photon

Why the LuxQED formula is used at 100 GeV?

LuxQED neglects higher twist corrections

$$\mathcal{O}\left(\frac{\Lambda}{\mu}\right)$$

For low μ , the integral is dominated by low Q^2 structure functions
non-perturbative!