

Higgs precision Higgs physics in electron–proton scattering at CERN

Thursday 18 July 2024 18:10 (17 minutes)

The Large Hadron-electron Collider and the Future Circular Collider in electron-hadron mode will make possible the study of DIS in the TeV regime providing electron-proton collisions with instantaneous luminosities of $10^{34} \text{ cm}^{-2} \text{ s}^{-1}$. With a charged current cross section around 200 (1000) fb at the LHeC (FCC-eh), Higgs bosons will be produced abundantly. We examine the opportunities for studying several of its couplings, particularly $H \rightarrow b\bar{b}$, $H \rightarrow c\bar{c}$, $H \rightarrow WW$, and Higgs to invisible. We also discuss the possibilities to measure anomalous Higgs couplings, and the implications of precise parton densities measured in DIS on Higgs physics. We finally address the complementarity in measuring Higgs couplings between the LHeC and the FCC-he and the respective hadronic colliders, the HL-LHC and the FCC-hh, and e^+e^- Higgs factories, but will also emphasise the gain in accuracy achievable by combining results between those colliders.

Alternate track

I read the instructions above

Yes

Presenter: KLEIN, Uta (University of Liverpool (GB))

Session Classification: Higgs Physics

Track Classification: 01. Higgs Physics