Probing the speed of sound in QGP with multi-particle [pT] cumulants in ALICE

Thursday 18 July 2024 15:02 (17 minutes)

The speed of sound squared, c_s^2 , a property of the quark-gluon plasma (QGP) connected to the QCD equation of state, can be extracted from ultra-central heavy-ion collisions, where the medium maintains a fixed size and the initial-state and thermal fluctuations dominate. We present the first ALICE measurements of the event-by-event mean transverse momentum, $\langle [p_{\rm T}] \rangle$, its average and higher-order fluctuations as a function of multiplicity using particle spectra and multi-particle $\langle [p_{\rm T}] \rangle$ cumulant techniques, in ultra-central Pb-Pb collisions at $\sqrt{s_{\rm NN}}=5.02$ TeV. The pronounced rise in $\langle [p_{\rm T}] \rangle$ and the sudden transition in higher-order fluctuations at high multiplicities are used to extract the c_s^2 and to probe the thermalization of the QGP. Our approach yields valuable insights into the thermalized nature of the QGP, contributing to a deeper understanding of the QCD equation of state.

Alternate track

I read the instructions above

Yes

Primary author: VAZQUEZ RUEDA, Omar (University of Houston (US))

Co-author: COLLABORATION, ALICE

Presenter: VAZQUEZ RUEDA, Omar (University of Houston (US))

Session Classification: Heavy Ions

Track Classification: 07. Heavy Ions