

# Searches for heavy resonances decaying into Z, W and Higgs bosons at CMS

*Saturday 20 July 2024 11:02 (17 minutes)*

A summary of searches for heavy resonances with masses exceeding 1 TeV decaying into pairs or triplets of bosons is presented, performed on data produced by LHC pp collisions at  $\sqrt{s}=13\text{TeV}$  and collected with the CMS detector during 2016 - 2018. The common feature of these analyses is the boosted topology, namely the decay products of the considered bosons (both electroweak W, Z bosons and the Higgs boson) are expected to be highly energetic and close in angle, leading to a non-trivial identification of the quarks and leptons in the final state. The exploitation of jet substructure techniques allows to increase the sensitivity of the searches where at least one boson decays hadronically. Various background estimation techniques are adopted, based on data-MC hybrid approaches or relying only in control regions in data. Results are interpreted in the context of the Warped Extra Dimension and Heavy Vector Triplet theoretical models, two possible scenarios beyond the standard model.

## Alternate track

### I read the instructions above

Yes

**Primary authors:** CMS; HUNDHAUSEN, Daniel (Hamburg University (DE))

**Presenter:** HUNDHAUSEN, Daniel (Hamburg University (DE))

**Session Classification:** Beyond the Standard Model

**Track Classification:** 03. Beyond the Standard Model