

Prospects for constraining light-quark electroweak couplings at Higgs factories


Krzysztof Mękała

DESY, Hamburg, Germany

Faculty of Physics, University of Warsaw, Poland

based on work in collaboration with D. Jeans, J. Reuter, J. Tian, A.F. Żarnecki

ICHEP 2024, Prague, July 2024



Quarks



Leptons






Higgs boson



Forces

A nice picture but it is not "self-explanatory":
it contains many free parameters.

Measuring precision observables allows
to constrain the SM parameters
and to search for New Physics.



Z decays to hadrons are constrained from LEP and SLC...

$$R_b = \Gamma(b\bar{b})/\Gamma(\text{hadrons})$$

$$\Gamma_{12}/\Gamma_8$$

OUR FIT is obtained by a simultaneous fit to several c - and b -quark measurements as explained in the note "The Z boson" and ref. LEP-SLC 06.

The Standard Model predicts $R_b=0.21581$ for $m_t=174.3$ GeV and $M_H=150$ GeV.

VALUE	DOCUMENT ID	TECN	COMMENT
0.21629 ± 0.00066 OUR FIT			
0.21594 ± 0.00094 ± 0.00075	1 ABE	05F SLD	$E_{\text{cm}}^{ee} = 91.28$ GeV
0.2174 ± 0.0015 ± 0.0028	2 ACCIARRI	00 L3	$E_{\text{cm}}^{ee} = 89-93$ GeV
0.2178 ± 0.0011 ± 0.0013	3 ABBIENDI	99B OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV
0.21634 ± 0.00067 ± 0.00060	4 ABREU	99B DLPH	$E_{\text{cm}}^{ee} = 88-94$ GeV
0.2159 ± 0.0009 ± 0.0011	5 BARATE	97F ALEP	$E_{\text{cm}}^{ee} = 88-94$ GeV

*Review of Particle Physics,
PDG, 2022*

$$R_c = \Gamma(c\bar{c})/\Gamma(\text{hadrons})$$

$$\Gamma_{11}/\Gamma_8$$

OUR FIT is obtained by a simultaneous fit to several c - and b -quark measurements as explained in the note "The Z boson" and ref. LEP-SLC 06.

The Standard Model predicts $R_c = 0.1723$ for $m_t = 174.3$ GeV and $M_H = 150$ GeV.

VALUE	DOCUMENT ID	TECN	COMMENT
0.1721 ± 0.0030 OUR FIT			
0.1744 ± 0.0031 ± 0.0021	1 ABE	05F SLD	$E_{\text{cm}}^{ee} = 91.28$ GeV
0.1665 ± 0.0051 ± 0.0081	2 ABREU	00 DLPH	$E_{\text{cm}}^{ee} = 88-94$ GeV
0.1698 ± 0.0069	3 BARATE	00B ALEP	$E_{\text{cm}}^{ee} = 88-94$ GeV
0.180 ± 0.011 ± 0.013	4 ACKERSTAFF	98E OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV
0.167 ± 0.011 ± 0.012	5 ALEXANDER	96R OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV

$$\Gamma((u\bar{u} + c\bar{c})/2)/\Gamma(\text{hadrons})$$

$$\Gamma_9/\Gamma_8$$

This quantity is the branching ratio of $Z \rightarrow$ "up-type" quarks to $Z \rightarrow$ hadrons. Except ACKERSTAFF 97T the values of $Z \rightarrow$ "up-type" and $Z \rightarrow$ "down-type" branchings are extracted from measurements of $\Gamma(\text{hadrons})$, and $\Gamma(Z \rightarrow \gamma + \text{jets})$ where γ is a high-energy (>5 or 7 GeV) isolated photon. As the experiments use different procedures and slightly different values of M_Z , $\Gamma(\text{hadrons})$ and α_S in their extraction procedures, our average has to be taken with caution.

VALUE	DOCUMENT ID	TECN	COMMENT
0.166 ± 0.009 OUR AVERAGE			
0.172 ^{+0.011} _{-0.010}	1 ABBIENDI	04E OPAL	$E_{\text{cm}}^{ee} = 91.2$ GeV
0.160 ± 0.019 ± 0.019	2 ACKERSTAFF	97T OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV
0.137 ^{+0.038} _{-0.054}	3 ABREU	95X DLPH	$E_{\text{cm}}^{ee} = 88-94$ GeV
0.137 ± 0.033	4 ADRIANI	93 L3	$E_{\text{cm}}^{ee} = 91.2$ GeV


Future e^+e^- colliders operating at the Z-pole would be a perfect place to study the couplings.



ILC



CEPC



FCC-ee

Source	$e^-e^+ \rightarrow c\bar{c}$				$e^-e^+ \rightarrow b\bar{b}$			
	$P_{e^-e^+}(-0.8, +0.3)$ R_c	$A_{FB}^{c\bar{c}}$	$P_{e^-e^+}(+0.8, -0.3)$ R_c	$A_{FB}^{c\bar{c}}$	$P_{e^-e^+}(-0.8, +0.3)$ R_b	$A_{FB}^{b\bar{b}}$	$P_{e^-e^+}(+0.8, -0.3)$ R_b	$A_{FB}^{b\bar{b}}$
Statistics	0.18%	0.38%	0.27%	0.52%	0.12%	0.24%	0.23%	0.70%
Preselection eff.	<0.01%	0.12%	0.02%	0.16%	<0.01%	0.08%	0.06%	0.12%
Background	0.01%	0.01%	0.02%	0.02%	0.01%	0.01%	0.06%	<0.01%
heavy quark mistag	0.11%	<0.01%	0.06%	<0.01%	0.12%	<0.01%	0.22%	<0.01%
<i>uds</i> mistag	0.03%	<0.01%	0.02%	<0.01%	0.08%	<0.01%	0.14%	<0.01%
Angular correlations	0.10%	0.10%	0.10%	0.10%	0.10%	0.10%	0.10%	0.10%
Beam Polarisation	<0.01%	<0.01%	0.02%	0.01%	<0.01%	0.01%	0.03%	0.15%
Systematics	0.15%	0.16%	0.12%	0.19%	0.18%	0.13%	0.29%	0.22%
Total	0.24%	0.41%	0.30%	0.55%	0.21%	0.27%	0.37%	0.73%

Electroweak observables at ILC250:
A. Irles *et al.*, [2306.11413]

The cross sections to heavy quarks could be well constrained at ILC thanks to excellent flavour-tagging.

But how to take the measurement if...

- tagging is imperfect (s quark)?
- tagging is unavailable (u, d quarks)?

How to measure Z couplings
to light quarks?

General idea

We want to measure quark couplings:

$$c_f = v_f^2 + a_f^2$$

They are given in the SM by:

$$v_f = 2I_{3,f} - 4Q_f \sin^2 \theta_W \quad a_f = 2I_{3,f}$$

Γ_{had} reads (exact at fixed order):

$$\Gamma_{had} = N_c \frac{G_\mu M_Z^3}{24\pi\sqrt{2}} \left(1 + \frac{\alpha_s}{\pi} + \dots \right) (3c_d + 2c_u)$$

and $\Gamma_{had+\gamma}$:

$$\Gamma_{had+\gamma} = N_c \frac{G_\mu M_Z^3}{24\pi\sqrt{2}} f(y_{cut}) \frac{\alpha}{2\pi} (3q_d^2 c_d + 2q_u^2 c_u)$$

The correction factor $f(y_{cut})$ to be determined for a given value of the resolution parameter y_{cut} .

Measurement at the Z-pole

We can measure radiative and non-radiative cross sections separately and disentangle the couplings c_d and c_u :

$$\sigma_{Z \rightarrow had} = \mathcal{C}_1 \cdot (3c_d + 2c_u)$$

$$\sigma_{Z \rightarrow had+\gamma} = \mathcal{C}_2 \cdot (3c_d + 8c_u)$$

Note: in this picture, the couplings are universal among quarks of the same type. This assumption can be lifted by employing heavy-flavour tagging.

How to generate Monte Carlo
events?

Analysis setup

We want to consider:

$$e^+ e^- \rightarrow q\bar{q}(\gamma)$$

Experimentally measured photons can originate not only from the Final State Radiation but also from the Initial State Radiation, hadronisation and decays...

One may encounter the following issues:


- **Matrix Element** calculations – divergent or very slow for low photon-emission angles;
- **ISR structure functions** – good for small angles, a proper matching procedure needed;
- **FSR showers** – important for QCD emissions, may cause double-counting;
- **hadron decays** – photons to be included properly.

Matching procedure – *Whizard* perspective

- matching: *soft* physics invisible in the detector, *hard* physics properly reconstructed
- soft ISR photons simulated using built-in structure functions
- soft FSR photons simulated using parton showers
- hard photons simulated using full ME calculations (0, 1, 2... ME γ samples)
 - momentum transfer and energy to define the soft and hard regimes

Efficiency of the matching procedure

- ILC@Z-pole: about 4% of *Whizard* events are rejected to avoid double-counting.



How to select events?

work in progress

Event reconstruction



Measurable photons can originate from:

- Initial State Radiation,
- Final State Radiation,
- hadronisation and decays.

The interesting information comes only from FSR so reconstruction criteria should reduce other contributions.

Z-pole: ISR naturally reduced (but often not in the *baseline* scenario)


Isolation criterion y_{cut}



$$y_{\text{cut}} = E_\gamma \cdot \sin(\theta_{\gamma q_i}^{\text{min}})$$

preliminary, ILC@Z-pole

Isolation criterion y_{cut}



$$\frac{q_u}{q_d} = \left(\frac{Q_u}{Q_d} \right)^2 \cdot \frac{N_u}{N_d} \cdot \frac{c_u}{c_d} \approx 2.1$$

preliminary, ILC@Z-pole

Conclusions

- Couplings of the Z boson to light quarks are poorly constrained but an improvement could be achieved at future lepton colliders.
- Proper assessment of the uncertainties requires deep understanding of theoretical calculations, event simulation and reconstruction.
- *Work in progress...*

Backup

Backup: matching criteria

$$q_- = \sqrt{4E_0 E_\gamma} \sin \frac{\theta_\gamma}{2}$$
$$q_+ = \sqrt{4E_0 E_\gamma} \cos \frac{\theta_\gamma}{2}$$

Simulating events with *Whizard* and *Pythia6* (shower and hadronisation)

- ME cuts:

- all γ 's:

$$q_\pm > 1 \text{ GeV} \text{ and } E > 1 \text{ GeV} \text{ and } M(\gamma, q_i) > 1 \text{ GeV}$$

- event selection:

- all ISR SF γ 's:

$$q_\pm < 1 \text{ GeV} \text{ or } E < 1 \text{ GeV} \text{ or } M(\gamma, q_i) < 1 \text{ GeV}$$

- all FSR shower γ 's whose parents are initial quarks:

$$q_\pm < 1 \text{ GeV} \text{ or } E < 1 \text{ GeV} \text{ or } M(\gamma, q_i) < 1 \text{ GeV}$$