

Triggering on electrons, photons, tau leptons, Jets and energy sums at HL-LHC with the upgraded CMS Level-1 Trigger

Thursday 18 July 2024 20:40 (20 minutes)

The High-Luminosity LHC will open an unprecedented window on the weak-scale nature of the universe, providing high-precision measurements of the Standard Model as well as searches for new physics beyond the standard model. The Compact Muon Solenoid (CMS) experiment is planning to replace entirely its trigger and data acquisition system to achieve this ambitious physics program. Efficiently collecting those datasets will be a challenging task, given the harsh environment of 200 proton-proton interactions per LHC bunch crossing. The new Level-1 trigger architecture for HL-LHC will improve performance with respect to Phase I through the addition of tracking information and subdetector upgrades leading to higher granularity and precision timing information. In this poster, we present a large panel of trigger algorithms for the upgraded Phase II trigger system, which benefit from the finer information to reconstruct optimally the physics objects. The expected performance will be presented.

Alternate track

I read the instructions above

Yes

Authors: CMS; PAINESIS, Haris (National and Kapodistrian University of Athens (GR))

Presenter: PAINESIS, Haris (National and Kapodistrian University of Athens (GR))

Session Classification: Poster Session 1

Track Classification: 14. Computing, AI and Data Handling