Likelihood and Deep Learning Analysis of the electron neutrino event sample at Intermediate Water Cherenkov Detector (IWCD) of the Hyper-Kamiokande experiment

Friday, 19 July 2024 20:40 (20 minutes)

Hyper-K is a next-generation long baseline neutrino experiment. One of its primary physics goals is to measure neutrino oscillation parameters precisely, including CP-asymmetry. As conventional vµ beam from J-PARC neutrino baseline contains only 1.5% of ve interaction of total, it is challenging to measure ve/ve(anti) scattering cross-section on nuclei. To reduce systematic uncertainty, IWCD will be built to study neutrino interaction rate with higher accuracy. The presented, simulated data comprises veCC0 π as main signal & NC π 0, vµCC are major background events. To reduce the backgrounds, initially a log-likelihood-based reconstruction algorithm to select candidate events was used, which however, sometimes struggles to distinguish π 0 events properly from electron-like events. Thus, ML-based framework has been developed to enhance the purity & signal efficiency rate of ve events. Implementing it notably enhances both the efficiency & purity of ve signals over the conventional approach.

Alternate track

I read the instructions above

Yes

Primary author: Ms MONDAL, Tanima (IIT Kharagpur, India)Presenter: Ms MONDAL, Tanima (IIT Kharagpur, India)Session Classification: Poster Session 2

Track Classification: 02. Neutrino Physics