Barium tagging in NEXT: toward an ultra-low background 0vββ decay detector.

Thursday 18 July 2024 20:40 (20 minutes)

Neutrinoless double beta decay experiments are pushing their sensitivities to reach half-lives on the order of 10^{28} years. A promising approach involves detecting the daughter ion generated in the decay. The NEXT collaboration is testing chemical sensors to identify the Ba²⁺ ion produced in the double beta decay of ¹³⁶Xe, coinciding with the emission of two electrons. This entails a challenge, since only a few ions per year would be produced in the NEXT chamber. Further the chemosensors must be compatible with the ultra-dry conditions of xenon. The R&D effort in NEXT towards Barium tagging is two fold: First, to build a Single-Molecule Fluorescence Imaging (SMFI) system with enough sensitivity to detect the signal from the chemosensors and capable of operating under a dry environment. Second, to develop a barium source reproducing the conditions of the NEXT detector. This two systems will be integrated to compose a reliable Barium-tagging sensor.

Alternate track

1. Detectors for Future Facilities, R&D, Novel Techniques

I read the instructions above

Yes

Author: HERRERO GOMEZ, Pablo (Hebrew University)

Co-authors: Dr SHOMRONI, Itay (Hebrew University); Dr ARAZI, Lior (Ben Gurion University of the Negev); Mr KAPON, Nir (Hebrew University)

Presenter: HERRERO GOMEZ, Pablo (Hebrew University)

Session Classification: Poster Session 1

Track Classification: 02. Neutrino Physics