An Improved Search for Unstable Sterile Neutrinos at IceCube

Philip Weigel
for the IceCube Collaboration

ICHEP 2024
July 20th, 2024

Massachusetts
Institute of
Technology

IceCube
South Pole Neutrino Observatory

Neutrino Oscillation Anomalies

- Several unresolved anomalies in various neutrino experiments







3+1 Sterile Neutrino Model

- Anomalies in short baseline neutrino experiments could be explained by a sterile neutrino
  - Introduce a new flavor and mass state, append a row and column to the PMNS matrix
  - These states do not interact weakly → only accessible through oscillations



$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \\ \nu_s \end{pmatrix} = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} & U_{e4} \\ U_{\mu 1} & U_{\mu 2} & U_{\mu 3} & U_{\mu 4} \\ U_{\tau 1} & U_{\tau 2} & U_{\tau 3} & U_{\tau 4} \\ U_{s1} & U_{s2} & U_{s3} & U_{s4} \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \\ \nu_4 \end{pmatrix}$$

$$|U_{e4}|^2 = \sin^2(\theta_{14})$$
$$|U_{\mu 4}|^2 = \sin^2(\theta_{24})\cos^2(\theta_{14})$$
$$|U_{\tau 4}|^2 = \sin^2(\theta_{34})\cos^2(\theta_{24})\cos^2(\theta_{34})$$

Atmospheric Neutrinos and IceCube

The conventional atmospheric muon neutrino flux originates from π and K decay-in-flight





Sterile Neutrino Oscillations in Matter

- To probe neutrino oscillations, IceCube can measure the energy and the zenith angle of the incoming neutrino
  - The zenith angle is a proxy for the baseline L, the distance that the neutrinos have traveled
- There are two main features in these oscillograms:



Sterile Neutrino Oscillations in Matter

- To probe neutrino oscillations, IceCube can measure the energy and the zenith angle of the incoming neutrino
  - The zenith angle is a proxy for the baseline L, the distance that the neutrinos have traveled
- There are two main features in these oscillograms:
  - Vacuum (L/E) oscillations



Sterile Neutrino Oscillations in Matter

- To probe neutrino oscillations, IceCube can measure the energy and the zenith angle of the incoming neutrino
  - The zenith angle is a proxy for the baseline L, the distance that the neutrinos have traveled
- There are two main features in these oscillograms:
  - Vacuum (L/E) oscillations
  - A matter-enhanced resonant disappearance for antineutrinos that travel through most of the Earth

$$E_{res} = \mp \frac{\Delta m^2 \cos(2\theta)}{\sqrt{2} G_F N_n}$$



Sterile Neutrino Oscillations in Matter

- To probe neutrino oscillations, IceCube can measure the energy and the zenith angle of the incoming neutrino
  - The zenith angle is a proxy for the baseline L, the distance that the neutrinos have traveled
- There are two main features in these oscillograms:
  - Vacuum (L/E) oscillations
  - A matter-enhanced resonant disappearance for antineutrinos that travel through most of the Earth

$$E_{res} = \mp \frac{\Delta m^2 \cos(2\theta)}{\sqrt{2} G_F N_n}$$



Improvements to Event Selection and Analysis

- To search for sterile neutrinos, we leverage the large flux of atmospheric muon (anti)neutrinos observed in IceCube
- Major improvements since the previous search:
  - BDT for removing atmospheric muon backgrounds
  - DNN-based energy reconstruction and classifier
  - More detailed ice systematic treatment
  - Improved atmospheric flux systematics
  - Broken power law for astrophysical flux







arXiv:2405.08077 [hep-ex]

Improvements to Event Selection and Analysis

- To search for sterile neutrinos, we leverage the large flux of atmospheric muon (anti)neutrinos observed in IceCube
- Major improvements since the previous search:
  - BDT for removing atmospheric muon backgrounds
  - DNN-based energy reconstruction and classifier
  - More detailed ice systematic treatment
  - Improved atmospheric flux systematics
  - Broken power law for astrophysical flux







arXiv:2405.08077 [hep-ex]

Latest 3+1 Sterile Neutrino Results

- Best Fit:

$$\Delta m^2_{41} = 3.5 \text{ eV}^2$$

$$\sin^2(2\theta_{24}) = 0.16$$

$$\text{p-value} = 3.1\% \longrightarrow 2.2\sigma$$

- Consistent with the previous sterile analyses in IceCube
- These results appeared on the arXiv in May, see: arXiv:2405.08070



Invisible Sterile Neutrino Decay



Adding a decay mechanism[1] dampens the sterile oscillation signature, which is preferred in the global fits[2]

$$\tau = \frac{1}{\Gamma} = \frac{16\pi}{g^2 m_4}$$

$\Delta m^2_{41} = 1.0\ \mathrm{eV}^2$

$\sin^2(2\theta_{24}) = 0.1$



E = 1 TeV

M. Moulai Thesis (MIT)

[1] *Phys. Rev. D 97, 055017*
[2] *J. High Energ. Phys.* **2023**, 58 (2023)

Results for 3+1+decay

- The best fit for the 3+1+decay search is the no-decay scenario
  - Best fit at $g^2 = 0$
  - p-value = 7.4% (3 dof)



Results for 3+1+decay

- The best fit for the 3+1+decay search is the no-decay scenario
  - Best fit at $g^2 = 0$
  - p-value = 7.4% (3 dof)



Results for 3+1+decay

- The best fit for the 3+1+decay search is the no-decay scenario
  - Best fit at $g^2 = 0$
  - p-value = 7.4% (3 dof)
- This result is consistent with the previous search from 2022
  - Previous best fit is excluded, but the LLH space was quite shallow so there is overlap



Results for 3+1+decay

- The best fit for the 3+1+decay search is the no-decay scenario
  - Best fit at $g^2 = 0$
  - p-value = 7.4% (3 dof)
- This result is consistent with the previous search from 2022
  - Previous best fit is excluded, but the LLH space was quite shallow so there is overlap
- Regions from the global fits to SBL experiments are excluded
  - Partially excluded at 99% CL
  - Mostly excluded at 95% CL



Systematic pulls



For details of the systematic treatment, see: [arXiv:2405:08077]

Systematic pulls



For details of the systematic treatment, see: arXiv:2405:08077

Systematic pulls



For details of the systematic treatment, see: arXiv:2405:08077

Conclusion

- IceCube has performed an improved search for unstable eV-scale sterile neutrinos using 10.7 years of muon neutrino data
  - Probed the invisible decay case, which is preferred over the ordinary 3+1 in the global fits
- No preference for decay was found, best fit point at $g^2 = 0$
  - Results are consistent with the previous analysis, but the previous best fit point is now excluded at >99% CL
- **This result puts constraints on the preferred region of the SBL experiments**

Thank you for listening!