

# Hidden Zeros and the Double Copy

*Saturday 20 July 2024 09:45 (15 minutes)*

Recently, Arkani-Hamed et al. proposed the existence of zeros in scattering amplitudes in certain quantum field theories including the cubic adjoint scalar theory  $\text{Tr}(\phi^3)$ , the  $SU(N)$  non-linear sigma model (NLSM) and Yang-Mills (YM) theory. These hidden zeros are special kinematic points where the amplitude vanishes and factorizes into a product of lower-point amplitudes, similar to factorization near poles. In our work, we show a close connection between the existence of such zeros and color-kinematics duality. In fact, the hidden zeros can be derived from the Bern-Carrasco-Johansson (BCJ) relations. We also show that these zeros extend via the Kawai-Lewellen-Tye (KLT) relations to special Galileon amplitudes and their corrections, evincing that these hidden zeros are also present in permutation-invariant amplitudes.

## Alternate track

### I read the instructions above

Yes

**Author:** OKTEM, Umut

**Co-authors:** BARTSCH, Christoph (Charles University Prague); TRNKA, Jaroslav; KAMPF, Karol (Charles University (CZ)); PARANJAPE, Shruti; BROWN, Taro

**Presenter:** OKTEM, Umut

**Session Classification:** Formal Theory

**Track Classification:** 10. Formal Theory