

XVth Quark Confinement and the Hadron Spectrum



Contribution ID: 135

Type: **Oral**

Anisotropic pressure and novel first-order phase transition in SU(3) Yang-Mills theory on $\mathbb{T}^2 \times \mathbb{R}^2$

Tuesday 20 August 2024 16:00 (20 minutes)

We consider a pure Yang-Mills theory on $\mathbb{T}^2 \times \mathbb{R}^2$ with boundary conditions imposed not only in the imaginary time direction but also in one spatial direction, and discuss thermodynamic quantities and their phase structures. The introduction of the boundary condition leads to the breaking of rotational symmetry, resulting in anisotropy of the pressure. Results from lattice QCD simulations [1] show that the anisotropic effect is suppressed until the spatial extension becomes significantly smaller near the critical temperature. This result is a very different behavior from that of massless free boson systems. In order to clarify the mechanism behind this result, we employ an effective model with two Polyakov loops along the time and spatial directions [2]. We show that introducing the interplay of two Polyakov loops well describes the lattice data in the high-temperature region, as suggested in a previous study [3]. Furthermore, we suggest the presence of a new first-order phase transition, which is distinguished from the confinement phase transition.

[1]Masakiyo Kitazawa et al, Phys. Rev. D 99, 094507 (2019)

[2]Daisuke Fujii, Akihiro Iwanaka, Masakiyo Kitazawa and Daiki Suenaga, arXiv:2402.17638 [hep-ph]

[3]Daiki Suenaga and Masakiyo Kitazawa, Phys. Rev. D 107, 074502 (2023)

Primary authors: FUJII, Daisuke (Japan Atomic Energy Agency); Mr IWANAKA, Akihiro (Research Center for Nuclear Physics); SUENAGA, Daiki (KMI, Nagoya University); KITAZAWA, Masakiyo

Presenter: FUJII, Daisuke (Japan Atomic Energy Agency)

Session Classification: Strongly-Coupled Theories and Dark Matter

Track Classification: G: Strongly-Coupled Theories and Dark Matter