

Plasma Wakefield Acceleration

AWAKE Experiment at CERN

Marlene Turner

Outline


- What are **plasma wakefields** and why are they interesting?
- How to accelerate **charged particles** using plasma wakefields?
 - Underlying physics concepts, state-of-the-art results
- What is the **AWAKE experiment**, and why is it important?
- The AWAKE experimental setup
- Latest AWAKE results
 - Ideas and plans for the future

Advanced Proton Driven Plasma Wakefield Acceleration Experiment



- Plasma ?
- Proton driven ?
- Wakefield acceleration ?
- Acceleration ?

Charged Particle Acceleration



- Acceleration of charged particles requires an electric field
- Charged particle will accelerate as long as it experiences the field

$1 \text{ TV} = 10^{12} \text{ V}$ 1.5V with battery length of $\sim 3\text{cm} \rightarrow 50 \text{ V/m}$


$1 \text{ GV} = 10^9 \text{ V}$

$1 \text{ MV} = 10^6 \text{ V}$ To reach $1 \text{ TeV} \rightarrow \sim 20 \text{ 000 million km}$


$1 \text{ kV} = 10^3 \text{ V}$ Distance Earth-Sun $\sim 152 \text{ million km}$



Charged Particle Acceleration




- Acceleration of charged particles requires an electric field
- Charged particle will accelerate as long as it experiences the field



- Even better:
 - Field travels together with the beam

Definition of Plasma and Plasma Wakefield

Plasma




Plasma: ionised gas (4th state of matter)

- **Quasi-neutrality:** the overall charge of a plasma is about zero.
- **Collective effects:** Charged particles must be close enough together that each particle influences many nearby charged particles.
- **Electrostatic interactions dominate** over collisions or ordinary gas kinetics.

Definition of Plasma and Plasma Wakefield


Plasma



Plasma: ionised gas (4th state of matter)


- **Quasi-neutrality:** the overall charge of a plasma is about zero.
- **Collective effects:** Charged particles must be close enough together that each particle influences many nearby charged particles.
- **Electrostatic interactions dominate** over collisions or ordinary gas kinetics.

Plasma Wakefields



Plasma Wakefields:

- are the **fields** created/sustained by collective motion of plasma particles.




Why use Plasmas for Charged Particle Acceleration?

Conventional technology:
metallic radiofrequency (RF) cavities



LHC cavities


New concept:
plasma wakefields acceleration
→ transient structures in plasma



Accelerating Gradient

RF cavities


Limited to $\sim 100 \text{ MV/m}$ due to electric breakdowns (ionization).



Plasma Wakefields

Plasma is already ionized or “broken-down” and can sustain electric fields $\sim 100 \text{ GV/m}$.


$$eE_{max} = 1 \left[\frac{\text{eV}}{\text{cm}} \right] \cdot n^{1/2} [\text{cm}^{-3}]$$



Accelerating Gradient

RF cavities

Limited to $\sim 100 \text{ MV/m}$ due to electric breakdowns (ionization).




→ Plasma wakefields can sustain order of magnitude higher fields

Plasma Wakefields

Plasma is already ionized or “broken-down” and can sustain electric fields $\sim 100 \text{ GV/m}$.

$$eE_{max} = 1 \left[\frac{\text{eV}}{\text{cm}} \right] \cdot n^{1/2} [\text{cm}^{-3}]$$




Structure exists only for a very short amount of time!

Circular and Linear Accelerators

Circular accelerators

Image from P. Arpaia et al.,
NIMA 985, 164652 (2021)




- Beam passes acceleration section multiple times.
- Max. energy (E) limited by synchrotron radiation losses
 $\propto E^4/(r^2m^4)$

- Advantage: beam passes accelerating section many times
- Disadvantage: synchrotron radiation losses

LHC tunnel:
 $p+p \rightarrow 14 \text{ TeV}$
 $e+e^- \rightarrow 209 \text{ GeV}$

Synchrotron Radiation



Synchrotron radiation is caused by leaving part of fields behind when the beam moves along the curve.

$$\propto E^4/(r^2 m^4)$$


Particle energy Bending radius Mass of the particle

→ Needs to be taken into account when accelerated charged particles are deflected in the radial direction.

Circular and Linear Accelerators


Circular accelerators

Image from P. Arpaia et al.,
NIMA 985, 164652 (2021)



- Beam passes acceleration section multiple times.
- Max. energy (E) limited by synchrotron radiation losses
 $\propto E^4/(r^2m^4)$

Linear accelerators



- Beam passes acceleration section multiple times.
- Negligible synchrotron radiation losses
- Accelerator length and accelerating gradient define final beam energy.

e.g. to accelerate electrons to 1 TeV (10^{12} eV):
100 MeV/m \times 10000 m or
100 GeV/m \times 10 m

Let Us Repeat...

- Plasma wakefields allow to accelerate charged particles with \sim 1-100 GeV/m
 - 10-1000 m to reach 1 TeV beam energy
- High gradients are important when using linear accelerators (e.g. for light particles) to minimize synchrotron radiation losses
 - For linear accelerators, their length defines the final beam energy

How to Create Plasma Wakefields?








Image from
<https://revbalance.com/improving-balance-for-wakesurfing/>



CEAN O'HAREZON 2013


Energy Source: The Driver




- Relativistic charged particle bunches or laser pulses
 - → Relativistic charged particle bunches carry almost purely **transverse electric** fields
- What we need → longitudinal electric field to accelerate charged particles

Trick:

- Use plasma to convert the transverse electric field of the proton bunch into a longitudinal electric field in the plasma.
- The more energy is available, the longer (distance-wise) these plasma wakefields can be sustained




How to Drive a Plasma Wave




Important to understand

- Plasma electron motion is mostly **transverse**
- Electrons do not move significantly longitudinally
- Rb ions are heavy and do not move significantly on the timescale of the electrons

How to Drive a Plasma Wave




Charge separation → electric field
(longitudinal and transverse)



Where should we place an electron bunch to be accelerated?



Plasma Wakefields



- ← Accelerating for negatively charged particles
- Decelerating for negatively charged particles
- ↓ Focusing for negatively charged particles
- ↑ Defocusing for negatively charged particles

Let Us Repeat...


- Plasma wakefields require: plasma, energy source (driver)
 - Place a particle beam (witness) to be accelerated
- Plasma acts as a transformer
 - Drive beam energy is transferred to the witness bunch



State-of-the-Art Results










State-of-the-Art Results

BELLA (Berkeley, California)



Petawatt Laser Guiding and Electron Beam Acceleration to 8 GeV in a
Laser-Heated Capillary Discharge Waveguide
A. J. Gonsalves et al., Phys. Rev. Lett. 122, 084801







State-of-the-Art Results




SLAC (Stanford, California)



Blumenfeld, I., Clayton, C., Decker, F.J. et al.
Energy doubling of 42 GeV electrons in a metre-scale plasma wakefield accelerator.
Nature **445**, 741–744 (2007).

First Demonstration of a Free Electron Laser Driven by a Plasma Wakefield Accelerator




Free-electron lasing at 27 nanometres based on a laser wakefield accelerator
Wentao Wang et al., *Nature* volume 595, pages 516–520 (2021)

The AWAKE Experiment @CERN



Plasma Wakefield Physics @ CERN



- CERN has very high energetic proton bunches available.
- Idea: use energy stored in the proton bunches to accelerate lighter particles e.g. electrons

Challenge: Bunch Length

Requirement



In order to create plasma wakefields efficiently, the drive bunch length has to be in the order of the plasma wavelength.

Challenge


The SPS proton bunches are ~ 10 cm long, and
The AWAKE plasma wavelength is 1.2 mm.

Solution



The experiment relies on the self-modulation instability
To micro-bunch the long proton beam into micro-bunches.



Formation of the Seeded Self-Modulation




The Seeded Self-Modulation




longitudinal
micro-bunch
structure of the
beam density

transverse
proton motion

The Seeded Self-Modulation




Each micro-bunch drives its own ‘low-amplitude’ wakefield




Resonant wakefield excitation.

Simulation Result



Experimental Realization @CERN

→ AWAKE Experiment



From a concept and an idea to reality !

AWAKE Components



Plasma


- Laser
- Rubidium vapor

Drive Bunch

- Proton beam (400 GeV/c)

Witness Bunch

- Electron beam (10-20 MeV)





750m proton beam line




Diagnostics:

- Proton
- Laser
- Electron





Experimental Layout





The AWAKE Plasma


Rubidium vapour cell.

The laser **ionizes** the outermost electron of each rubidium atom.

Desired **plasma density**: $\sim 1\text{-}10 \times 10^{14}$ electrons/cm³.



The AWAKE Experiment @CERN




- Proton bunch momentum: 400 GeV/c
- 3×10^{11} protons/bunch
- Bunch length: $\sigma_z = \sim 10$ cm
- Radial bunch size at plasma entrance: $\sigma_r = 0.2$ mm

Diagnostics


Relevant measurements:

- What is the plasma density?
- Did the proton beam self-modulated over the 10 m of plasma?
- What is the energy of the accelerated electrons?




Self-Modulation Diagnostics

Streak camera measurement





Foil emits waves up to the plasma wavelength of the foil including:

- radiation in the optical range (OTR).
- Coherent radiation (CTR) for wavelengths bigger than the structure of the micro-bunches




Accelerated Electron Energy Measurement



- Electrons will be injected with an energy around 10-20 MeV.
- Accelerated electrons are sent through a **dipole magnet** and deposit energy on a scintillating screen which is imaged by a camera.



Typical final energy distribution of the accelerated electron beam after 10 m plasma:



A. Petrenko, CERN



Let Us Repeat...


- To realize the AWAKE experiment at CERN, we need:
 - Plasma (vapor source + laser)
 - Proton bunch (wakefield driver)
 - Electron bunch (witness – to be accelerated)
- Diagnostics are key to a successful measurement
 - AWAKE diagnostics include:
 - Screens (to know beam positions and verify that SSM was successful)
 - Streak camera (time resolved images of the proton bunch)
 - Electron spectrometer (energy of the accelerated witness bunch)

AWAKE Experimental Results



AWAKE Run 1 (2016-2018)

1. **Self-modulate** a long (compared λ_{pe}) 400 GeV/c proton bunch in plasma.




2. **Accelerate** externally injected 10- 20 MeV electrons to GeV energies (2018).




Self-Modulation Measurement Results

Plasma off:




Plasma on:






Shortly after we have observed the Seeded-Self Modulation for the first time!

First Electron Acceleration




No electrons accelerated.



Accelerated electrons.

Electron Acceleration Results




AWAKE Collaboration, *Nature* volume 561, pages 363–367 (2018)




**Shortly after we
have observed
electron
acceleration for
the first time!**

Most Recently: New Plasma Source Technology




Coming up Next in AWAKE: New Plasma Source



- Allows to adjust the plasma density along the 10 m
- More stable SSM, → higher wakefield amplitudes

AWAKE until ~2030




First AWAKE Particle Physics Applications

Example I: Dark Matter Experiment

These experiments use the collisions of an electron beam with a fixed-target or a dump to generate the dark photon via Bremsstrahlung (electron and proton beams) or meson production.


The products of the collisions are mostly absorbed in the dump and the dark photon is searched for as a displaced vertex with two opposite charged tracks in the decay volume of the experiment.



From Caldwell: <https://arxiv.org/pdf/1812.11164.pdf>

First AWAKE Particle Physics Applications

Example II: Electron-Proton Collisions




Diagrams of neutral-current (top) and charged-current (bottom) deep-inelastic electron–proton scattering processes. Image credit: DESY.

- **Collide:**

- 50 GeV electrons with 7 TeV LHC protons
- \sim TeV electrons with 7 TeV LHC protons

Plasma-based collider design



Caldwell, A., Wing, M. VHEeP: a very high energy electron–proton collider. *Eur. Phys. J. C* **76**, 463 (2016).
<https://doi.org/10.1140/epjc/s10052-016-4316-1>

Physics cases:

- Study of the sub-structure and spin structure of the proton and photon
- Determine if partons are fundamental point-like objects
- Clarifying the underlying physics leading to the energy dependence of cross sections
- Leptoquark production: hypothetical particles that would interact with quarks and leptons

Summary and Conclusions

- **Plasma wakefield acceleration** is a novel technique to accelerate charged particles
 - Advantage: **Very high accelerating gradient**, compact accelerators
- Proof of principle acceleration has been demonstrated
 - Next step: aim for high beam quality at high repetition rate → First applications
- AWAKE is a proof-of-principle accelerator R&D experiment at CERN:
 - Only proton-driven wakefield acceleration experiment worldwide
 - The experiment opens a pathway towards particle physics applications
- AWAKE uses a:
 - 400 GeV SPS proton beam as drive beam
 - 10-20 MeV electrons as witness beam
 - 4.5 TW laser beam to create the plasma
 - 10 m long rubidium vapor source
- Final Goal: Design high quality & high energy electron accelerator based on acquired knowledge.



home.cern