

# Accessing Orbital Angular Momentum from TMDs and GPDs

Matthias Burkardt

New Mexico State University

September 1, 2011

- Probabilistic interpretation of GPDs as Fourier transforms of impact parameter dependent PDFs
  - $H(x, 0, -\Delta_{\perp}^2) \rightarrow q(x, \mathbf{b}_{\perp})$
  - $\tilde{H}(x, 0, -\Delta_{\perp}^2) \rightarrow \Delta q(x, \mathbf{b}_{\perp})$
  - $E(x, 0, -\Delta_{\perp}^2) \rightarrow \perp$  deformation of PDFs when the target is  $\perp$  polarized





↪ Ji relation

- Chromodynamik lensing and  $\perp$  single-spin asymmetries (SSA)

transverse distortion of PDFs  
+ final state interactions }  $\Rightarrow$

↪ SSA in  $\gamma N \rightarrow \pi + X$

- quark-gluon correlations  $\rightarrow \perp$  force on  $q$  in DIS
- Summary



- form factors:  $\overleftrightarrow{FT} \rho(\vec{r})$
- $GPDs(x, \vec{\Delta})$ : form factor for quarks with momentum fraction  $x$
- ↪ suitable FT of  $GPDs$  should provide spatial distribution of quarks with momentum fraction  $x$
- careful: cannot measure longitudinal momentum ( $x$ ) and longitudinal position simultaneously (Heisenberg)
- ↪ consider purely transverse momentum transfer





### Impact Parameter Dependent Quark Distributions

$$q(x, \mathbf{b}_\perp) = \int \frac{d^2 \Delta_\perp}{(2\pi)^2} H(x, \xi = 0, -\Delta_\perp^2) e^{-i\mathbf{b}_\perp \cdot \Delta_\perp}$$

$q(x, \mathbf{b}_\perp)$  = parton distribution as a function of the separation  $\mathbf{b}_\perp$  from the transverse center of momentum  $\mathbf{R}_\perp \equiv \sum_{i \in q, g} \mathbf{r}_{\perp, i} x_i$   
 MB, Phys. Rev. D62, 071503 (2000)



- No relativistic corrections (Galilean subgroup!)
- ↪ corollary: interpretation of 2d-FT of  $F_1(Q^2)$  as charge density in transverse plane also free of relativistic corrections
- probabilistic interpretation

$q(x, \mathbf{b}_\perp)$  for unpol. p



### unpolarized proton

- $q(x, \mathbf{b}_\perp) = \int \frac{d^2\Delta_\perp}{(2\pi)^2} H(x, 0, -\Delta_\perp^2) e^{-i\mathbf{b}_\perp \cdot \Delta_\perp}$
  - $x$  = momentum fraction of the quark
  - $\vec{b}_\perp$  =  $\perp$  distance of quark from  $\perp$  center of momentum
  - small  $x$ : large 'meson cloud'
  - larger  $x$ : compact 'valence core'
  - $x \rightarrow 1$ : active quark becomes center of momentum
- $\hookrightarrow \vec{b}_\perp \rightarrow 0$  (narrow distribution) for  $x \rightarrow 1$




proton polarized in  $+\hat{x}$  direction

no axial symmetry!

$$q(x, \mathbf{b}_\perp) = \int \frac{d^2 \Delta_\perp}{(2\pi)^2} H_q(x, 0, -\Delta_\perp^2) e^{-i\mathbf{b}_\perp \cdot \Delta_\perp} \\ - \frac{1}{2M} \frac{\partial}{\partial b_y} \int \frac{d^2 \Delta_\perp}{(2\pi)^2} E_q(x, 0, -\Delta_\perp^2) e^{-i\mathbf{b}_\perp \cdot \Delta_\perp}$$

Physics: relevant density in DIS is  $j^+ \equiv j^0 + j^3$  and left-right asymmetry from  $j^3$




proton polarized in  $+\hat{x}$  direction

$$q(x, \mathbf{b}_\perp) = \int \frac{d^2 \Delta_\perp}{(2\pi)^2} H_q(x, 0, -\Delta_\perp^2) e^{-i\mathbf{b}_\perp \cdot \Delta_\perp} \\ - \frac{1}{2M} \frac{\partial}{\partial b_y} \int \frac{d^2 \Delta_\perp}{(2\pi)^2} E_q(x, 0, -\Delta_\perp^2) e^{-i\mathbf{b}_\perp \cdot \Delta_\perp}$$

sign & magnitude of the average shift

model-independently related to p/n  
anomalous magnetic moments:

$$\langle b_y^q \rangle \equiv \int dx \int d^2 b_\perp q(x, \mathbf{b}_\perp) b_y \\ = \frac{1}{2M} \int dx E_q(x, 0, 0) = \frac{\kappa_q}{2M}$$



sign & magnitude of the average shift  
model-independently related to p/n  
anomalous magnetic moments:

$$\begin{aligned}\langle b_y^q \rangle &\equiv \int dx \int d^2 b_\perp q(x, \mathbf{b}_\perp) b_y \\ &= \frac{1}{2M} \int dx E_q(x, 0, 0) = \frac{\kappa_q}{2M}\end{aligned}$$

$$\kappa^P = 1.913 = \frac{2}{3}\kappa_u^P - \frac{1}{3}\kappa_d^P + \dots$$


- $u$ -quarks:  $\kappa_u^P = 2\kappa_p + \kappa_n = 1.673$

↪ shift in  $+\hat{y}$  direction

- $d$ -quarks:  $\kappa_d^P = 2\kappa_n + \kappa_p = -2.033$

↪ shift in  $-\hat{y}$  direction

- $\langle b_y^q \rangle = \mathcal{O}(\pm 0.2 \text{ fm})$  !!!!




### anomalous gravito-magnetic moment

- $B(0) \equiv \sum_{i \in q, g} B_i(0) = 0$   
( $\rightarrow$  S.J. Brodsky)
  - $B_i(0)$  = shift of CoM for flavor  $i$  in  $\perp$  pol. nucleon relative to nucleon CoM
- $\hookrightarrow B(0) = 0 \iff$  CoM of quarks and gluons sum up to CoM of nucleon
- lattice (LHPC, QCDSF)  $u$  and  $d$  quarks seem to almost saturate  $\sum_{i \in q, g} B_i(0) = 0$



example: semi-inclusive deep-inelastic scattering (SIDIS)  $\gamma p \rightarrow \pi X$



- $u, d$  distributions in  $\perp$  polarized proton have left-right asymmetry in  $\perp$  position space (T-even!); sign 'determined' by  $\kappa_u$  &  $\kappa_d$
  - attractive FSI deflects active quark towards the CoM
- $\hookrightarrow$  FSI translates position space distortion (before the quark is knocked out) in  $+\hat{y}$ -direction into momentum asymmetry that favors  $-\hat{y}$  direction  $\rightarrow$  'chromodynamic lensing'

$\Rightarrow$

$\kappa_p, \kappa_n \longleftrightarrow$  sign of SSA!!!!!!! (MB,2004)

- confirmed by HERMES (and recent COMPASS)  $p$  data; consistent with vanishing isoscalar Sivers (COMPASS)

- treat FSI to lowest order in  $g$

$\hookrightarrow$

$$\langle k_q^i \rangle = -\frac{g}{4p^+} \int \frac{d^2 \mathbf{b}_\perp}{2\pi} \frac{b^i}{|\mathbf{b}_\perp|^2} \left\langle p, s \left| \bar{q}(0) \gamma^+ \frac{\lambda_a}{2} q(0) \rho_a(\mathbf{b}_\perp) \right| p, s \right\rangle$$

with  $\rho_a(\mathbf{b}_\perp) = \int dr^- \rho_a(r^-, \mathbf{b}_\perp)$  summed over all quarks and gluons

$\hookrightarrow$  SSA related to dipole moment of parton density-density correlations


- similar density-density correlations as appear in multiparton correlations ( $\rightarrow$  M.Diehl)

- GPDs (N polarized in  $+\hat{x}$  direction):  $u \rightarrow +\hat{y}$  and  $d \rightarrow -\hat{y}$

$\hookrightarrow$  expect density density correlation to show same asymmetry  $\langle b^y \bar{u}(0) \gamma^+ \frac{\lambda_a}{2} u(0) \rho_a(\mathbf{b}_\perp) \rangle > 0$

$\hookrightarrow$  sign of SSA opposite to sign of distortion in position space

'Chromodynamic lensing' mechanism for  $\perp$  SSA requires long range coherence of color field!



before 'dressing'

active quark 'dressed' with glue

QCD-evolution: long-range color decoherence:

- after 'dressing' itself with a gluon, previously **red** quark more likely to be **blue** or **green**

↪ attraction to far-away spectators mostly gone

- only attracted to close-by (high  $Q^2$ )  $g$  from dressing
- high  $Q^2$ :  $q$  at low  $x$  likely to have dressed itself with perturbative gluon!

↪ 'Chromodynamic lensing' mechanism suppressed for high  $Q^2$  & small  $x$ ?

sea quarks

- consider pert.  $\bar{q}$  from splitting of *glue*
- $f_{1T,g}^\perp$  expected to be small due to sum rule

$$\sum_{i \in q, g} \int dx \int d^2 \mathbf{k}_\perp \mathbf{k}_\perp^2 f_{1T}^{\perp, i}(x, \mathbf{k}_\perp) = 0$$

- expect  $f_{1T, \bar{q}}^\perp$  also to be small for pert.  $\bar{q}$

## Total (Spin+Orbital) Quark Angular Momentum

$$J_q^x = L_q^x + S_q^x = \int d^3r [yT_q^{0z}(\vec{r}) - zT_q^{0y}(\vec{r})]$$

- $T_q^{\mu\nu}(\vec{r})$  energy momentum tensor ( $T_q^{\mu\nu}(\vec{r}) = T_q^{\nu\mu}(\vec{r})$ )
- $T_q^{0i}(\vec{r})$  momentum density [ $P_q^i = \int d^3r T_q^{0i}(\vec{r})$ ]
- think:  $(\vec{r} \times \vec{p})^x = yp^z - zp^y$

relate to impact parameter dependent quark distributions  $q(x, \mathbf{r}_\perp)$ :

Consider spherically symmetric wave packet with nucleon polarized in  $+\hat{x}$  direction

- eigenstate under rotations about  $x$ -axis

$\hookrightarrow$  both terms in  $J_q^x$  equal:

$$J_q^x = 2 \int d^3r y T_q^{0z}(\vec{r}) = \int d^3r y [T_q^{0z}(\vec{r}) + T_q^{z0}(\vec{r})]$$

- $\int d^3r y T_q^{00}(\vec{r}) = 0 = \int d^3r y T_q^{zz}(\vec{r})$

$$\Rightarrow J_q^x = \int d^3r y T_q^{++}(\vec{r}) \quad \text{with} \quad T^{++} \equiv T^{00} + T^{0z} + T^{z0} + T^{zz}$$

relate to impact parameter dependent quark distributions  $q(x, \mathbf{r}_\perp)$ :

Consider spherically symmetric wave packet with nucleon polarized in  $+\hat{x}$  direction

- eigenstate under rotations about  $x$ -axis

↪ both terms in  $J_q^x$  equal:

$$J_q^x = 2 \int d^3r y T_q^{0z}(\vec{r}) = \int d^3r y [T_q^{0z}(\vec{r}) + T_q^{z0}(\vec{r})]$$

- $\int d^3r y T_q^{00}(\vec{r}) = 0 = \int d^3r y T_q^{zz}(\vec{r})$

$$\Rightarrow J_q^x = \int d^3r y T_q^{++}(\vec{r}) \quad \text{with} \quad T^{++} \equiv T^{00} + T^{0z} + T^{z0} + T^{zz}$$

- $\int dx x q(x, \mathbf{r}_\perp) = \frac{1}{2m_N} \int dz T^{++}(\vec{r})$   
(note: here  $x$  is momentum fraction and not  $r^x$ )

↪  $J_q^x = m_N \int dx x r^y q(x, \mathbf{r}_\perp)$

- before applying this result to  $\perp$  shifted PDFs, need to consider 'overall  $\perp$  shift' of CoM for  $\perp$  polarized target...

spherically symmetric wave packet has center of momentum off-center:

- relativistic effect  $\rightarrow$  use Dirac wave packet for nucleon

$$\psi = \left( \begin{array}{c} f(r) \\ \frac{\vec{\sigma} \cdot \vec{p}}{E + M_N} f(r) \end{array} \right) \chi \quad \text{with} \quad \chi = \frac{1}{\sqrt{2}} \left( \begin{array}{c} 1 \\ 1 \end{array} \right)$$

$\int d^3r f^2(r) = 1$ , take limit of large 'radius' for wave packet

- evaluate  $T_q^{0z} = \frac{i}{2} \bar{q} (\gamma^0 \partial^z + \gamma^z \partial^0) q$  in this state
- $\psi^\dagger \partial_z \psi$  even under  $y \rightarrow -y$ , i.e. no contribution to  $\langle y T_q^{0z} \rangle$
- use  $i\psi^\dagger \gamma^0 \gamma^z \partial^0 \psi = E\psi^\dagger \gamma^0 \gamma^z \psi$

$$\begin{aligned} \langle T^{0z} y \rangle &= E \int d^3r \psi^\dagger \gamma^0 \gamma^z \psi y = E \int d^3r \psi^\dagger \left( \begin{array}{cc} 0 & \sigma^z \\ \sigma^z & 0 \end{array} \right) \psi y \\ &= \frac{2E}{E + M_N} \int d^3r \chi^\dagger \sigma^z \sigma^y \chi f(r) (-i) \partial^y f(r) y \\ &= \frac{E}{E + M_N} \int d^3r f^2(r) \xrightarrow{R \rightarrow \infty} \frac{1}{2} \end{aligned}$$

$\hookrightarrow$   $p$  pol. in  $+\hat{x}$  direction has CoM shifted by  $\frac{1}{2M_N}$  in  $+\hat{y}$  direction!

spherically symmetric wave packet has center of momentum off-center:

- relativistic effect  $\rightarrow$  use Dirac wave packet for nucleon

$$\psi = \begin{pmatrix} f(r) \\ \frac{\vec{\sigma} \cdot \vec{p}}{E + M_N} f(r) \end{pmatrix} \chi \quad \text{with} \quad \chi = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$


$$\langle T^{0z} y \rangle \xrightarrow{R \rightarrow \infty} \frac{1}{2}$$

$\hookrightarrow p$  pol. in  $+\hat{x}$  direction has CoM shifted by  $\frac{1}{2M_N}$  in  $+\hat{y}$  direction!

origin of 'shift' of CoM

- nucleon polarization:  $\odot$
- counterclockwise momentum density from lower component
- $p \sim \frac{1}{R}$ , but  $y \sim R$

$\hookrightarrow \langle T^{++} y \rangle = \mathcal{O}(1)$



relate to impact parameter dependent quark distributions  $q(x, \mathbf{b}_\perp)$ :

- Thus  $J_q^x = m_N \int dx x r^y q(x, \mathbf{r}_\perp)$  with  $b^y = r^y - \frac{1}{2m_N}$ , where  $q(x, \mathbf{r}_\perp)$  is distribution relative to CoM of whole nucleon
- recall:  $q(x, \mathbf{b}_\perp)$  for nucleon polarized in  $+\hat{x}$  direction

$$q(x, \mathbf{b}_\perp) = \int \frac{d^2 \Delta_\perp}{(2\pi)^2} H_q(x, 0, -\Delta_\perp^2) e^{-i\mathbf{b}_\perp \cdot \Delta_\perp} \\ - \frac{1}{2M_N} \frac{\partial}{\partial b_y} \int \frac{d^2 \Delta_\perp}{(2\pi)^2} E_q(x, 0, -\Delta_\perp^2) e^{-i\mathbf{b}_\perp \cdot \Delta_\perp}$$

$$\Rightarrow J_q^x = M_N \int dx x r^y q(x, \mathbf{r}_\perp) = \int dx x \left( m_N b^y + \frac{1}{2} \right) q(x, \mathbf{r}_\perp) \\ = \frac{1}{2} \int dx x [H(x, 0, 0) + E(x, 0, 0)]$$



relate to impact parameter dependent quark distributions  $q(x, \mathbf{b}_\perp)$ :


- Thus  $J_q^x = m_N \int dx x r^y q(x, \mathbf{r}_\perp)$  with  $b^y = r^y - \frac{1}{2m_N}$ , where  $q(x, \mathbf{r}_\perp)$  is distribution relative to CoM of whole nucleon
- recall:  $q(x, \mathbf{b}_\perp)$  for nucleon polarized in  $+\hat{x}$  direction

$$q(x, \mathbf{b}_\perp) = \int \frac{d^2 \Delta_\perp}{(2\pi)^2} H_q(x, 0, -\Delta_\perp^2) e^{-i\mathbf{b}_\perp \cdot \Delta_\perp} \\ - \frac{1}{2M_N} \frac{\partial}{\partial b_y} \int \frac{d^2 \Delta_\perp}{(2\pi)^2} E_q(x, 0, -\Delta_\perp^2) e^{-i\mathbf{b}_\perp \cdot \Delta_\perp}$$

$$\Rightarrow J_q^x = M_N \int dx x r^y q(x, \mathbf{r}_\perp) = \int dx x \left( m_N b^y + \frac{1}{2} \right) q(x, \mathbf{r}_\perp) \\ = \frac{1}{2} \int dx x [H(x, 0, 0) + E(x, 0, 0)]$$

- X.Ji (1996): rotational invariance  $\Rightarrow$  apply to all components of  $\vec{J}$
- partonic interpretation exists only for  $\perp$  components!

## Ji decomposition



$$\frac{1}{2} = \sum_q \frac{1}{2} \Delta q + L_q + J_g$$

$$\frac{1}{2} \Delta q = \frac{1}{2} \int d^3x \langle P, S | q^\dagger(\vec{x}) \Sigma^3 q(\vec{x}) | P, S \rangle$$

$$L_q = \int d^3x \langle P, S | q^\dagger(\vec{x}) (\vec{x} \times i\vec{D})^3 q(\vec{x}) | P, S \rangle$$

$$J_g = \int d^3x \langle P, S | [\vec{x} \times (\vec{E} \times \vec{B})]^3 | P, S \rangle$$

- $i\vec{D} = i\vec{\partial} - g\vec{A}$

## Ji decomposition

- $\Delta q$  from polarized DIS
- $J_q \equiv \frac{1}{2} \Delta q + L_q = \frac{1}{2} \int_0^1 dx [H_q(x, 0, 0) + E_q(x, 0, 0)]$  from DVCS
- $J_g$  in principle from gluon-GPDs; in practice  $J_g = \frac{1}{2} - J_q$
- spin 1  $\rightarrow$  Goldstein, Liuti, ..


$$\begin{aligned}
 \vec{J}_\gamma &= \int d^3r \vec{r} \times (\vec{E} \times \vec{B}) = \int d^3r \vec{r} \times [\vec{E} \times (\vec{\nabla} \times \vec{A})] \\
 &= \int d^3r \left[ E^j (\vec{r} \times \vec{\nabla}) A^j - \vec{r} \times (\vec{E} \cdot \vec{\nabla}) \vec{A} \right] \\
 &= \int d^3r \left[ E^j (\vec{r} \times \vec{\nabla}) A^j + (\vec{r} \times \vec{A}) \vec{\nabla} \cdot \vec{E} + \vec{E} \times \vec{A} \right]
 \end{aligned}$$

- replace  $2^{nd}$  term (eq. of motion  $\vec{\nabla} \cdot \vec{E} = ej^0 = e\psi^\dagger\psi$ ), yielding

$$\vec{J}_\gamma = \int d^3r \left[ \psi^\dagger \vec{r} \times e\vec{A}\psi + E^j (\vec{x} \times \vec{\nabla}) A^j + \vec{E} \times \vec{A} \right]$$

- $\psi^\dagger \vec{r} \times e\vec{A}\psi$  cancels similar term in electron OAM  $\psi^\dagger \vec{r} \times (\vec{p} - e\vec{A})\psi$
- ↪ decomposing  $\vec{J}_\gamma$  into spin and orbital also shuffles angular momentum from photons to electrons!

## Ji decomposition



$$\frac{1}{2} = \sum_q \frac{1}{2} \Delta q + L_q + J_g$$


$$\frac{1}{2} \Delta q = \frac{1}{2} \int d^3x \langle P, S | q^\dagger(\vec{x}) \Sigma^3 q(\vec{x}) | P, S \rangle$$

$$L_q = \int d^3x \langle P, S | q^\dagger(\vec{x}) (\vec{x} \times i\vec{D})^3 q(\vec{x}) | P, S \rangle$$

$$J_g = \int d^3x \langle P, S | \left[ \vec{x} \times (\vec{E} \times \vec{B}) \right]^3 | P, S \rangle$$

- $i\vec{D} = i\vec{\partial} - g\vec{A}$

## Jaffe decomposition



light-cone framework & gauge  $A^+ = 0$

$$\frac{1}{2} = \sum_q \frac{1}{2} \Delta q + \mathcal{L}_q + \Delta G + \mathcal{L}_g$$

$$\mathcal{L}_q = \int d^3r \langle P, S | \bar{q}(\vec{r}) \gamma^+ (\vec{r} \times i\vec{\partial})^z q(\vec{r}) | P, S \rangle$$


$$\Delta G = \varepsilon^{+-ij} \int d^3r \langle P, S | \text{Tr} F^{+i} A^j | P, S \rangle$$

$$\mathcal{L}_g = 2 \int d^3r \langle P, S | \text{Tr} F^{+j} (\vec{x} \times i\vec{\partial})^z A^j | P, S \rangle$$

## Jaffe decomposition

- $\Delta q$  from polarized DIS
- $\Delta G$  from  $\Delta g(x)$   
( $\vec{p} \leftarrow \vec{p}$  &  $\frac{d}{d \ln Q^2} \Delta q(x)$ )
- $\Delta G$  gauge invariant! Nonlocal for  $A^+ \neq 0$
- no exp./lattice access to  $\mathcal{L}_q, \mathcal{L}_g$
- only  $\mathcal{L} \equiv \mathcal{L}_g + \sum_q \mathcal{L}_q$ , by subtraction  
 $\mathcal{L} = \frac{1}{2} - \Delta G - \sum_q \frac{1}{2} \Delta q$

## Jaffe decomposition



light-cone framework & gauge  $A^+ = 0$

$$\frac{1}{2} = \sum_q \frac{1}{2} \Delta q + \mathcal{L}_q + \Delta G + \mathcal{L}_g$$

$$\mathcal{L}_q = \int d^3 r \langle P, S | \bar{q}(\vec{r}) \gamma^+ (\vec{r} \times i\vec{\partial})^z q(\vec{r}) | P, S \rangle$$

$$\Delta G = \varepsilon^{+-ij} \int d^3 r \langle P, S | \text{Tr} F^{+i} A^j | P, S \rangle$$

$$\mathcal{L}_g = 2 \int d^3 r \langle P, S | \text{Tr} F^{+j} (\vec{x} \times i\vec{\partial})^z A^j | P, S \rangle$$

- $L_q$  matrix element of

$$q^\dagger \left[ \vec{r} \times \left( i\vec{\partial} - g\vec{A} \right) \right]^z q = \bar{q} \gamma^0 \left[ \vec{r} \times \left( i\vec{\partial} - g\vec{A} \right) \right]^z q$$


- $\mathcal{L}_q^z$  matrix element of  $(\gamma^+ = \gamma^0 + \gamma^z)$

$$\bar{q} \gamma^+ \left[ \vec{r} \times i\vec{\partial} \right]^z q \Big|_{A^+=0}$$

- (for  $\vec{p} = 0$ ) matrix element of  $\bar{q} \gamma^z \left[ \vec{r} \times \left( i\vec{\partial} - g\vec{A} \right) \right]^z q$  vanishes (parity!)
- ↪  $L_q$  identical to matrix element of  $\bar{q} \gamma^+ \left[ \vec{r} \times \left( i\vec{\partial} - g\vec{A} \right) \right]^z q$  (nucleon at rest)
- ↪ even in light-cone gauge,  $L_q^z$  and  $\mathcal{L}_q^z$  still differ by matrix element of  $q^\dagger \left( \vec{r} \times g\vec{A} \right)^z q \Big|_{A^+=0} = q^\dagger (r^x gA^y - r^y gA^x) q \Big|_{A^+=0}$

## scalar diquark model

- 'mother functions'  $\psi_s^S(x, \mathbf{k}_\perp)$
- ↪  $\mathcal{L}_q$  from  $|\psi_s^S(x, \mathbf{k}_\perp)|^2$
- GPDs from overlap integrals of  $\psi^\dagger \psi$
- ↪  $L_q$  from Ji
- $L_q = \mathcal{L}_q$ .  
No surprise since  $L_q - \mathcal{L}_q \sim \langle q^\dagger \vec{r} \times \vec{A} q \rangle$  and no  $\vec{A}$  in scalar diquark model
- $L_q(x) \neq \mathcal{L}_q(x)$



M.B. + Hikmat BC,  
PRD **79**, 071501 (2009)

QED for dressed  $e^-$  in QED

- 'mother functions'  $\psi_{sh}^S(x, \mathbf{k}_\perp)$
- ↪  $\mathcal{L}_q$  from  $|\psi_{sh}^S(x, \mathbf{k}_\perp)|^2$
- GPDs from overlap integrals of  $\psi^\dagger \psi$
- ↪  $L_q$  from Ji
- $\mathcal{L}_e = L_e + \frac{\alpha}{4\pi} \neq L_e$
- Wakamatsu-pizza?
- Goldstein-pizza?

Bakker Leader Trueman





Bakker Leader Trueman

 $\perp$  pol. nucleon at  $p \rightarrow \infty$ 

$$J_q^x = L_q^x + S_q^x = \int d^3r [yT_q^{0z}(\vec{r}) - zT_q^{0y}(\vec{r})]$$

BIG small

$\hookrightarrow p$  pol. in  $+\hat{x}$  direction has CoM shifted by  $\frac{1}{2M_N}$  in  $+\hat{y}$  direction!

- for wave packet centered around origin:


$$J^x \sim \frac{1}{2M_N} P_N \longrightarrow \infty$$

$\hookrightarrow$  subtract that infinity


- quark flavor  $q$  has CoM shifted by

$$\frac{1}{2M_N} \int dx x E(x_q, 0, 0)$$

- $J_q^x \sim \frac{1}{2M_N} \int dx x E(x_q, 0, 0) P_N \longrightarrow \infty$



$q$  with polarization  $\odot$




↪ counterclockwise current from lower component


↪  $q$  distribution shifted to top

unpolarized target

- all  $q$  polns. equally likely



$q$  with polarization  $\odot$




↪ counterclockwise current from lower component


↪  $q$  distribution shifted to top

unpolarized target

•  $q$  with pol.  $\uparrow$  shifted to left



$q$  with polarization  $\odot$




↪ counterclockwise current from lower component


↪  $q$  distribution shifted to top

unpolarized target

•  $q$  with pol.  $\downarrow$  shifted to right



$q$  with polarization  $\odot$




↪ counterclockwise current from lower component


↪  $q$  distribution shifted to top

unpolarized target

•  $q$  with pol. → shifted to top



$q$  with polarization  $\odot$




↪ counterclockwise current from lower component


↪  $q$  distribution shifted to top

unpolarized target


- $q$  with pol.  $\leftarrow$  shifted to bottom




$q$  with polarization  $\odot$



lattice calculations (QCDSF)



unpolarized target



- transversity distribution in unpol. target described by chirally odd GPD  $\bar{E}_T$
- $\bar{E}_T > 0$  for  $u$  &  $d$  (QCDSF)
- connection  $h_1^\perp(x, \mathbf{k}_\perp) \leftrightarrow \bar{E}_T$  similar to  $f_{1T}^\perp(x, \mathbf{k}_\perp) \leftrightarrow E$ .
- ↳  $h_1^\perp(x, \mathbf{k}_\perp) < 0$  for  $u/p, d/p, u/\pi, \bar{d}/\pi, \dots$  (MB+BH, 2008)
- different valence quarks add coherently  $|h_1^\perp| > |f_1^\perp|$  (MB+BH; Musch)

## higher twist in polarized DIS

- $\sigma_{LL} \propto g_1 - \frac{2Mx}{\nu} g_2$
  - $g_1 = \frac{1}{2} \sum_q e_q^2 g_1^q$  with  $g_1^q = q^\uparrow(x) + \bar{q}^\uparrow(x) - q^\downarrow(x) - \bar{q}^\downarrow(x)$
  - $g_2$  involves quark-gluon correlations
- ↪ no parton interpret. as difference between number densities for  $g_2$
- for  $\perp$  pol. target,  $g_1$  &  $g_2$  contribute equally

$$\sigma_{LT} \propto g_T \equiv g_1 + g_2$$

- ↪ 'clean' separation between  $g_2$  and  $\frac{1}{Q^2}$  corrections to  $g_1$
- What can one learn from  $g_2$ ?
  - $g_2 = g_2^{WW} + \bar{g}_2$  with  $g_2^{WW}(x) \equiv -g_1(x) + \int_x^1 \frac{dy}{y} g_1(y)$

$$d_2 \equiv 3 \int dx x^2 \bar{g}_2(x) = \frac{1}{2MP^{+2}S^x} \langle P, S | \bar{q}(0) g G^{+y}(0) \gamma^+ q(0) | P, S \rangle$$



$$d_2 \equiv 3 \int dx x^2 \bar{g}_2(x) = \frac{1}{2MP^{+2}S^x} \langle P, S | \bar{q}(0) g G^{+y}(0) \gamma^+ q(0) | P, S \rangle$$


$$\sqrt{2}G^{+y} = G^{0y} + G^{zy} = -E^y + B^x$$

↪ write  $d_2 = \frac{1}{4} (\chi_E + 2\chi_M)$  with

$$2M^2 \vec{S} \chi_E = \langle P, S | \vec{j}_a \times \vec{E}_a | P, S \rangle$$

$$2M^2 \vec{S} \chi_B = \langle P, S | \vec{j}_a^0 \times \vec{B}_a | P, S \rangle$$

called color-**electric** and **magnetic**  
'polarizabilities'



$$d_2 \equiv 3 \int dx x^2 \bar{g}_2(x) = \frac{1}{2MP^{+2}S^x} \langle P, S | \bar{q}(0) g G^{+y}(0) \gamma^+ q(0) | P, S \rangle$$


$$\sqrt{2}G^{+y} = G^{0y} + G^{zy} = -E^y + B^x$$

↪ write  $d_2 = \frac{1}{4} (\chi_E + 2\chi_M)$  with

$$2M^2 \vec{S} \chi_E = \langle P, S | \vec{j}_a \times \vec{E}_a | P, S \rangle$$

$$2M^2 \vec{S} \chi_B = \langle P, S | \vec{j}_a^0 \times \vec{B}_a | P, S \rangle$$

called color-**electric** and **magnetic**  
'polarizabilities'



names misleading:

these are **not** the response of the nucleon due to an applied color field!

$$d_2 \equiv 3 \int dx x^2 \bar{g}_2(x) = \frac{1}{2MP^{+2}S^x} \langle P, S | \bar{q}(0) g G^{+y}(0) \gamma^+ q(0) | P, S \rangle$$

$$\sqrt{2}G^{+y} = G^{0y} + G^{zy} = -E^y + B^x$$

$$d_2 \equiv 3 \int dx x^2 \bar{g}_2(x) = \frac{1}{2MP^{+2}S^x} \langle P, S | \bar{q}(0) g G^{+y}(0) \gamma^+ q(0) | P, S \rangle$$

$$\sqrt{2}G^{+y} = G^{0y} + G^{zy} = -E^y + B^x = -\left(\vec{E} + \vec{v} \times \vec{B}\right)^y \text{ for } \vec{v} = (0, 0, -1)$$

$\hookrightarrow d_2 \leftrightarrow$  average **color Lorentz force** acting on quark moving with  $v = c$  in  $-\hat{z}$  direction in the instant after being struck by  $\gamma^*$

$$\langle F^y \rangle = -2M^2 d_2 = -\frac{M}{P^{+2}S^x} \langle P, S | \bar{q}(0) g G^{+y}(0) \gamma^+ q(0) | P, S \rangle$$

cf. Qiu-Sterman matrix element  $\langle k_\perp^y \rangle \equiv \int_0^1 dx \int d^2 k_\perp k_\perp^2 f_{1T}^\perp(x, k_\perp^2)$

$$\langle k_\perp^y \rangle = -\frac{1}{2p^+} \left\langle P, S \left| \bar{q}(0) \int_0^\infty dx^- g G^{+y}(x^-) \gamma^+ q(0) \right| P, S \right\rangle$$

semi-classical interpretation: average  $k_\perp$  in SIDIS obtained by correlating the quark density with the transverse impulse acquired from (color) Lorentz force acting on struck quark along its trajectory to (light-cone) infinity

matrix element defining  $d_2$

$\leftrightarrow$

$1^{st}$  integration point in QS-integral

## color Lorentz force

$d_2 \leftrightarrow$  average **color Lorentz force** acting on quark moving with  $v = c$  in  $-\hat{z}$  direction in the instant after being struck by  $\gamma^*$

$$\langle F^y \rangle = -2M^2 d_2 = -\frac{M}{P^{+2} S_x} \langle P, S | \bar{q}(0) g G^{+y}(0) \gamma^+ q(0) | P, S \rangle$$

sign of  $d_2$ 

- $\kappa_q/p \rightarrow$  sign of deformation
- $\hookrightarrow$  direction of average force
- $\hookrightarrow d_2^u > 0, d_2^d < 0$
- cf.  $f_{1T}^{\perp u} < 0, f_{1T}^{\perp d} < 0$

magnitude of  $d_2$ 

- $\langle F^y \rangle = -2M^2 d_2 = -10 \frac{\text{GeV}}{f_m}$
- expect partial cancellation of forces in SSA
- $\hookrightarrow |\langle F^y \rangle| \ll \sigma \approx 1 \frac{\text{GeV}}{f_m}$
- $\hookrightarrow d_2 = \mathcal{O}(0.01)$

## lattice (Göckeler et al., 2005)

$$d_2^u \approx 0.010, d_2^d \approx -0.0056$$

please: up to date lattice calcs.

## color Lorentz force

$e_2 \leftrightarrow$  average **color Lorentz force** (in  $\hat{y}$ -direction) acting on quark (with transversity  $\hat{x}$ ) moving with  $v = c$  in  $-\hat{z}$  direction in the instant after being struck by  $\gamma^*$

$$\langle F^y \rangle = M^2 e_2 \equiv M^2 \int_0^1 dx \bar{e}_2(x) = \frac{M}{4P^{+2}} \sum_{i=1,2} \langle P | \bar{q}(0) g G^{+i}(0) \sigma^{+i} q(0) | P \rangle$$

## chirally even

- GPD  $E_q \Rightarrow \mathbf{b}_\perp$  deformation of unpol.  $q$  distr. in  $\perp$  pol. target

$$\hookrightarrow f_{1T}^\perp$$

$$\hookrightarrow d_2 \equiv \int dx x^2 \bar{g}_2 \text{ force}$$

## chirally odd

- GPD  $\bar{E}_T \Rightarrow \mathbf{b}_\perp$  deformation of quarks with transversity in unpol. target

$$\hookrightarrow h_1^\perp$$

$$\hookrightarrow e_2 \equiv \int dx x^2 \bar{e}_2 \text{ force}$$

## lattice (Göckeler et al., 2005)

$$d_2^u \approx 0.010, d_2^d \approx -0.0056$$

please: up to date lattice calcs.

## lattice

B.Musch ( $\rightarrow$  H.Avakian)

subtraction of  $m_q$  term?

- Deeply Virtual Compton Scattering (DVCS)  $\rightarrow$  GPDs
- $\hookrightarrow$  impact parameter dependent PDFs  $q(x, \mathbf{b}_\perp)$
- $E^q(x, 0, -\Delta_\perp^2) \leftrightarrow \kappa_{q/p}$  (contribution from quark flavor  $q$  to anomalous magnetic moment)
- $E^q(x, 0, -\Delta_\perp^2) \rightarrow \perp$  deformation of PDFs for  $\perp$  polarized target
- $\perp$  deformation  $\leftrightarrow$  (sign of) SSA (Sivers; Boer-Mulders)
- parton interpretation for Ji-relation
- $L_q \neq \mathcal{L}_q$
- higher-twist  $(\int dx x^2 \bar{g}_2(x), \int dx x^2 \bar{e}(x)) \leftrightarrow \perp$  force in DIS
- $\perp$  deformation  $\leftrightarrow$  (sign of) quark-gluon correlations  $(\int dx x^2 \bar{g}_2(x), \int dx x^2 \bar{e}(x))$