

# FAIROS-HEP

Kyle Stuart Cranmer (Wisconsin)  
Mike Hildreth (Notre Dame)  
Peter Elmer (Princeton)



FAIR and Open Science in High Energy Physics  
OAC-2226378, OAC-2226379 and OAC-2226380



# FAIROS-HEP

Recently, the US National Science Foundation funded a new Research Coordination Network project titled “FAIROS-HEP”.

**F**indable

**A**ccessible

**I**nteroperable

**R**eusable

**O**pen **S**cience

# The NSF's FAIROS Research Coordination Networks

Findable, Accessible, Interoperable, Reusable, Open Science Research Coordination Networks (FAIROS RCN) program represent a pooled investment of over \$12.5 million in open science from all directorates comprising NSF.

**FAIROS RCN supports groups of investigators to communicate, innovate, coordinate, and standardize research practices, training, and educational activities across disciplinary, organizational, geographic, and international boundaries to achieve the goals of FAIR and other open-science guiding principles.**

Research coordination networks are a form of awards that NSF makes to advance scientific practices and standards broadly across multiple research fields. These RCN awards will be for **three-year projects**.

# FAIROS-HEP Continues a Legacy of Contributions

DASPOS (2012-2016)

- <https://daspos.crc.nd.edu/>
- Contributions to RECAST led to REANA as a spinoff project now led by CERN
- Supported REANA Common Workflow Language

DIANA-HEP (2015-2021)

- <https://diana-hep.org/>
- Contributions to REANA, RECAST, launched pyhf likelihood publishing, Active Learning for reinterpretation
- Supported GitHub -> Zenodo DOI minting

IRIS-HEP (2018-?)

- <https://iris-hep.org/>
- Major contributions to likelihood publishing, HEPData integration,

SCAILFIN (2018-2021)

- <https://scaifin.github.io/>
- Contributions to REANA (Slurm and HPC backends, applications built on top of REANA, etc.), Active Learning for reinterpretation

FAIROS-HEP (2022-2025)

- <https://fairos-hep.org/> (under construction)
- **Continue the legacy of contributions, help coordinate the ecosystem**

# What is FAIROS-HEP?


The FAIROS-HEP project aims to connect groups of researchers thinking about FAIR data in HEP and other experts in this field to envision a more cohesive infrastructure around data and publications in HEP.

- By focusing on FAIR data practices and **how data and software can be linked to physics results**, we hope to **build a network of researchers** thinking about how we can create a “**living publication**” to preserve and extend physics results.
- The project includes **some funding for building infrastructure** as well as **future workshops connecting groups**.

# Kickoff workshop

The goals of the workshop were to:

1. Assess progress by each experiment in producing reusable data,
2. Establish updated ideas regarding the use cases for data access, interoperability, and reuse across the different experiments and experimental fields,
3. Define what data and associated information supports the use cases, and
4. Identify a preliminary set of access methods and infrastructure that would support these use cases.



The screenshot shows the Indico event page for the FAIROS-HEP Kick-Off Workshop. The page title is "FAIROS-HEP Kick-Off Workshop". The event dates are "8-10 Feb 2023" at "CERN", with the time zone "Europe/Zurich timezone". A search bar is visible in the top right corner. The left sidebar contains a navigation menu with the following items: "Overview" (selected), "Scientific Programme", "Timetable", "Contribution List", "My Conference", and "My Contributions". The main content area contains the following text:

Recently, the US National Science Foundation funded a new Research Coordination Network project titled "FAIROS-HEP". The project aims to connect groups of researchers thinking about FAIR data in HEP and other experts in this field to envision a more cohesive infrastructure around data and publications in HEP. By focusing on FAIR data practices and how data and software can be linked to physics results, we hope to build a network of researchers thinking about how we can create a "living publication" to preserve and extend physics results. The project includes some funding for building infrastructure as well as future workshops connecting groups.

This is our kick-off workshop, located at CERN.

The goals of the workshop will be to:

1. Assess progress by each experiment in producing reusable data,
2. Establish updated ideas regarding the use cases for data access, interoperability, and reuse across the different experiments and experimental fields,
3. Define what data and associated information supports the use cases, and
4. Identify a preliminary set of access methods and infrastructure that would support these use cases.


We look forward to engaging with you in this project.

<https://indico.cern.ch/event/1234612/>

**Technical Recommendations:** Provide initial direction for which elements of the cyber ecosystem will be most relevant for the first round of technical improvements. Initiate investigations.

# IRIS-HEP

The IRIS-HEP Analysis Systems focus area extends to reuse and preservation, but it has not been a major area of activity. FAIROS-HEP will coordinate closely.



# The Kinds of Developments that FAIROS-HEP Would Like to Nurture



# Open Science & Beyond

The field is at a tipping point. CERN has publicly embraced Open Science and the experiments are adopting new policies.

But we also realize Open Data is not the end of the story.  
**Reuse is key!**

nature  
physics

PERSPECTIVE

<https://doi.org/10.1038/s41567-018-0342-2>

Corrected: Publisher Correction

OPEN

## Open is not enough

Xiaoli Chen<sup>1,2</sup>, Sünje Dallmeier-Tiessen<sup>1\*</sup>, Robin Dasler<sup>1,11</sup>, Sebastian Feger<sup>1,3</sup>, Pamfilos Fokianos<sup>1</sup>, Jose Benito Gonzalez<sup>1</sup>, Harri Hirvonsalo<sup>1,4,12</sup>, Dinos Kousidis<sup>1</sup>, Artemis Lavasa<sup>1</sup>, Salvatore Mele<sup>1</sup>, Diego Rodriguez Rodriguez<sup>1</sup>, Tibor Šimko<sup>1\*</sup>, Tim Smith<sup>1</sup>, Ana Trisovic<sup>1,5\*</sup>, Anna Trzcinska<sup>1</sup>, Ioannis Tsanaksidis<sup>1</sup>, Markus Zimmermann<sup>1</sup>, Kyle Cranmer<sup>6</sup>, Lukas Heinrich<sup>6</sup>, Gordon Watts<sup>7</sup>, Michael Hildreth<sup>8</sup>, Lara Lloret Iglesias<sup>9</sup>, Kati Lassila-Perini<sup>4</sup> and Sebastian Neubert<sup>10</sup>

The solutions adopted by the high-energy physics community to foster reproducible research are examples of best practices that could be embraced more widely. This first experience suggests that reproducibility requires going beyond openness.



# Highlight: 2020 CERN homepage

HEPData  
@HEPData

Thanks @KyleCranmer for your support and promotion of @HEPData over several years. Looking forward to future collaboration with @iris\_hep on #pyhf likelihoods and more.

Kyle Cranmer @KyleCranmer · Jan 29

I would like to applaud @STFC\_Matters for funding @HEPData, a vital piece of cyberinfrastructure for HEP. The @NSF has been supporting HEP software and cyberinfrastructure with DASPOS, @diana\_hep and @iris\_hep. @iris\_hep looks forward to collaborating with you! [twitter.com/HEPData/status...](https://twitter.com/HEPData/status...)

1:15 PM · Jan 30, 2020 · Twitter Web App

The screenshot shows the CERN homepage with a navigation bar (ABOUT, NEWS, SCIENCE, RESOURCES, SEARCH, EN) and a large featured article about the CERN Council appointing Fabiola Gianotti. Below this is a 'LATEST NEWS' section with three articles: 'New open release allows theorists to explore...', 'Relive 2019 at CERN', and 'Dive into the world of accelerators'. A blue arrow points from the first article to the right.

Featured on CERN homepage

The screenshot shows a news article on the CERN website. The title is 'New open release allows theorists to explore LHC data in a new way'. The sub-headline reads 'The ATLAS collaboration releases full analysis likelihoods, a first for an LHC experiment'. The article is dated 9 JANUARY, 2020 and is by Katarina Anthony. The main image shows a laptop displaying a website with a plot, next to a small potted plant and a printed document.





News › News › Topic: Knowledge sharing

Voir en [français](#)

## New open release allows theorists to explore LHC data in a new way

The ATLAS collaboration releases full analysis likelihoods, a first for an LHC experiment

9 JANUARY, 2020 | By Katarina Anthony



Courtesy of CERN

## ATLAS releases 'full orchestra' of analysis instruments

01/14/21 | By Stephanie Melchor

The ATLAS collaboration has begun to publish likelihood functions, information that will allow researchers to better understand and use their experiment's data in future analyses.



# Published Probability Models

## Published Probability Models

Updated list of HEPData entries for publications using `histFactory` JSON statistical models:

- Search for charginos and neutralinos in final states with two boosted hadronically decaying bosons and missing transverse momentum in pp collisions at  $\sqrt{s}=13$  TeV with the ATLAS detector. 2021. doi:10.17182/hepdata.104458
- Measurement of the  $t\bar{t}$  production cross section in pp collisions at  $\sqrt{s}=13$  TeV with the ATLAS detector. 2021. doi:10.17182/hepdata.105039
- Search for R-parity violating supersymmetry in a final state containing leptons and many jets with the ATLAS experiment using  $\sqrt{s}=13$  TeV proton-proton collision data. 2021. doi:10.17182/hepdata.104860
- Search for chargino-neutralino pair production in final states with three leptons and missing transverse momentum in  $\sqrt{s}=13$  TeV pp collisions with the ATLAS detector. 2021. doi:10.17182/hepdata.95751
- Measurements of the inclusive and differential production cross sections of a top-quark-antiquark pair in association with a Z boson at  $\sqrt{s}=13$  TeV with the ATLAS detector. 2021. doi:10.17182/hepdata.100351
- Search for pair production of third-generation scalar leptoquarks decaying into a top quark and a  $\tau$ -lepton in pp collisions at  $\sqrt{s}=13$  TeV with the ATLAS detector. 2021. doi:10.17182/hepdata.100174.
- Search for squarks and gluinos in final states with one isolated lepton, jets, and missing transverse momentum at  $\sqrt{s}=13$  TeV with the ATLAS detector. 2021. doi:10.17182/hepdata.97041
- Search for trilepton resonances from chargino and neutralino pair production in  $\sqrt{s}=13$  TeV pp collisions with the ATLAS detector. 2020. doi:10.17182/hepdata.99806.
- Search for displaced leptons in  $\sqrt{s}=13$  TeV pp collisions with the ATLAS detector. 2020. doi:10.17182/hepdata.98796.
- Search for squarks and gluinos in final states with jets and missing transverse momentum using 139 fb<sup>-1</sup> of  $\sqrt{s}=13$  TeV pp collision data with the ATLAS detector. 2020. doi:10.17182/hepdata.95664.
- Measurement of the  $t\bar{t}$  production cross-section in the lepton+jets channel at  $\sqrt{s}=13$  TeV with the ATLAS experiment. 2020. doi:10.17182/hepdata.95748.
- Search for long-lived, massive particles in events with a displaced vertex and a muon with large impact parameter in pp collisions at  $\sqrt{s}=13$  TeV with the ATLAS detector. 2020. doi:10.17182/hepdata.91760
- Search for chargino-neutralino production with mass splittings near the electroweak scale in three-lepton final states in  $\sqrt{s}=13$  TeV pp collisions with the ATLAS detector. 2019. doi:10.17182/hepdata.91127.
- Searches for electroweak production of supersymmetric particles with compressed mass spectra in  $\sqrt{s}=13$  TeV pp collisions with the ATLAS detector. 2019. doi:10.17182/hepdata.91374
- Search for direct stau production in events with two hadronic  $\tau$ -leptons in  $\sqrt{s}=13$  TeV pp collisions with the ATLAS detector. 2019. doi:10.17182/hepdata.92006.
- Search for direct production of electroweakinos in final states with one lepton, missing transverse momentum and a Higgs boson decaying into two b-jets in (pp) collisions at  $\sqrt{s}=13$  TeV with the ATLAS detector. 2019. doi:10.17182/hepdata.90607.
- Search for squarks and gluinos in final states with same-sign leptons and jets using 139 fb<sup>-1</sup> of data collected with the ATLAS detector. 2019. doi:10.17182/hepdata.91214.
- Search for bottom-squark pair production with the ATLAS detector in final states containing Higgs bosons, b-jets and missing transverse momentum. 2019. doi:10.17182/hepdata.89408.

Search for bottom-squark pair production with the ATLAS detector in final states containing Higgs bosons,  $b$ -jets and missing transverse momentum

The ATLAS collaboration

Aad, Georges , Abbott, Brad , Abbott, Dale Charles , Abdinov, Ovsat , Abed Abud, Adam , Abeling, Kira , Abhayasinghe, Deshan Kavishka , Abidi, Syed Haider , Abouzeid, Ossama , Abraham, Nicola

**JHEP 12 (2019) 060, 2019.**

<https://doi.org/10.17182/hepdata.89408.v3>

Journal

INSPIRE

Resources

 HistFactory

# Reuse in <30 lines of code

```
1 import json
2 import cabinetry
3 import pyhf
4 from cabinetry.model_utils import prediction
5 from pyhf.contrib.utils import download
6
7 # download the ATLAS bottom-squarks analysis probability models from HEPData
8 download("https://www.hepdata.net/record/resource/1935437?view=true", "bottom-squarks")
9
10 # construct a workspace from a background-only model and a signal hypothesis
11 bkg_only_workspace = pyhf.Workspace(json.load(open("bottom-squarks/RegionC/BkgOnly.json")))
12 patchset = pyhf.PatchSet(json.load(open("bottom-squarks/RegionC/patchset.json")))
13 workspace = patchset.apply(bkg_only_workspace, "sbottom_600_280_150")
14
15 # construct the probability model and observations
16 model, data = cabinetry.model_utils.model_and_data(workspace)
17
18 # produce visualizations of the pre-fit model and observed data
19 prefit_model = prediction(model)
20 cabinetry.visualize.data_mc(prefit_model, data)
21
22 # fit the model to the observed data
23 fit_results = cabinetry.fit.fit(model, data)
24
25 # produce visualizations of the post-fit model and observed data
26 postfit_model = prediction(model, fit_results=fit_results)
27 cabinetry.visualize.data_mc(postfit_model, data)
```






Figure 3: Pre-fit (left) and post-fit (right) visualizations of a selected signal hypothesis for four signal regions of the ATLAS search [41] of a bottom-squark of mass 600 GeV with a second-lightest neutralino of mass 280 GeV and lightest supersymmetric particle of mass 150 GeV generated from the full statistical models published in Ref. [20] using code from Ref. [40].



## A SModelS interface for pyhf likelihoods

#6

Gaël Alguero (LPSC, Grenoble), Sabine Kraml (LPSC, Grenoble), Wolfgang Waltenberger (Vienna, OAW and Vienna U.) (Sep 3, 2020)

Published in: *Comput.Phys.Commun.* 264 (2021) 107909 • e-Print: [2009.01809 \[hep-ph\]](https://arxiv.org/abs/2009.01809)

[pdf](#) [DOI](#) [cite](#) [claim](#)

[reference search](#)

[20 citations](#)

# Community Contributions

## Reinterpretation and Long-Term Preservation of Data and Code #1

Stephen Bailey (LBL, Berkeley), K.S. Cranmer (Wisconsin U., Madison), Matthew Feickert (Wisconsin U., Madison), Rob Fine (Los Alamos), Sabine Kraml (LPSC, Grenoble) et al. (Sep 16, 2022)

Contribution to: [2022 Snowmass Summer Study](#) · e-Print: [2209.08054](#) [physics.comp-ph]

[pdf](#) [cite](#) [claim](#) [reference search](#) [2 citations](#)

## Data and Analysis Preservation, Recasting, and Reinterpretation #2

Stephen Bailey (LBL, Berkeley), Christian Bierlich (Lund U. (main)), Andy Buckley (Glasgow U.), Jon Butterworth (University Coll. London), Kyle Cranmer (New York U.) et al. (Mar 18, 2022)

Contribution to: [2022 Snowmass Summer Study](#) · e-Print: [2203.10057](#) [hep-ph]

[pdf](#) [cite](#) [claim](#) [reference search](#) [4 citations](#)

## Signal region combination with full and simplified likelihoods in MadAnalysis 5 #2

Gaël Alguero (LPSC, Grenoble and Annecy, LAPTH), Jack Y. Araz (Durham U., IPPP), Benjamin Fuks (Paris, LPTHE), Sabine Kraml (LPSC, Grenoble) (Jun 29, 2022)

Published in: *SciPost Phys.* 14 (2023) 009 · e-Print: [2206.14870](#) [hep-ph]

[pdf](#) [DOI](#) [cite](#) [claim](#) [reference search](#) [1 citation](#)

## Publishing statistical models: Getting the most out of particle physics experiments #4

Kyle Cranmer (New York U.), Sabine Kraml (LPSC, Grenoble), Harrison B. Prosper (Florida State U.), Philip Bechtle (Bonn U.), Florian U. Bernlochner (Bonn U.) et al. (Sep 10, 2021)

Published in: *SciPost Phys.* 12 (2022) 1, 037, *SciPost Phys.* 12 (2022) 037 · e-Print: [2109.04981](#) [hep-ph]

[pdf](#) [DOI](#) [cite](#) [claim](#) [reference search](#) [23 citations](#)

## Reinterpretation of LHC Results for New Physics: Status and Recommendations after Run 2 #8

LHC Reinterpretation Forum Collaboration · Waleed Abdallah (Harish-Chandra Res. Inst. and Cairo U.) et al. (Mar 19, 2020)

Published in: *SciPost Phys.* 9 (2020) 2, 022 · e-Print: [2003.07868](#) [hep-ph]

[pdf](#) [links](#) [DOI](#) [cite](#) [claim](#) [reference search](#) [57 citations](#)

SciPost Physics

Submission

## Publishing statistical models: Getting the most out of particle physics experiments

Kyle Cranmer<sup>1\*</sup>, Sabine Kraml<sup>2†</sup>, Harrison B. Prosper<sup>3§</sup> (editors), Philip Bechtle<sup>4</sup>, Florian U. Bernlochner<sup>4</sup>, Itay M. Bloch<sup>5</sup>, Enzo Canonero<sup>6</sup>, Marcin Chrzascz<sup>7</sup>, Andrea Coccaro<sup>8</sup>, Jan Conrad<sup>9</sup>, Glen Cowan<sup>10</sup>, Matthew Feickert<sup>11</sup>, Nahuel Ferreira Iachellini<sup>12,13</sup>, Andrew Fowlie<sup>14</sup>, Lukas Heinrich<sup>15</sup>, Alexander Held<sup>1</sup>, Thomas Kuhr<sup>3,16</sup>, Anders Kveltestad<sup>17</sup>, Maeve Madigan<sup>18</sup>, Farvah Mahmoudi<sup>15,19</sup>, Knut Dundas Morá<sup>20</sup>, Mark S. Neubauer<sup>11</sup>, Maurizio Pierini<sup>15</sup>, Juan Rojo<sup>8</sup>, Sezen Sekmen<sup>22</sup>, Luca Silvestrini<sup>23</sup>, Veronica Sanz<sup>24,25</sup>, Giordon Stark<sup>26</sup>, Riccardo Torre<sup>8</sup>, Robert Thorne<sup>27</sup>, Wolfgang Waltenberger<sup>28</sup>, Nicholas Wardle<sup>29</sup>, Jonas Wittbrodt<sup>30</sup>

**1** New York University, USA **2** LPSC Grenoble, France **3** Florida State University, USA **4** University of Bonn, Germany **5** School of Physics and Astronomy, Tel-Aviv University, Israel **6** University of Genova, Italy **7** Institute of Nuclear Physics, Polish Academy of Sciences, Krakow, Poland **8** INFN, Sezione di Genova, Italy **9** Oskar Klein Centre, Stockholm University, Sweden **10** Royal Holloway, University of London, UK **11** University of Illinois at Urbana-Champaign, USA **12** Max Planck Institute for Physics, Munich, Germany **13** Exzellenzcluster ORIGINS, Garching, Germany **14** Nanjing Normal University, Nanjing, PRC **15** CERN, Switzerland **16** Ludwig-Maximilians-Universität München, Germany **17** University of Oslo, Norway **18** DAMTP, University of Cambridge, UK **19** Lyon University, France **20** Columbia University 10027, USA **21** VU Amsterdam and Nikhef, The Netherlands **22** Kyungpook National University, Daegu, Korea **23** INFN, Sezione di Roma, Italy **24** University of Sussex, UK **25** IFIC, Universidad de Valencia-CSIC, Spain **26** SCIPP, UC Santa Cruz, CA, USA **27** University College London, UK **28** HEPHY and University of Vienna, Austria **29** Imperial College London, UK **30** Lund University, Sweden

\* kyle.cranmer@nyu.edu, † sabine.kraml@lpsc.in2p3.fr, § harry@hep.fsu.edu  
(mailing list: [open-likelihoods@cern.ch](#))

September 9, 2021

## Abstract

The statistical models used to derive the results of experimental analyses are of incredible scientific value and are essential information for analysis preservation and reuse. In this paper, we make the scientific case for systematically publishing the full statistical models and discuss the technical developments that make this practical. By means of a variety of physics cases — including parton distribution functions, Higgs boson measurements, effective field theory interpretations, direct searches for new physics, heavy flavor physics, direct dark matter detection, world averages, and beyond the Standard Model global fits — we illustrate how detailed information on the statistical modelling can enhance the short- and long-term impact of experimental results.


1

arXiv:2109.04981v1 [hep-ph] 10 Sep 2021

# Preservation & Reinterpretation


First results using the RECAST reinterpretation framework and publishing full statistical likelihoods (using pyhf)





**ATLAS PUB Note**  
ATL-PHYS-PUB-2019-029



5th August 2019




**Reproducing searches for new physics with the ATLAS experiment through publication of full statistical likelihoods**

The ATLAS Collaboration

The ATLAS Collaboration is starting to publicly provide likelihoods associated with statistical fits used in searches for new physics on HEPData. These likelihoods adhere to a specification first defined by the BiStFactory p.d.f. template. This note introduces a JSON schema that fully describes the BiStFactory statistical model and is sufficient to reproduce key results from published ATLAS analyses. This is peer independent of its implementation in ROOT and it can be used to run statistical analysis outside of the ROOT and RooStats/ROOT Framework. The first of these likelihoods published on HEPData is from a search for bottom-squark pair production. Using two independent implementations of the model, one in ROOT and one in pure Python, the limits on the bottom-squark mass are reproduced, underscoring the implementation independence and long-term viability of the archived data.





© 2019 CERN for the benefit of the ATLAS Collaboration. Reproduction of this article or parts of it is allowed as specified in the CC-BY-4.0 license.



**ATLAS PUB Note**  
ATL-PHYS-PUB-2019-032



11th August 2019



**RECAST framework reinterpretation of an ATLAS Dark Matter Search constraining a model of a dark Higgs boson decaying to two  $b$ -quarks**

The ATLAS Collaboration

The reinterpretation of a search for dark matter produced in association with a Higgs boson decaying to  $b$ -quarks performed with RECAST, a software framework designed to facilitate the reinterpretation of existing searches for new physics, is presented. Reinterpretation using RECAST is enabled through the sustainable preservation of the original data analysis as re-executable declarative workflows using modern cloud technologies and integrated with the wider CERN Analysis Preservation efforts. The reinterpretation targets a model predicting dark matter production in association with a hypothetical dark Higgs boson decaying into  $b$ -quarks where the mass of the dark Higgs boson  $m_H$  is a free parameter, necessitating a faithful reinterpretation of the analysis. The dataset has an integrated luminosity of  $79.8 \text{ fb}^{-1}$  and was recorded with the ATLAS detector at the Large Hadron Collider at a centre-of-mass energy of  $\sqrt{s} = 13 \text{ TeV}$ . Constraints on the parameter space of the dark Higgs model for a fixed choice of dark matter mass  $m_\chi = 200 \text{ GeV}$  exclude model configurations with a mediator mass up to  $3.2 \text{ TeV}$ .

© 2019 CERN for the benefit of the ATLAS Collaboration. Reproduction of this article or parts of it is allowed as specified in the CC-BY-4.0 license.


## Scalable ATLAS pMSSM computational workflows using containerised REANA reusable analysis platform

M. Donadoni<sup>1</sup>, M. Feickert<sup>2</sup>, L. Heinrich<sup>3</sup>, Y. Liu<sup>4</sup>, A. Mečoniš<sup>1</sup>, V. Moiseievkov<sup>1</sup>, G. Stark<sup>5</sup>, T. Šimko<sup>1</sup>, M. Vidal García<sup>1</sup>

- <sup>1</sup>CERN, Geneva, Switzerland
- <sup>2</sup>University of Madison, Wisconsin, United States
- <sup>3</sup>Max-Planck-Institut für Physik, München, Germany
- <sup>4</sup>Sun Yat-sen University, Guangzhou, China
- <sup>5</sup>University of California, Santa Cruz, United States

### ATLAS pMSSM searches


We have developed a streamlined framework for large-scale pMSSM reinterpretations of ATLAS analyses of LHC Run 2 using containerised computational workflows. The project is looking to assess the global coverage of BSM physics and requires running numerous computational workflows representing pMSSM model points.



ATLAS pMSSM studies from LHC Run 1, [arXiv:1508.06602](#)  
K. Cranmer, I. Yavin, RECAST [arXiv:1510.0550](#)

### REANA reusable analysis platform


The computational workflows were run at scale using the REANA reusable analysis platform. The workflows typically run on Kubernetes clusters.



The architecture of the REANA cloud platform. Users can use a command-line client and a web interface to submit containerised workflows that are then orchestrated on supported compute backends.

### Selection of analyses

Following ATLAS analysis preservation policies, many ATLAS analyses have been preserved as containerised Yadage workflows. After validation they are added to a curated selection of analyses suitable for the pMSSM study.



ATLAS SUSY analyses preserved on GitLab.

### Scaling out to O(5k) workflows

We have improved the REANA platform scheduling in order to maximise the scheduling throughput of incoming workflows.




A scalability test submitting 200 workflows every 10 minutes. A cluster with 448 cores (top) cannot keep up with the load. A cluster with 1072 cores (bottom) can comfortably hold the incoming workload.

Cluster Size	Throughput
448 cores	~100 workflows/min
1072 cores	~250 workflows/min

The benchmark tests were running in the CERN Computer Centre and on the Google Cloud Platform public cloud. The REANA scheduling parameters were optimised to maximise CPU utilisation for the pMSSM workflows taking into account the three time-consuming staging jobs per one workflow run.

### Computational workflow

One typical containerised workflow consists of three time-consuming staging steps that run in parallel and the fitting steps that run afterwards. The workflow dependency graph is simple; the complexity relies in having to run O(5k) of these workflows in order to cover sufficient number of pMSSM model points.



A typical pMSSM workflow. The computational runtime is about 10 minutes without systematics (real payload) and about 10 hours with all systematics (real payload).

### Conclusions

- ATLAS searches for supersymmetry are effectively preserved with computational workflow recipes enabling their future reuse and reinterpretation.
- We have launched several ATLAS pMSSM workflows on REANA and studied the workflow turnaround throughput rate as a function of increasing Kubernetes cluster size.
- REANA platform has been extended to support the workload of many concurrent workflows. The solution was benchmarked on medium to large clusters (from 500 to 5000 cores).
- It is essential to adapt cluster parameters to the type of workflows in order to ensure best throughput and cluster resource utilisation (CPU per node, RAM per node).
- The developed system is ready to run large-scale ATLAS pMSSM reinterpretations of LHC Run 2 analyses.



# FAIROS-HEP Developed Infrastructure

Explorer About Gallery Help Contact

Perform Fit New Workspace Reset Params Download Plot

Plots

Histogram  Pull Plot

Channels

- WyCR (custom parameters)
- SR 1fj (custom parameters)
- ttyCR (custom parameters)
- SR 0fj (custom parameters)

Parameters Sort  Impact  Name

tW ME generator

ttbar XS

Lumi

hfakeweight unconv 20 TOT

The figure displays four stacked plots, each representing a different channel: WyCR, SR 1fj, ttyCR, and SR 0fj. Each plot consists of two vertically stacked panels. The top panel shows the total number of events per bin, broken down into various physics components represented by different colors in the stacked histogram. The bottom panel shows the ratio of data to model (data/model) per bin, with data points represented by black diamonds and a shaded uncertainty band. The x-axis for all plots is labeled 'bin' and the y-axis for the top panel is 'events'.



# FAIROS-HEP: Future Workshop Trajectory

**Workshop 2:** Broader Community Engagement and Theory Reinterpretation

Attendees: Particle Physics Experimentalists and Theorists

Location: TBD

**Workshop 3:** Specific Reuse Case: Deriving EFT Results from Future LHC Data

Attendees: Participants from the LHC experiments and theorists working on EFT interpretations

Location: US University, TBD

**Workshop 4:** Broader Engagement: CNI and External Science Partners

Attendees: Core RCN members, CNI membership, External Science Partners

Location: Semi-Annual CNI Membership Meeting

**Workshop 5:** Reuse Case: Kinematic RECASTing for New Physics Discovery

Attendees: Participants from the LHC experiments and theorists working on new physics searches

Location: US University

Thank you!

Questions?

# The NSF's FAIROS Research Coordination Networks

We are one of 10 funded projects:

## DBER+ Commons

- FAIR in Education research

## MaRCN

- Open science in materials

## FAIR facilities and instruments

- PIDs for research instruments

## FARR

- Best practices for ML/AI

## Geospatial Big Data Infrastructure

- Environmental research

## Paleobio/zooarchaeology databases

- Community-coordinated resources

## SEEKCommons project

- Bridge social and environ. sciences

## NoCTURN

- Non-clinical tomography

## REPETO

- Reproducibility in CS Education