



TECTOR

Total weight : 14,000 tonnes
 Overall diameter : 15.0 m
 Overall length : 28.7 m
 Magnetic field : 3.8 T

STEEL RETURN YOKE
 12,500 tonnes

SILICON TRACKERS
 Pixel (100x150 μm) $\sim 16\text{m}^2 \sim 66\text{M}$ channels
 Microstrips (80x180 μm) $\sim 200\text{m}^2 \sim 9.6\text{M}$ channels

SUPERCONDUCTING SOLENOID
 Niobium titanium coil carrying $\sim 18,000\text{A}$

MUON CHAMBERS
 Barrel: 250 Drift Tube, 480 Resistive Plate Chambers
 Endcaps: 468 Cathode Strip, 432 Resistive Plate Chambers

PRESHOWER
 Silicon strips $\sim 16\text{m}^2 \sim 137,000$ channels

FORWARD CALORIMETER
 Steel + Quartz fibres $\sim 2,000$ channels

Polarization Effects in Processes of Dimuon Production

V. Shalaev and S. Shmatov

Yerevan. 11.09.2023

CRYSTAL
 ELECTROMAGNETIC
 CALORIMETER (ECAL)
 76,000 scintillating PbWO₄ crystals

HADRON CALORIMETER (HCAL)
 Brass + Plastic scintillator $\sim 7,000$ channels


CONFERENCE ON HIGH ENERGY PHYSICS




The Drell-Yan Process and Angular Coefficients

Why study dimuons at LHC?

- Important Standard model benchmark channel
- Search for new physic
- Can be used to explore proton inner structure
- Important background source for many BSM processes



	mass → ≈2.3 MeV/c² charge → 2/3 spin → 1/2	≈1.275 GeV/c² 2/3 1/2	≈173.07 GeV/c² 2/3 1/2	0 0 1	≈126 GeV/c² 0 0
	u up	c charm	t top	g gluon	H Higgs boson
QUARKS	≈4.8 MeV/c² -1/3 1/2	≈95 MeV/c² -1/3 1/2	≈4.18 GeV/c² -1/3 1/2	0 0 1	
	d down	s strange	b bottom	γ photon	
	0.511 MeV/c² -1 1/2	105.7 MeV/c² -1 1/2	1.777 GeV/c² -1 1/2	91.2 GeV/c² 0 1	
	e electron	μ muon	τ tau	Z Z boson	
LEPTONS	<2.2 eV/c² 0 1/2	<0.17 MeV/c² 0 1/2	<15.5 MeV/c² 0 1/2	80.4 GeV/c² ±1 1	
	ν_e electron neutrino	ν_μ muon neutrino	ν_τ tau neutrino	W W boson	
				Gauge bosons	



$$\frac{d^2\sigma}{d\cos\theta^*d\phi^*} \propto \left[(1 + \cos^2\theta^*) + A_0 \frac{1}{2}(1 - 3\cos^2\theta^*) + A_1 \sin(2\theta^*) \cos\phi^* + A_2 \frac{1}{2} \sin^2\theta^* \cos(2\phi^*) \right. \\ \left. + A_3 \sin\theta^* \cos\phi^* + A_4 \cos\theta^* + A_5 \sin^2\theta^* \sin(2\phi^*) + A_6 \sin(2\theta^*) \sin\phi^* + A_7 \sin\theta^* \sin\phi^* \right].$$


where θ^* and ϕ^* are the polar and azimuthal angles of l^- (e^- or μ^-) in the rest frame of γ^*/Z (Collins-Soper) and coefficients $A_0 - A_7$ are functions of p_T, Y, M kinematic variables, polarised and unpolarized cross sections




The Drell-Yan Process and Angular Coefficients

Why study dimuons at LHC?

- Important Standard model benchmark channel
- Search for new physics
- **Can be used to explore proton inner structure**
- Important background source for many BSM processes



mass →	≈2.3 MeV/c²	≈1.275 GeV/c²	≈173.07 GeV/c²	0	≈126 GeV/c²
charge →	2/3	2/3	2/3	0	0
spin →	1/2	1/2	1/2	1	0
	u up	c charm	t top	g gluon	H Higgs boson
QUARKS					
	≈4.8 MeV/c²	≈95 MeV/c²	≈4.18 GeV/c²	0	
	-1/3	-1/3	-1/3	0	
	1/2	1/2	1/2	1	
	d down	s strange	b bottom	γ photon	
	0.511 MeV/c²	105.7 MeV/c²	1.777 GeV/c²	91.2 GeV/c²	
	-1	-1	-1	0	
	1/2	1/2	1/2	1	
	e electron	μ muon	τ tau	Z Z boson	
LEPTONS					
	<2.2 eV/c²	<0.17 MeV/c²	<15.5 MeV/c²	80.4 GeV/c²	
	0	0	0	±1	
	1/2	1/2	1/2	1	
	ν_e electron neutrino	ν_μ muon neutrino	ν_τ tau neutrino	W W boson	
					GAUGE BOSONS




$$\frac{d^2\sigma}{d\cos\theta^*d\phi^*} \propto \left[(1 + \cos^2\theta^*) + A_0 \frac{1}{2} (1 - 3\cos^2\theta^*) + A_1 \sin(2\theta^*) \cos\phi^* + A_2 \frac{1}{2} \sin^2\theta^* \cos(2\phi^*) \right. \\ \left. + A_3 \sin\theta^* \cos\phi^* + A_4 \cos\theta^* + A_5 \sin^2\theta^* \sin(2\phi^*) + A_6 \sin(2\theta^*) \sin\phi^* + A_7 \sin\theta^* \sin\phi^* \right].$$

where θ^* and ϕ^* are the polar and azimuthal angles of l^- (e^- or μ^-) in the rest frame of γ^*/Z (Collins-Soper) and coefficients $A_0 - A_7$ are functions of p_T, Y, M kinematic variables, polarised and unpolarized cross sections





Process Dynamics








Process Dynamics









Process Dynamics







Process Dynamics






Process Dynamics




$$\frac{d^2\sigma}{d\cos\theta^*d\phi^*} \propto \left[(1 + \cos^2\theta^*) + A_0 \frac{1}{2}(1 - 3\cos^2\theta^*) + A_1 \sin(2\theta^*) \cos\phi^* + A_2 \frac{1}{2} \sin^2\theta^* \cos(2\phi^*) \right. \\ \left. + A_3 \sin\theta^* \cos\phi^* + A_4 \cos\theta^* + A_5 \sin^2\theta^* \sin(2\phi^*) + A_6 \sin(2\theta^*) \sin\phi^* + A_7 \sin\theta^* \sin\phi^* \right].$$








Process Dynamics

- Non zero partons transverse momentum
- Correlations between spin and parton transverse momentum




All A_i depends on PDF's!






Process Dynamics



- Non zero partons transverse momentum
- Correlations between spin and parton transverse momentum



All A_i depends on PDF's!



- Higher orders QCD effects produce complicated P^T distribution of partons




In Z-boson rest frame




Process Dynamics


- Non zero partons transverse momentum
- Correlations between spin and parton transverse momentum




All A_i depends on PDF's!



- LO: $A_0 - A_2 = 0$
- NLO: $A_0 - A_2 > 0$




- Higher orders QCD effects produce complicated P^T distribution of partons







Process Dynamics

- Non zero partons transverse momentum
- Correlations between spin and parton transverse momentum




All A_i depends on PDF's




EPJ C 76 (2016) 325

A_3 to A_7 are sensitive to V-A structure of EWK









- LO: $A_0 - A_2 = 0$
- NLO: $A_0 - A_2 > 0$



In Z-boson rest frame




Measurements of Drell-Yan Angular Coefficients at the LHC



Angular coefficients A_{0-7} (A_{0-4}) are obtained in bins of dilepton (dimuon) transverse momentum and rapidity $80 < M_{\mu+\mu-} < 100$ GeV by ATLAS (CMS) collaboration at 8 TeV

- ✓ Lam-Tung relation $A_0 = A_2$ (related with rotation invariance) violation was observed
- ✓ Forward-Backward Asymmetry A_{FB} was also measured at 7, 8, and 13 TeV
- ✓ Experimental data of CMS and ATLAS experiments are in agreement with each other and with SM NNLO predictions, but some deviations are exist at high p_T^Z
- ✓ A_{0-7} measurements at 13 TeV are in progress (CMS)



Polarization Study with the SPD NICA



- $\sqrt{s_{pp}} = 27 \text{ GeV}$ is not really well known region
- QCD sub-processes not well-described at this energy range
- Polarized beams give a possibility to study Transverse Momentum Dependent PDFs

	unpolarized	longitudinally pol.	transversely pol.
QUARK	f_1 number density		f_{1T}^\perp Sivers
longitudinally pol.		g_{1L} helicity	g_{1T} transversity
transversely pol.	h_1^\perp Boer-Mulders	h_{1L}^\perp pretzelosity	h_1 transversity

Polarization Study with the SPD NICA



- $\sqrt{s_{pp}} = 27 \text{ GeV}$ is not really well known region
- QCD sub-processes not well-described at this energy range
- Polarized beams give a possibility to study Transverse Momentum Dependent PDFs

	unpolarized	longitudinally pol.	transversely pol.
unpolarized	f_1 number density		f_{1T}^\perp Sivers
longitudinally pol.		g_{1L} helicity	g_{1T} pretzelosity
transversely pol.	h_1^\perp Boer-Mulders	h_{1L}^\perp transversity	h_1^\perp pretzelosity

QUARK

unpolarized
longitudinally pol.
transversely pol.

Study of Drell – Yan process at NICA energies is inefficient: $S/B \approx 4.6 \times 10^{-4}$ (before cuts)

The SPD Collaboration made a decision to suspend the study of such reactions

(A. Skachkova Report
31.08.23 Minsk)

Polarization Study with the SPD NICA



- $\sqrt{s_{pp}} = 27 \text{ GeV}$ is not really well known region
- QCD sub-processes not well-described at this energy range
- Polarized beams give a possibility to study Transverse Momentum Dependent PDFs



	unpolarized	longitudinally pol.	transversely pol.
unpolarized	f_1 number density		f_{1T}^\perp Sivers
longitudinally pol.		g_{1L} helicity	g_{1T}
transversely pol.	h_1^\perp Boer-Mulders	h_{1L}^\perp	h_1^\perp transversity h_{1T}^\perp pretzelosity

Study of Drell – Yan process at NICA energies is inefficient: $S/B \approx 4.6 \times 10^{-4}$ (before cuts)

The SPD Collaboration made a decision to suspend the study of such reactions

(A. Skachkova Report 31.08. Minsk)

- ✓ Dominated by gluon-gluon fusion
- ✓ High cross-section
- ✓ J/ψ can be easily reconstructed from the $\mu^+ \mu^-$ - decay
- ✓ Factorization of $c\bar{c}$ pair is not well understood theoretically



Polarization Study with the SPD NICA



- $\sqrt{s_{pp}} = 27 \text{ GeV}$ is not really well known region
- QCD sub-processes not well-described at this energy range
- Polarized beams give a possibility to study Transverse Momentum Dependent PDFs


	unpolarized	longitudinally pol.	transversely pol.
QUARK	f_1 number density		f_{1T}^\perp Sivers
longitudinally pol.		g_{1L} helicity	g_{1T} transversity
transversely pol.	h_1^\perp Boer-Mulders	h_{1L}^\perp pretzelosity	h_{1T}^\perp pretzelosity

Study of Drell – Yan process at NICA energies is inefficient: $S/B \approx 4.6 \times 10^{-4}$ (before cuts)

The SPD Collaboration made a decision to suspend the study of such reactions

(A. Skachkova Report 31.08. Minsk)

- ✓ Dominated by gluon-gluon fusion
- ✓ High cross-section
- ✓ J/ψ can be easily reconstructed from the $\mu^+ \mu^-$ - decay
- ✓ Factorization of $c\bar{c}$ pair is not well understood theoretically




$$\sigma_{\sqrt{s_{pp}}=27 \text{ GeV}}(J/\psi) = 200 \text{ nb}$$

$$\text{Br}: J/\psi \rightarrow \mu^+ \mu^- = 0.05961$$

$$\sigma(J/\psi \rightarrow \mu^+ \mu^-) = 12 \text{ nb}$$

Main background: $\pi^+ \pi^- \rightarrow \mu^+ \mu^-$



Polarization Study with the SPD NICA





- $\sqrt{s_{pp}} = 27 \text{ GeV}$ is not really well known region
- QCD sub-processes not well-described at this energy range
- Polarized beams give a possibility to study Transverse Momentum Dependent PDFs

	unpolarized	longitudinally pol.	transversely pol.
QUARK	f_1 number density		f_{1T}^\perp Sivers
transversely pol. longitudinally pol.		g_{1L} helicity	g_{1T} Boer-Mulders
transversely pol. longitudinally pol.	h_1^\perp Boer-Mulders	h_{1L}^\perp Boer-Mulders	h_{1T}^\perp pretzelocity

Study of Drell – Yan process at NICA energies is inefficient: $S/B \approx 4.6 \times 10^{-4}$ (before cuts)


The SPD Collaboration made a decision to suspend the study of such reactions (A. Skachkova Report 31.08. Minsk)



$$\sigma_{\sqrt{s_{pp}=27 \text{ GeV}}}(J/\psi) = 200 \text{ nb}$$

$$\text{Br}: J/\psi \rightarrow \mu^+ \mu^- = 0.05961$$

$$\sigma(J/\psi \rightarrow \mu^+ \mu^-) = 12 \text{ nb}$$



Could be suppressed with selection conditions!

Main background: $\pi^+ \pi^- \rightarrow \mu^+ \mu^-$

Conclusions



Результаты:

- ✓ Full set of angular polarization coefficients A_{0-7} was measured in Z-peak by ATLAS and CMS (A_{0-4}) in pp-collisions at 8 TeV
- ✓ A_{0-7} measurements at 13 TeV with the CMS is almost done. Estimation of higher order effects is under the process

All measurements (8, 13 TeV) are in a good agreement with a Standard model predictions!

- ✓ Studying of J/ψ production under SPD conditions is started with MC modeling. Possibility to perform this kind of research at NICA is approved

Prospects:

- Publish 13 TeV A_i measurements results
- Studying of J/ψ production with simulation of SPD detector response

Thanks for your attention!



CMS Draft Analysis Note



The content of this note is intended for CMS internal use and distribution only

2021/10/02

Archive Hash: 9583ef3-D

Archive Date: 2021/08/27

The Drell-Yan Angular Coefficients Measurement at 13 TeV

the CMS Collaboration
CERN

Abstract

The polarization of the Z-boson is presented triple differentially, in bins of Z-boson rapidity, transverse momentum and dilepton invariant mass. A data set of Z-bosons decaying to muons at a pp collision energy of 13 TeV and with an integrated luminosity of 138.7 fb^{-1} is used. The polarization of the Z-boson does affect the acceptance for precision measurements and is important for the modeling of the kinematics of leptons, as the polarization governs the Z-boson decay. The seven polarization coefficients that are measured in the Collins-Soper frame. Fair agreement is observed between the data and simulation.



Заключение



Результаты:

- ✓ Полный набор угловых поляризационных коэффициентов A_{0-7} был измерен в Z-пике экспериментами ATLAS и CMS (A_{0-4}) при 8 ТэВ
- ✓ Анализ по измерению A_{0-7} в эксперименте CMS при 13 ТэВ находится на стадии завершения. Исследуется влияние высших порядков с помощью различных Монте-Карло генераторов

Все измерения (8, 13 ТэВ) находятся в хорошем согласии с предсказаниями Стандартной модели!

Перспективы:

- Завершения процедуры утверждения результатов при 13 ТэВ
- Расширение программы исследований с использованием генераторов
- Изучение поляризационных эффектов в процессах рождения J/ψ в эксперименте SPD на коллайдере NICA

Спасибо за внимание!



Эксперимент “Компактный мюонный соленоид”




Компактный мюонной соленоид (CMS) – многоцелевая детекторная система, созданная для работы на пучках Большого адронного коллайдера (LHC) при энергии пучков протонов до 14 ТэВ в с.ц.м. и светимости до $10^{34} \text{ см}^{-2}\text{с}^{-1}$

Детекторные систем спроектированы для измерения физических объектов (фотонов, электронов, мюонов, струй и недостающей энергии) с поперечной энергией до нескольких ТэВ

CMS Integrated Luminosity Delivered, pp

Data included from 2010-03-30 11:22 to 2018-10-26 08:23 UTC



CMS DETECTOR

Total weight : 14,000 tonnes
 Overall diameter : 15.0 m
 Overall length : 28.7 m
 Magnetic field : 3.8 T

STEEL RETURN YOKE
 12,500 tonnes

SILICON TRACKERS
 Pixel (100x150 μm) ~16m² ~66M channels
 Microstrips (80x180 μm) ~200m² ~9.6M channels

SUPERCONDUCTING SOLENOID
 Niobium titanium coil carrying ~18,000A

MUON CHAMBERS
 Barrel: 250 Drift Tube, 480 Resistive Plate Chambers
 Endcaps: 468 Cathode Strip, 432 Resistive Plate Cham

PRESHOWER
 Silicon strips ~16m² ~137,000 cha


FORWARD CALORIMETER
 Steel + Quartz fibres ~2,000 Channels

CRYSTAL ELECTROMAGNETIC CALORIMETER (ECAL)
 ~76,000 scintillating PbWO₄ crystals

HADRON CALORIMETER (HCAL)
 Brass + Plastic scintillator ~7,000 channels

- Длина – 22 метра
- Диаметр – 15 метров
- Магнитное поле – 3.8 T
- Вес – 14 000 тонн


Comparison of the five angular coefficients A_i and A₀ – A₂ measured in the Collins–Soper frame in bins of p_T for |Y| < 1



PLB 750 (2015) 154

- A_i increasing with p_T enlargement (except A₄)
- A₀ ≈ A₂ as it is predicted
- Lum–Tung violation are presented
- Well described by theory
- But...

Comparison of the five angular coefficients A_i and A₀ – A₂ measured in the Collins–Soper frame in bins of p_T for 1 < |Y| < 2.1



PLB 750 (2015) 154

- ... some deviations between data and MC at high p_T!



A_i well described by Standard Model but we should get more precise information about NLO (NNLO) effects at high p_T!



Расчет угловых коэффициентов с помощью различных МК генераторов событий



Для оценки вклада высших порядков в значения A_i были использованы МК-генераторы PYTHIA8, POWHEG и FEWZ. $A_0 - A_7$ извлекались путем аппроксимации распределений событий по $\cos\theta^*, \varphi^*$. Результаты моделирования различных генераторов сравнивались между собой, а также с результатами анализа



- Демонстрируется существенная разница между расчетами в лидирующем порядке (LO), и высшими порядками (NLO и NNLO). Моделирование в NNLO требует более тонкой настройки структурных функций.





Angular Coefficients Values

- At LO only A_4 is non-zero
- $A_0 = A_2$ at LO QCD (Lum-Tung relation) but it is violated at higher orders

Lum-Tung relation can be violated for non-zero p_T of Z-bosons due to:

- higher orders and twists
- QCD vacuum structure



$A_0 - A_2$ related to the Z-boson polarisation
 $A_3 - A_6$ sensitive V-A structure of the couplings in parity violation terms

- A_i is dependent on PDF
- A_4 related to the forward-backward asymmetry.
- $A_5 - A_7$ are expected to become non-zero only NNLO, but are small for NLO processes, usually taken to be zero.



Datasets for Data



Experimental data at $\sqrt{s} = 13 \text{ TeV}$. Results for 2016 data ($\sim 37.2 \text{ fb}^{-1}$) are presented.
Work with the full Run-2 statistic is ongoing

Year	Dataset
2016	/SingleMuon/Run2016B-05Feb2018_ver2-v1/MINIAOD /SingleMuon/Run2016C-05Feb2018-v1/NANOAOD /SingleMuon/Run2016D-05Feb2018-v1/NANOAOD /SingleMuon/Run2016E-05Feb2018-v1/NANOAOD /SingleMuon/Run2016F-05Feb2018-v1/NANOAOD /SingleMuon/Run2016G-05Feb2018-v1/NANOAOD /SingleMuon/Run2016H-05Feb2018_ver2-v1/NANOAOD /SingleMuon/Run2016H-05Feb2018_ver3-v1/NANOAOD
2017	/SingleMuon/Run2017B-31Mar2018-v1/NANOAOD /SingleMuon/Run2017C-31Mar2018-v1/NANOAOD /SingleMuon/Run2017D-31Mar2018-v1/NANOAOD /SingleMuon/Run2017E-31Mar2018-v1/NANOAOD /SingleMuon/Run2017F-31Mar2018-v1/NANOAOD /SingleMuon/Run2017H-31Mar2018-v1/NANOAOD
2018	/SingleMuon/Run2018A-14Sep2018_ver3-v1/NANOAOD /SingleMuon/Run2018B-14Sep2018_ver2-v1/NANOAOD /SingleMuon/Run2018C-14Sep2018_ver1-v1/NANOAOD /SingleMuon/Run2018D-14Sep2018_ver2-v1/NANOAOD

Year	Dataset
2016	/DYJetsToLL_M-50....to Inf_TuneCUETP8M1_13TeV-amcatnl0FXFX- pythia8/RunIISummer16NanoAOD- PUMoriond17_05Feb2018_94X_mcRun2_asymptotic_v2_ext2-v1/NANOAODSIM

Events Selection conditions:

- $p_T^\mu > 20 \text{ GeV}$, $|\eta| < 2.4$
- Tight Muon Selection
- Isolation: $R_{\text{ellso}} < 0.15$, $R=0.4$
- HLT_IsoTkMu24 || HLT_IsoMu24

Binning

- Z region
 - $p_T = 0, 10, 20, 35, 55, 80, 120, 200, 400, \text{inf}$.
 - $|Y| = 0, 1, 2.4$
 - $M = 81 - 101$
- 50toInf
 - $p_T = 0, 10, 20, 40, 600$.
 - $|Y| = 0, 0.35, 0.9, 1.35, 2.4$
 - $M = 50, 81, 101, \text{inf}$ ²⁷


The Collins-Soper coordinate system is chosen in such a way that the Z-axis bisects the angle between the interacting quarks

$$\cos \theta^* = \frac{p_z^{(\ell^+\ell^-)}}{|p_z^{(\ell^+\ell^-)}|} \frac{2}{m(Z/\gamma^*)\sqrt{m(Z/\gamma^*)^2 + p_T(Z/\gamma^*)^2}} (P_1^+ P_2^- - P_1^- P_2^+)$$


$$P_i^\pm = \frac{1}{\sqrt{2}} (E_i \pm p_{z,i}),$$

$$\tan \varphi^* = \frac{\sqrt{M_{l^+l^-}^2 + (p_T^{l^+l^-})^2}}{M_{l^+l^-}} \cdot \frac{\vec{\Delta}_r \cdot \widehat{R}_T}{\vec{\Delta}_r \cdot \widehat{h}}$$

Where $M_{l^+l^-}$ is the dilepton invariant mass, $\vec{\Delta}_r = l^- - l^+$, where l^- and l^+ are the respective four-momenta of the particle (electron, muon) and antiparticle (positron, antimuon), \widehat{h} is a transverse unit vector in the direction of $p_T^{l^+l^-}$ and \widehat{R}_T is a transverse unit vector in the direction $\vec{P}_A \times \vec{Q}$, \vec{P}_A is a vector pointing along the negative z-axis, $\vec{P}_A = (0, 0, -1)$, and \vec{Q} is the four-momentum of the dilepton pair.



Lum-Tung relation ($A_0 = A_2$) is satisfied when $\varphi_1 = 0!$



But $\varphi_1 \neq 0$ at NLO and higher!

Lum-Tung violated



Template Method




CMS measured only first five coefficients (while ATLAS measured all 8 coefficients):

$$\frac{d^2\sigma}{d\theta^* d\phi^*} = \sum_{i=0}^5 \sigma^i = P_5(1 + \cos^2 \theta^*) + P_0 \frac{1}{2}(1 - 3 \cos^2 \theta^*) + P_1 \sin(2\theta^*) \cos \phi^* \\ + P_2 \frac{1}{2} \sin^2 \theta^* \cos(2\phi^*) + P_3 \sin \theta^* \cos \phi^* + P_4 \cos \theta^*$$

P_i coefficients relates with A_i as: $A_i = \frac{P_i}{P_5}$

- Fill $\cos \theta^*$, φ^* histogram at gen and reco level
- Reweight Reco events by $\frac{1 + \cos^2 \theta^*}{N_{gen}(\cos \theta^*, \varphi^*)}$
 $\frac{1 - 3 \cos^2 \theta^*}{2 N_{gen}(\cos \theta^*, \varphi^*)}$, ... to get templates H_i for all of the coefficients. Here we divide by $N_{gen}(\cos \theta^*, \varphi^*)$ to get rid of polarization



- Angular coefficients can be directly obtained by minimizing the objective function:

$$\chi^2 = \frac{\left(\text{data}^{j,k} - \left(\sum_{i=0}^5 P_i H_i^{j,k} + H_{\text{Bkg}}^{j,k} \right) \right)^2}{\text{data}^{j,k}}$$





Templates at 13 TeV (Generator Level)

(MADGRAPH+PYTHIA8, CUETP8M1,NLO)



Templates for the six fit parameters P_0 - P_5 on generator level obtained for the first bin of p_T (10-20 GeV) bin for the rapidity bin $|Y| < 1$



Templates for the six fit parameters P_0 - P_5 on generator level obtained for the first bin of p_T (200-400 GeV) bin for the rapidity bin $1 < |Y| < 2.4$.



Шаблоны получены в восьми бинах p_T и двух бинах быстроты!





Templates at 13 TeV (Generator Level)

(MADGRAPH+PYTHIA8, CUETP8M1)



Templates for the six fit parameters P_0 - P_5 on generator level obtained for the first bin of p_T (0-10 GeV) bin for the rapidity bin $1 < |Y| < 2.4$.




Templates for the six fit parameters P_0 - P_5 on generator level at $\sqrt{s} = 13$ TeV for the same p_T bin and $|Y| < 1$.

Templates obtained for eight p_T and two rapidity bins!




Измерение асимметрии по φ_{CS}^* . Предварительные результаты



В соответствии с теоретической моделью распределение мюонов по углу φ_{CS}^* должно быть симметричным относительно нуля

$$A_{\varphi_{CS}^*} = \frac{N_{\varphi_{CS}^{*-}} - N_{\varphi_{CS}^{*+}}}{N_{\varphi_{CS}^{*-}} + N_{\varphi_{CS}^{*+}}}$$



Асимметрия $A_{\varphi_{CS}^*}$ была впервые измерена в рамках анализа по изучению угловых распределений мюонов в процессе Дрелла-Яна в зависимости от инвариантной массы мюонной пары, в четырех интервалах по скорости и поперечному импульсу.

Измеренные значения соответствуют ожидаемым в рамках используемой модели

Исследование предложено О.В. Теряевым (JINR)