

Contribution ID: 299



Type: Parallel contribution

Higgs Production through Top-prime decays at the LHC

Thursday 11 August 2011 11:30 (20 minutes)

We explore LHC signatures of vectorlike quarks, which are hypothetical fermions whose left- and righthanded components have the same electroweak quantum numbers. We consider interactions of such a quark, top-prime, with the top quark via a Yukawa coupling and with a bottom quark through a W boson. We look at Higgs production through the decay of the top-prime in a top-prime pair production channel through QCD at the LHC with $\sqrt{s} = 7$ TeV. Such a process depends only on the top-prime mass. In this channel, we consider semi-leptonic W-boson decays. This choice is dictated by the reduction of QCD background and a higher cross section than the corresponding much cleaner di-leptonic channel. We suggest a background discrimination strategy involving *b*-tagging and a lepton in the final state. The possibility of the top-prime decaying into a light (120 GeV/ c^2) and a relatively heavier Higgs (150 GeV/ c^2) will be explored. The mixing angle between the top and the top-prime, which is a parameter of the model, has been chosen judiciously so that the analysis remains as model independent as possible.

Primary authors: Prof. DOBRESCU, Bogdan (Theoretical Physics Department, Fermi National Accelerator Laboratory); Prof. NARAIN, Meenakshi (Brown University); Ms BHATTACHARYA, Saptaparna (Brown University)

Presenter: Ms BHATTACHARYA, Saptaparna (Brown University)

Session Classification: Beyond the Standard Model

Track Classification: Beyond the Standard Model