



Contribution ID: 299

Type: **Parallel contribution**

## Higgs Production through Top-prime decays at the LHC

*Thursday, 11 August 2011 11:30 (20 minutes)*

We explore LHC signatures of vectorlike quarks, which are hypothetical fermions whose left- and right-handed components have the same electroweak quantum numbers. We consider interactions of such a quark, top-prime, with the top quark via a Yukawa coupling and with a bottom quark through a  $W$  boson. We look at Higgs production through the decay of the top-prime in a top-prime pair production channel through QCD at the LHC with  $\sqrt{s} = 7$  TeV. Such a process depends only on the top-prime mass. In this channel, we consider semi-leptonic  $W$ -boson decays. This choice is dictated by the reduction of QCD background and a higher cross section than the corresponding much cleaner di-leptonic channel. We suggest a background discrimination strategy involving  $b$ -tagging and a lepton in the final state. The possibility of the top-prime decaying into a light ( $120 \text{ GeV}/c^2$ ) and a relatively heavier Higgs ( $150 \text{ GeV}/c^2$ ) will be explored. The mixing angle between the top and the top-prime, which is a parameter of the model, has been chosen judiciously so that the analysis remains as model independent as possible.

**Primary authors:** Prof. DOBRESCU, Bogdan (Theoretical Physics Department, Fermi National Accelerator Laboratory); Prof. NARAIN, Meenakshi (Brown University); Ms BHATTACHARYA, Saptaparna (Brown University)

**Presenter:** Ms BHATTACHARYA, Saptaparna (Brown University)

**Session Classification:** Beyond the Standard Model

**Track Classification:** Beyond the Standard Model