

American Physical Society Conference  
Division of Particles and Fields  
Brown University, Providence RI, August 9–13, 2011


Measurement of the Cross section for Prompt  
Isolated Diphoton Production in  $p\bar{p}$  Collisions  
at  $\sqrt{s} = 1.96$  TeV

Costas Vellidis


FNAL

On behalf of the CDF Collaboration


# Prompt diphoton production at hadron colliders




- **Prompt photons** = photons produced directly in perturbative scattering or via parton fragmentation (as opposed to non-perturbative photon production in meson decays).
- Main source of prompt diphoton production at hadron colliders via QCD interactions.




# Prompt diphoton production at hadron colliders




- **Prompt photons** = photons produced directly in perturbative scattering or via parton fragmentation (as opposed to non-perturbative photon production in meson decays).
- Main source of prompt diphoton production at hadron colliders via QCD interactions.



- **Main background:  $\gamma$ +jet and dijet**, with one or two jets misidentified as photons  $\rightarrow$  reducible background.




# Prompt diphoton production at hadron colliders




- **Prompt photons** = photons produced directly in perturbative scattering or via parton fragmentation (as opposed to non-perturbative photon production in meson decays).
- At much smaller rate, prompt diphotons may originate from more exotic (and exciting!) production mechanisms:

- Higgs decay




- Extra dimensions




- SUSY

- ...



# Prompt diphoton production at hadron colliders




- **Prompt photons** = photons produced directly in perturbative scattering or via parton fragmentation (as opposed to non-perturbative photon production in meson decays).
- At much smaller rate, prompt diphotons may originate from more exotic (and exciting!) production mechanisms:
  - Higgs decay

Precise measurements of QCD  $\gamma\gamma$  production should put us on solid footing to search for new physics:


- Validate/improve theoretical predictions for irreducible (QCD  $\gamma\gamma$ ) background.
- Develop/demonstrate good control over reducible backgrounds.

- SUSY
- ...



# Previous Tevatron measurements

- CDF publication in Run II with 207 pb<sup>-1</sup>. *PRL 95, 022003 (2005)*
- Event selection:  $p_{T1(2)}=14(13)$  GeV,  $|\eta_{1,2}|<0.9$ ,  $\Delta R(\gamma,\gamma)<0.3$ ,  $E_T^{iso}<1$  GeV.




- $p_T(\gamma\gamma) > 25$  GeV region in data dominated by events with  $p_T(\gamma\gamma) > M(\gamma\gamma)$  and  $\Delta\phi(\gamma,\gamma) < \pi/2 \rightarrow$  potentially large fragmentation contributions.
- Large sensitivity of theoretical prediction on isolation requirement.

Here the Pythia prediction uses only matrix element based production of photons

# Previous Tevatron measurements

- D0 publication in Run II with  $4.2 \text{ fb}^{-1}$  *PLB 690, 108 (2010)*
- $p_{T1(2)}=21(20) \text{ GeV}/c$ ,  $|\eta_{1,2}|<1$ ,  $\Delta R(\gamma,\gamma)>0.4$ ,  $(E_{\text{tot}}^{R=0.4} - E_{\text{em}}^{R=0.2})/E_{\text{em}}^{R=0.2} < 0.1$ ,  $p_T(\gamma\gamma) < M(\gamma\gamma)$




- Good agreement between data and RESBOS for  $M_{\gamma\gamma} > 50 \text{ GeV}/c^2$
- Need for a resummed calculation
- Data spectrum harder than predicted
- Observable nearly insensitive to experimental effects
- Supports conclusion from  $p_T(\gamma\gamma)$  measurement

(\* Overall normalization uncertainty (7.3%) not included in data error bars.

Here the Pythia prediction uses only matrix element based production of photons


# Photon identification and event selection



- Photons are selected offline from EM clusters, reconstructed within a cone of radius  $R=0.4$  in the  $\eta$ - $\phi$  plane, and requiring:
    - Fiducial to the central calorimeter:  $|\eta| < 1.1$  Avoids divergence in NLO calculation
    - $E_T \geq 17$  GeV (1<sup>st</sup>  $\gamma$  in the event), 15 GeV (2<sup>nd</sup>  $\gamma$ )
    - Isolated in the calorimeter:  $I_{\text{cal}} = E_{\text{tot}}(R=0.4) - E_{\text{EM}}(R=0.4) \leq 2$  GeV
    - Low HAD fraction:  $E_{\text{HAD}}/E_{\text{EM}} \leq 0.055 + 0.00045 \times E_{\text{tot}}/\text{GeV}$
    - At most one track in cluster with  $p_T^{\text{trk}} \leq 1$  GeV/c +  $0.005 \times E_T^\gamma/c$
    - Shower profile:  $\chi^2_{\text{CES}} \leq 20$
    - $E_T$  of 2<sup>nd</sup> CES cluster  $\leq 2.4$  GeV +  $0.01 \times E_T$
- } Imply that  $\Delta R(\gamma, \gamma) \geq 0.4$



# Photon identification and event selection



- Photons are selected offline from EM clusters, reconstructed within a cone of radius  $R=0.4$  in the  $\eta$ - $\phi$  plane, and requiring:
    - Fiducial to the central calorimeter:  $|\eta| < 1.1$  Avoids divergence in NLO calculation
    - $E_T \geq 17$  GeV (1<sup>st</sup>  $\gamma$  in the event), 15 GeV (2<sup>nd</sup>  $\gamma$ )
    - Isolated in the calorimeter:  $I_{\text{cal}} = E_{\text{tot}}(R=0.4) - E_{\text{EM}}(R=0.4) \leq 2$  GeV
- } Imply that  $\Delta R(\gamma, \gamma) \geq 0.4$

Selected events correspond to **5.4 fb<sup>-1</sup>** of integrated luminosity

- Shower profile:  $\chi^2_{\text{CES}} \leq 20$
- $E_T$  of 2<sup>nd</sup> CES cluster  $\leq 2.4$  GeV + 0.01  $\times$   $E_T$

## Background

$$\frac{d\sigma}{dX} = \frac{N_{\gamma\gamma}}{\varepsilon \cdot A \cdot L \cdot \Delta}$$

Jets misidentified as photons: dijet and  $\gamma$ +jet

- Fluctuations in jet fragmentation to leading  $\pi^0$  or  $\eta^0$  meson ( $\pi^0, \eta^0 \rightarrow \gamma\gamma$ )
- Normalization and shape estimated from MC using **track isolation**:  $I_{trk} = \sum_{\text{tracks in } R < 0.4}^{|z_{vtx} - z_{trk}| < 5 \text{ cm}} p_T^{trk}$
- Sensitive only to underlying event and jet fragmentation (for fake  $\gamma$ ), immune to multiple interactions (due to z-cut) and calorimeter leakage
- Good resolution in low- $E_T$  region, where background is most important
- Uses charged particles only


## Background

$$\frac{d\sigma}{dX} = \frac{N_{\gamma\gamma}}{\varepsilon \cdot A \cdot L \cdot \Delta}$$

Jets misidentified as photons: dijet and  $\gamma$ +jet

- Fluctuations in jet fragmentation to leading  $\pi^0$  or  $\eta^0$  meson ( $\pi^0, \eta^0 \rightarrow \gamma\gamma$ )
- Normalization and shape estimated from MC using **track isolation**:  $I_{trk} = \sum_{\text{tracks in } R < 0.4}^{|z_{vtx} - z_{trk}| < 5\text{cm}} p_T^{trk}$
- Sensitive only to underlying event and jet fragmentation (for fake  $\gamma$ ), immune to multiple interactions (due to z-cut) and calorimeter leakage
- Good resolution in low- $E_T$  region, where background is most important
- Uses charged particles only

Substantially different shape of signal and background  $I_{trk}$  distributions can be used to characterize true and fake  $\gamma$



## Background estimation: 4×4 matrix method

- Use the track isolation cut for each photon to compute a per-event weight under the different hypotheses ( $\gamma\gamma$ ,  $\gamma$ +jet and dijet):

$$\begin{pmatrix} w_{jj} \\ w_{j\gamma} \\ w_{\gamma j} \\ w_{\gamma\gamma} \end{pmatrix} = E^{-1} \times \begin{pmatrix} w_{ff} \\ w_{fp} \\ w_{pf} \\ w_{pp} \end{pmatrix} \begin{array}{l} \text{Both photons fail} \\ \text{Leading fail, trailing passes} \\ \text{Leading passes, trailing fails} \\ \text{Both photons pass} \end{array}$$



$$E = \begin{pmatrix} (1 - \epsilon_{j1})(1 - \epsilon_{j2}) & (1 - \epsilon_{j1})(1 - \epsilon_{\gamma 2}) & (1 - \epsilon_{\gamma 1})(1 - \epsilon_{j2}) & (1 - \epsilon_{\gamma 1})(1 - \epsilon_{\gamma 2}) \\ (1 - \epsilon_{j1})\epsilon_{j2} & (1 - \epsilon_{j1})\epsilon_{\gamma 2} & (1 - \epsilon_{\gamma 1})\epsilon_{j2} & (1 - \epsilon_{\gamma 1})\epsilon_{\gamma 2} \\ \epsilon_{j1}(1 - \epsilon_{j2}) & \epsilon_{j1}(1 - \epsilon_{\gamma 2}) & \epsilon_{\gamma 1}(1 - \epsilon_{j2}) & \epsilon_{\gamma 1}(1 - \epsilon_{\gamma 2}) \\ \epsilon_{j1}\epsilon_{j2} & \epsilon_{j1}\epsilon_{\gamma 2} & \epsilon_{\gamma 1}\epsilon_{j2} & \epsilon_{\gamma 1}\epsilon_{\gamma 2} \end{pmatrix}$$

- For instance, if leading passes/trailing fails, the event weight is:
- Estimated number of prompt diphoton events bin-by-bin is given by the sum of  $\gamma\gamma$  weights:

$$\begin{pmatrix} w_{ff} \\ w_{fp} \\ w_{pf} \\ w_{pp} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}$$

$$N_{\gamma\gamma} = \sum_{i=1}^{N_{data}} w_{\gamma\gamma}^i$$

# Experimental systematic uncertainties



- Total systematic uncertainty ~15-30%, smoothly varying with the kinematic variables considered
- Main source is background subtraction, followed by overall normalization (efficiencies: 7%; integrated luminosity: 6%; UE correction: 6%)


## Theoretical predictions

- **DIPHOX**: Fixed-order NLO calculation including non-perturbative fragmentations (T. Binoth *et al.*, Phys. Rev. D **63**, 114016 (2001))
- **RESBOS**: Low- $p_T$  resummed calculation smoothly matched to high- $p_T$  NLO (T. Balazs *et al.*, Phys. Rev. D **76**, 013008 (2007))
- **PYTHIA** 6.2.16 parton-shower calculation (no k-factor applied) (T. Sjöstrand *et al.*, Comp. Phys. Comm. **135**, 238 (2001))

# Theoretical predictions

- **DIPHOX**: Fixed-order NLO calculation including non-perturbative fragmentations (T. Binoth *et al.*, Phys. Rev. D **63**, 114016 (2001))
- **RESBOS**: Low- $p_T$  resummed calculation smoothly matched to high- $p_T$  NLO (T. Balazs *et al.*, Phys. Rev. D **76**, 013008 (2007))
- **PYTHIA** 6.2.16 parton-shower calculation (no k-factor applied) (T. Sjöstrand *et al.*, Comp. Phys. Comm. **135**, 238 (2001))


Two separate calculations, one involving (a – b) only (“PYTHIA  $\gamma\gamma$ ”) and one involving (a – d) (“PYTHIA  $\gamma\gamma+\gamma j$ ”), are compared with the data



# Theoretical predictions

- **DIPHOX**: Fixed-order NLO calculation including non-perturbative fragmentations (T. Binoth *et al.*, Phys. Rev. D **63**, 114016 (2001))
- **RESBOS**: Low- $p_T$  resummed calculation smoothly matched to high- $p_T$  NLO (T. Balazs *et al.*, Phys. Rev. D **76**, 013008 (2007))
- **PYTHIA 6.2.16** parton-shower calculation (no k-factor applied) (T. Sjöstrand *et al.*, Comp. Phys. Comm. **135**, 238 (2001))

Two separate calculations, one involving (a – b) only (“PYTHIA  $\gamma\gamma$ ”) and one involving (a – d) (“PYTHIA  $\gamma\gamma+\gamma j$ ”), are compared with the data



- NLO theoretical uncertainties:
  - PDFs: 3-6%; use 44 eigenvectors from CTE6.1M
  - Renormalization/factorization/fragmentation scales: ~10-20% depending on the observable; all scales simultaneously varied by  $\times 2$  up and down




# Theoretical predictions

- **DIPHOX**: Fixed-order NLO calculation including non-perturbative fragmentations (T. Binoth *et al.*, Phys. Rev. D **63**,114016 (2001))
- **RESBOS**: Low- $p_T$  resummed calculation smoothly matched to high- $p_T$  NLO (T. Balazs *et al.*, Phys. Rev. D **76**, 013008 (2007))




- **PYTHIA 6.2.16** p8 (T.Sjöstrand *et al.*, Two separate calculations (a – b) only (“PYTHIA”) (a – d) (“PYTHIA”) the data

	Total cross section (pb)
Data	$12.5 \pm 0.2_{\text{stat}} \pm 3.7_{\text{syst}}$
RESBOS	$11.3 \pm 2.4_{\text{syst}}$
DIPHOX	$10.6 \pm 0.6_{\text{syst}}$
PYTHIA $\gamma\gamma + \gamma j$	9.2
PYTHIA $\gamma\gamma$	5.0



- NLO theoretical uncertainties:
  - PDFs: 3-6%; use 44 eigenvectors from CTE6.1M
  - Renormalization/factorization/fragmentation scales: ~10-20% depending on the observable; all scales simultaneously varied by  $\times 2$  up and down

# Differential cross sections






- Good agreement between data and theory for  $M_{\gamma\gamma} > 30$  GeV/c<sup>2</sup>

- Resummation important for  $p_T(\gamma\gamma) > 20$  GeV/c
- Fragmentations cause excess of data over theory for  $p_T(\gamma\gamma) = 20 - 50$  GeV/c

- Resummation important for  $\Delta\phi_{\gamma\gamma} > 2.2$  rad
- Data spectrum harder than predicted

# Differential cross sections






- Good agreement between data and RESBOS
- Good agreement between data and DIPHOX, except for  $0.7 < z < 0.8$

- Good agreement between data and theory

- Observable sensitive to PDFs
- Good agreement between data and theory, except for  $|\cos\theta^*| \rightarrow 1$

# Differential cross sections for $p_T(\gamma\gamma) < M(\gamma\gamma)$




- Good agreement between data and theory


- “Shoulder” in data for  $p_T(\gamma\gamma) = 20 - 50 \text{ GeV}/c$  significantly reduced

- Discrepancies between data and theory for  $\Delta\phi_{\gamma\gamma} < 1.7 \text{ rad}$  reduced


# Differential cross sections for $p_T(\gamma\gamma) < M(\gamma\gamma)$



- Good agreement between data and RESBOS
- Good agreement between data and DIPHOX, except for  $0.7 < z < 0.8$



- Good agreement between data and theory



- Good agreement between data and theory

## Summary and conclusions

- Reported measurements of differential cross sections for direct diphoton production at  $\sqrt{s}=1.96$  TeV using  $5.4 \text{ fb}^{-1}$ .


$$\frac{d\sigma}{dM_{\gamma\gamma}} \quad \frac{d\sigma}{dp_T^{\gamma\gamma}} \quad \frac{d\sigma}{d\Delta\phi_{\gamma\gamma}} \quad \frac{d\sigma}{dz} \quad \frac{d\sigma}{dy_{\gamma\gamma}} \quad \frac{d\sigma}{d\cos\theta^*}$$

- Measurements are compared to state-of-art theoretical predictions such as **DIPHOX**, **RESBOS**, and **PYTHIA**. Overall agreement between data and theory, within known limitations, is observed.
- **Resummation** matched with NLO pQCD calculations works well at low  $p_T(\gamma\gamma)$  ( $<20 \text{ GeV}/c$ ) and large  $\Delta\phi_{\gamma\gamma}$  ( $>2.2$  rad).
- **Fragmentations** appear to be not under good control in sensitive kinematic regions [ $M(\gamma\gamma) < 60 \text{ GeV}/c^2$ ,  $20 \text{ GeV}/c < p_T(\gamma\gamma) < 50 \text{ GeV}/c$ ,  $\Delta\phi_{\gamma\gamma} < 1$  rad].
- Data-to-theory comparisons show best agreement for  $p_T(\gamma\gamma) < M(\gamma\gamma)$ , where theoretical uncertainties are smaller and predictions are less sensitive to the isolation requirement.
- **Parton-shower PYTHIA** Monte Carlo, which in previous analyses limited to matrix-element-based simulations was found to fail reproducing the data, now provides a description of the data competitive with full NLO calculations by **including ISR and FSR photons**
- A PRL ([arXiv:1106.5123](https://arxiv.org/abs/1106.5123)) and a PRD ([arXiv:1106.5131](https://arxiv.org/abs/1106.5131)) have been submitted


Backup slides

# Fragmentation contributions


- Collinear singularity in final state photon radiation off a parton can be handled e.g. via fragmentation functions.



## Single-photon fragmentation



## Double-photon fragmentation



Low-mass/small-angle diphoton pairs



Not included in any theoretical prediction!

- Fragmentation contributions can be suppressed via:
  - experimental photon isolation requirements (can only be approximated in theory)
  - $p_T(\gamma\gamma) < M(\gamma\gamma)$

$$E_T^{iso} = \sum_{\text{partons or hadrons within } R < 0.4} E_T - E_{T\gamma}$$





# Resummation of initial state gluons

- At fixed  $M(\gamma\gamma)$ , the differential cross section as a function of  $p_T(\gamma\gamma)$  at  $O(\alpha_s)$  given by:

$$\frac{d\sigma}{dp_{T\gamma\gamma}^2} = \sigma_0 \frac{\alpha_s}{\pi} \frac{1}{p_{T\gamma\gamma}^2} \left[ a_1 \ln\left(\frac{M_{\gamma\gamma}^2}{p_{T\gamma\gamma}^2}\right) + a_0 \right]$$

Fixed-order calculation less reliable for  $p_T(\gamma\gamma) \ll M(\gamma\gamma)$  and diverges as  $p_T(\gamma\gamma) \rightarrow 0$ .  
 [Also when  $\Delta\phi(\gamma,\gamma) \rightarrow \pi$ .]




# Resummation of initial state gluons


- At fixed  $M(\gamma\gamma)$ , the differential cross section as a function of  $p_T(\gamma\gamma)$  is given by:

$$\frac{d\sigma}{dp_{T\gamma\gamma}^2} = \sigma_0 \frac{\alpha_s}{\pi} \frac{1}{p_{T\gamma\gamma}^2} \left[ a_1 \ln\left(\frac{M_{\gamma\gamma}^2}{p_{T\gamma\gamma}^2}\right) + a_0 \right]$$

Fixed-order calculation less reliable for  $p_T(\gamma\gamma) \ll M(\gamma\gamma)$  and diverges as  $p_T(\gamma\gamma) \rightarrow 0$ .  
 [Also when  $\Delta\phi(\gamma,\gamma) \rightarrow \pi$ .]



Physical description of the  $p_T(\gamma\gamma)$  and  $\Delta\phi(\gamma,\gamma)$  distributions requires all-order resummation of soft and collinear logarithms.



# Resummation of initial state gluons


- At fixed  $M(\gamma\gamma)$ , the differential cross section as a function of  $p_T(\gamma\gamma)$  is given by:

$$\frac{d\sigma}{dp_{T\gamma\gamma}^2} = \sigma_0 \frac{\alpha_s}{\pi} \frac{1}{p_{T\gamma\gamma}^2} \left[ a_1 \ln\left(\frac{M_{\gamma\gamma}^2}{p_{T\gamma\gamma}^2}\right) + a_0 \right]$$


PRD 76, 013009 (2007)

Fixed-order calculation less reliable for  $p_T(\gamma\gamma) \ll M(\gamma\gamma)$  and diverges as  $p_T(\gamma\gamma) \rightarrow 0$ .


[Also when  $\Delta\phi(\gamma,\gamma) \rightarrow \pi$ .]



Only small effect on  $M(\gamma\gamma)$  from resummation



# Data set



x27 more luminosity than previous CDF publication!

# Triggers

## Diphoton-12

- **L1:**
  - EM  $E_T \geq 8$  GeV
  - $E_{\text{HAD}}/E_{\text{EM}} \leq 0.125$
  - $N_{\text{cluster}} = 2$
- **L2:**
  - EM  $E_T \geq 10$  GeV
  - $E_{\text{HAD}}/E_{\text{EM}} \leq 0.125$
  - $N_{\text{cluster}} = 2$
  - Isolation  $\leq 3$  GeV  
or IsoFraction  $\leq 0.15$
- **L3:**
  - EM  $E_T \geq 12$  GeV
  - $E_{\text{HAD}}/E_{\text{EM}} \leq 0.055 + 0.00045 \times E_{\text{tot}}/\text{GeV}$
  - $N_{\text{cluster}} = 2$
  - Isolation  $\leq 2$  GeV  
or IsoFraction  $\leq 0.1$
  - Shower profile:  $\chi^2_{\text{CES}} \leq 20$



“OR”

## Diphoton-18 Same as diphoton-12 except:

- **L2:**
  - EM  $E_T \geq 16$  GeV
  - No isolation
- **L3:**
  - EM  $E_T \geq 18$  GeV
  - No isolation

Trigger efficiency after offline event selection: 100% for  $E_T \geq 15$  GeV

# Photon characterization using track isolation



For a single  $\gamma$ , a weight can be defined to characterize it as signal or background:

- $\varepsilon = 1$  (0) if  $l_{\text{trk}} < (\geq) 1 \text{ GeV}/c$
- $\varepsilon_s =$  signal efficiency for  $l_{\text{trk}} < 1 \text{ GeV}/c$
- $\varepsilon_b =$  background efficiency for  $l_{\text{trk}} < 1 \text{ GeV}/c$


$$W = \frac{\varepsilon - \varepsilon_b}{\varepsilon_s - \varepsilon_b}$$

Both modeled by  $ae^{-E_T/b} + c$



Cut chosen at  $l_{\text{trk}} = 1 \text{ GeV}/c$ , where  $\varepsilon_s - \varepsilon_b = \text{max}$ , to optimize resolution

# Background estimation: 4x4 matrix method

- Relative uncertainties for photon and jet track ISO efficiencies estimated as a function of  $E_T$  using MC.
- Compared data and MC in complementary cones (same  $\theta$ ,  $\phi \pm \pi/2$  with true photon cones, assumed to collect same amount of underlying event):




→ Data and MC consistent to within 3%.



# Background estimation: 4x4 matrix method

- Relative uncertainties for photon and jet track ISO efficiencies estimated as a function of  $E_T$  using MC.
- Compared data and MC in complementary cones (same  $\theta$ ,  $\phi \pm \pi/2$  with true photon cones, assumed to collect same amount of underlying event):







→ Data and MC consistent to within 3%.







# Signal fraction

$$\text{Signal fraction} = \frac{N_{\gamma\gamma}}{N_{data}}$$



# Signal fraction

$$\text{Signal fraction} = \frac{N_{\gamma\gamma}}{N_{data}}$$



- Average ~40%
- Better at high mass:  
 60-80% for  $M(\gamma\gamma) \sim 80-150$  GeV/c<sup>2</sup>  
 ~80% for  $M(\gamma\gamma) > 150$  GeV/c<sup>2</sup>
- Better at high  $p_T(\gamma\gamma)$ :  
 ~70% for  $p_T(\gamma\gamma) > 100$  GeV/c

## Acceptance × efficiency


$$\frac{d\sigma}{dX} = \frac{N_{\gamma\gamma}}{\varepsilon \cdot A \cdot L \cdot \Delta}$$

- Defined as:

*Number of events with two reconstructed EM clusters passing all cuts*

*Number of events with two generator-level photons passing kinematic and isolation cuts*

- Estimated using detector- and trigger-simulated and reconstructed PYTHIA events reweighted to match the data



# Acceptance × efficiency


$$\frac{d\sigma}{dX} = \frac{N_{\gamma\gamma}}{\varepsilon \cdot A \cdot L \cdot \Delta}$$

- Defined as:

*Number of events with two reconstructed EM clusters passing all cuts*

*Number of events with two generator-level photons passing kinematic and isolation cuts*

- Estimated using detector- and trigger-simulated and reconstructed PYTHIA events reweighted to match the data
- RESBOS and DIPHOX do not include non-perturbative effects: underlying event and hadronization  
→ lower efficiency of the isolation cut relative to PYTHIA  
(PYTHIA events are removed from the isolated denominator of the efficiency due to the underlying event)
- Correction estimated by convoluting PYTHIA UE isolation energy with DIPHOX energy in the isolation cone  
→ constant per event factor of 0.88 applied to the data



# Acceptance × efficiency


$$\frac{d\sigma}{dX} = \frac{N_{\gamma\gamma}}{\varepsilon \cdot A \cdot L \cdot \Delta}$$

- Defined as:

*Number of events with two reconstructed EM clusters passing all cuts*

*Number of events with two generator-level photons passing kinematic and isolation cuts*

- Estimated using detector- and trigger-simulated and reconstructed PYTHIA events reweighted to match the data
- RESBOS and DIPHOX do not include non-perturbative effects: underlying event and hadronization
  - lower efficiency of the isolation cut relative to PYTHIA (PYTHIA events are removed from the isolated denominator of the efficiency due to the underlying event)
- Correction estimated by convoluting PYTHIA UE isolation energy with DIPHOX energy in the isolation cone
  - constant per event factor of 0.88 applied to the data



Uncertainties in the efficiency estimation:

- 3% from material uncertainty
- 1.5% from the EM energy scale
- 3% from trigger efficiency uncertainty
- 6% (3% per photon) from UE correction

Average efficiency ~40%  
 Total systematic uncertainty: ~7-15%  
 Comparable statistical uncertainty

## Corrections and tests

- EM energy scale set by tuning the reconstructed  $Z^0 \rightarrow e^+e^-$  mass to the world average by Gaussian fitting in the window  $M_{ee} = 86-96 \text{ GeV}/c^2$ 
  - correction applied as a function of time before event selection to account for a few events below the energy threshold which the correction pushes above threshold

## Corrections and tests

- EM energy scale set by tuning the reconstructed  $Z^0 \rightarrow e^+e^-$  mass to the world average by Gaussian fitting in the window  $M_{ee} = 86-96 \text{ GeV}/c^2$ 
  - correction applied as a function of time before event selection to account for a few events below the energy threshold which the correction pushes above threshold
- Measurement of the  $Z^0 \rightarrow e^+e^-$  cross section tests the cross section measurement procedures:
  - Trigger efficiency
  - Ability of MC to predict event selection efficiency
  - Efficiency corrections
  - Luminosity

“Photon-like”  $e^+e^-$  selection applied with special requirements:


  - Two tracks allowed in cluster
  - Leading  $p_T^{\text{trk}}$  cut applied on the 2<sup>nd</sup> track in cluster
  - Track isolation corrected subtracting leading  $p_T^{\text{trk}}$
  - $0.8 \leq E/p \leq 1.2$  cut applied to eliminate hard radiation

Measured/published ratio in the window  $M_{ee} = 65-115 \text{ GeV}/c^2$ :  $1.007 \pm 0.01$  with 5% RMS over time

## Corrections and tests

- EM energy scale set by tuning the reconstructed  $Z^0 \rightarrow e^+e^-$  mass to the world average by Gaussian fitting in the window  $M_{ee} = 86-96 \text{ GeV}/c^2$ 
    - correction applied as a function of time before event selection to account for a few events below the energy threshold which the correction pushes above threshold
  
  - Measurement of the  $Z^0 \rightarrow e^+e^-$  cross section tests the cross section measurement procedures:
    - Trigger efficiency
    - Ability of MC to predict event selection efficiency
    - Efficiency corrections
    - Luminosity
- “Photon-like”  $e^+e^-$  selection applied with special requirements:
- Two tracks allowed in cluster
  - Leading  $p_T^{\text{trk}}$  cut applied on the 2<sup>nd</sup> track in cluster
  - Track isolation corrected subtracting leading  $p_T^{\text{trk}}$
  - $0.8 \leq E/p \leq 1.2$  cut applied to eliminate hard radiation

Measured/published ratio in  $M_{ee} = 65-115 \text{ GeV}/c^2$  window:  $1.007 \pm 0.01$  with 5% RMS over time






- Experimental effects (photon energy resolution, misvertexing) lead to event migration

$$\text{Purity (bin } i) = N(\text{gen bin } i \text{ AND reco bin } i) / N(\text{reco bin } i)$$



→ The acceptance correction also accounts for this




# Matrix element and radiation contributions in Pythia




# Data-to-theory cross section ratios



# Data-to-theory cross section ratios





**Resummation**




**Fragmentations**


# Data-to-theory cross section ratios



# Data-to-theory cross section ratios





Resummation





Fragmentations




# Data-to-theory cross section ratios for $p_T(\gamma\gamma) < M(\gamma\gamma)$



# Data-to-theory cross section ratios for $p_T(\gamma\gamma) < M(\gamma\gamma)$




# Differential cross sections for $p_T(\gamma\gamma) > M(\gamma\gamma)$




- Theory underestimates the data at the peak  $M_{\gamma\gamma} \sim 30$  GeV/c<sup>2</sup>
- Theory underestimates the data for  $p_T(\gamma\gamma) < 90$  GeV/c
- Theory underestimates the data for  $\Delta\phi_{\gamma\gamma} < 1.7$  rad




# Differential cross sections for $p_T(\gamma\gamma) > M(\gamma\gamma)$



- Theory underestimates the data



- Theory underestimates the data



- Theory underestimates the data