



Contribution ID: 360

Type: **Parallel contribution**

## Searches for Diboson Production in the Lepton + MET + Jets Final State in ATLAS

*Wednesday, 10 August 2011 18:15 (19 minutes)*

The study of diboson production at high energy colliders tests the electroweak sector of the standard model (SM) and provides a sensitive probe of new physics beyond the SM. An important example is the production of a Higgs boson with mass greater than  $140 \text{ GeV}/c^2$  which decays primarily to W boson pairs. The diboson decay channel where one W boson decays to leptons and the other vector boson decays to quarks leading to high energy jets is particularly interesting due to its large branching fraction as compared to all-leptonic channels but is also challenging due to large backgrounds, particularly from W+jets.

We present searches for diboson production in the lepton + MET+ jets final state using  $\sqrt{s}=7 \text{ TeV}$  collision data collected by the ATLAS detector during the 2011 run. Particular emphasis is placed on searches for (1) the SM Higgs boson with mass above the W pair production threshold and (2) SM WW+WZ production.

**Primary author:** NEUBAUER, Mark (University of Illinois at Urbana-Champaign)

**Presenter:** NEUBAUER, Mark (University of Illinois at Urbana-Champaign)

**Session Classification:** Higgs Physics

**Track Classification:** Higgs Physics