



Contribution ID: 385

Type: **Parallel contribution**

Search for the Standard Model Higgs boson in $H \rightarrow ZZ$ decay channels with the ATLAS detector

Tuesday, 9 August 2011 14:00 (20 minutes)

The SM Higgs boson in the medium and high mass ranges has a large branching ratio for decays to a pair of neutral weak bosons. Three decay modes of the Z boson pair have been explored by ATLAS. One Z is typically produced on-shell, which can be tagged using leptonic decay products. The decay of the second Z^* leads to three independent search channels: $llqq$, $ll\nu\nu$ and $llll$. Background compositions and topologies differ among these channels. The $llqq$ search can use jet information to reduce top and Z +jets backgrounds; the $ll\nu\nu$ search requires a good understanding of missing transverse energy, while the 4 lepton ('golden') channel is almost background-free and, owing to its low production rate, needs excellent lepton efficiencies in order to be sensitive.

This talk summarizes results in all three $H \rightarrow ZZ$ channels using data collected in 2011.

Primary author: CARRILLO MONTOYA, G (Department of Physics-University of Wisconsin)

Presenter: CARRILLO MONTOYA, G (Department of Physics-University of Wisconsin)

Session Classification: Higgs Physics

Track Classification: Higgs Physics