

# SHADOWS

## Search for Hidden And Dark Objects With the SPS

<sup>(1)</sup> *CERN, European Organization for Nuclear Research, CH-1211 Geneva 23, Switzerland*

<sup>(2)</sup> *INFN, Sezione di Napoli, Napoli, Italy*

<sup>(3)</sup> *Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany*

<sup>(4)</sup> *INFN, Sezione di Ferrara, Ferrara, Italy*

<sup>(5)</sup> *INFN Laboratori Nazionali di Frascati, Frascati (Rome), Italy*

<sup>(6)</sup> *INFN, Sezione di Roma III, Roma, Italy*

<sup>(7)</sup> *Johannes Gutenberg Universität Mainz, Mainz, Germany*

<sup>(8)</sup> *INFN, Sezione di Bologna, Bologna, Italy*

<sup>(9)</sup> *Karlsruhe Institute of Technology (KIT), Karlsruhe, Germany*

<sup>(10)</sup> *PARTREC and University of Groningen, Groningen, The Netherland*

<sup>(11)</sup> *INFN, Sezione di Roma III, Roma, Italy*

<sup>(12)</sup> *University of Freiburg, Freiburg, Germany*

<sup>(13)</sup> *Charles University, Prague, Czech Republic*

<sup>(14)</sup> *Royal Holloway, University of London, UK*

<sup>(15)</sup> *INFN, Sezione di Roma1, Roma, Italy*

<sup>(16)</sup> *University of Bologna, Bologna, Italy*

**+ the invaluable support of the PBC-CBWG, PBC-ECN3 Task Force, NA-CONS team, and CERN EP-DT Group.**

150<sup>th</sup> SPSC meeting – Open Session - 5 September 2023

# European Strategy for Particle Physics recommendations

[https://cds.cern.ch/record/2721370/files/CERN-ESU-015\\_2020%20Update%20European%20Strategy.pdf](https://cds.cern.ch/record/2721370/files/CERN-ESU-015_2020%20Update%20European%20Strategy.pdf)



#### 4. Other essential scientific activities for particle physics:

- a) *The quest for dark matter and the exploration of flavour and fundamental symmetries are crucial components of the search for new physics.*
- *This search can be done in many ways, for example through precision measurements of flavour physics and electric or magnetic dipole moments, and searches for axions, dark sector candidates and feebly interacting particles.*
- *There are many options to address such physics topics including energy-frontier colliders, accelerator and non-accelerator experiments. A diverse programme that is complementary to the energy frontier is an essential part of the European particle physics Strategy.*

SHADOWS and HIKE can explore simultaneously the multi-TeV region via precision measurements and low-mass NP with very feeble couplings becoming main players in the future CERN diversity programme.

# SHADOWS: 20 very intense (and exciting) months..

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH (CERN)

**Expression of Interest, 6 January 2022**

SHADOWS

Search for Hidden And Dark Objects With the SPS

*Expression of Interest*

W. Baldini<sup>(1)</sup>, A. Balla<sup>(2)</sup>, J. Bernhard<sup>(3)</sup>, A. Calcaterra<sup>(2)</sup>, V. Cafaro<sup>(4)</sup>, N. Charitonidis<sup>(3)</sup>, A. Ceccucci<sup>(3)</sup>, V. Cicero<sup>(4)</sup>, P. Ciambrone<sup>(2)</sup>, H. Danielsson<sup>(3)</sup>, A. De Roeck<sup>(3)</sup>, F. Duval<sup>(3)</sup>, G. D'Alessandro<sup>(3)</sup>, G. Felici<sup>(2)</sup>, L. Foggetta<sup>(2)</sup>, L. Gatignon<sup>(5)</sup>, A. Gerbershagen<sup>(3)</sup>, V. Giordano<sup>(4)</sup>, G. Lanfranchi<sup>(2)</sup>, I. Lax<sup>(4)</sup>, A. Montanari<sup>(4)</sup>, R. Murphy<sup>(3)</sup>, T. Napolitano<sup>(2)</sup>, A. Paoloni<sup>(2)</sup>, G. Papalino<sup>(2)</sup>, T. Rovelli<sup>(4)</sup>, A. Saputi<sup>(2)</sup>, S. Schuchmann<sup>(6)</sup>, F. Stummer<sup>(7)</sup>, G. Torromeo<sup>(4)</sup>, N. Tosi<sup>(4)</sup>, A. Vannozzi<sup>(2)</sup>.

(1) INFN, Sezione di Ferrara, Ferrara, Italy  
 (2) INFN, Laboratori Nazionali di Frascati, Frascati (Rome), Italy,  
 (3) CERN  
 (4) INFN, Sezione di Bologna, Bologna, Italy  
 (5) University of Lancaster, Lancaster, UK  
 (6) University of Mainz, Germany  
 (7) Royal Holloway, University of London, UK

13.01.2022 CERN-SPSC-2022-030 / 262C-EOI-033

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH (CERN)

**Letter of Intent, 4 November 2022**

SHADOWS

Search for Hidden And Dark Objects With the SPS

*Letter of Intent*

M. Alviggi<sup>(1)</sup>, S. Bachmann<sup>(2)</sup>, W. Baldini<sup>(3)</sup>, A. Balla<sup>(4)</sup>, M. Biglietti<sup>(8)</sup>, V. Büscher<sup>(11)</sup>, A. Calcaterra<sup>(4)</sup>, V. Cafaro<sup>(5)</sup>, N. Charitonidis<sup>(6)</sup>, A. Ceccucci<sup>(6)</sup>, V. Cicero<sup>(5)</sup>, P. Ciambrone<sup>(4)</sup>, H. Danielsson<sup>(6)</sup>, M. Dellapietra<sup>(1)</sup>, A. De Roeck<sup>(6)</sup>, F. Duval<sup>(6)</sup>, G. Felici<sup>(4)</sup>, T. Ferber<sup>(7)</sup>, L. Foggetta<sup>(4)</sup>, M. Gatta<sup>(4)</sup>, A. Gerbershagen<sup>(13)</sup>, V. Giordano<sup>(5)</sup>, S. Hansmann-Menzemer<sup>(2)</sup>, P. Iengo<sup>(1)</sup>, M. Iodice<sup>(8)</sup>, K. Jakobs<sup>(9)</sup>, M. Klute<sup>(7)</sup>, K. Köneke<sup>(9)</sup>, M. Koval<sup>(10)</sup>, G. Lanfranchi<sup>(4)</sup>, A. Laudrain<sup>(11)</sup>, I. Lax<sup>(5)</sup>, B. Leverington<sup>(2)</sup>, P. Lichard<sup>(6)</sup>, K. Massri<sup>(6)</sup>, A. Montanari<sup>(5)</sup>, R. Murphy<sup>(6,12)</sup>, T. Napolitano<sup>(4)</sup>, F. Neuhaus<sup>(11)</sup>, L. J. Nevay<sup>(6)</sup>, A. Paoloni<sup>(4)</sup>, G. Papalino<sup>(4)</sup>, U. Parzefall<sup>(9)</sup>, S. Ritter<sup>(11)</sup>, T. Rovelli<sup>(5,14)</sup>, A. Saputi<sup>(3)</sup>, B. Schmidt<sup>(6)</sup>, M. Schott<sup>(11)</sup>, H.C. Schultz-Coulon<sup>(2)</sup>, G. Sekhniaidze<sup>(1)</sup>, F. Stummer<sup>(6,12)</sup>, G. Torromeo<sup>(5)</sup>, N. Tosi<sup>(5)</sup>, U. Uwer<sup>(2)</sup>, M. van Dijk<sup>(6)</sup>, A. Vannozzi<sup>(4)</sup>, R. Wanke<sup>(11)</sup>, C. Weiser<sup>(9)</sup>, P. Wertelaers<sup>(6)</sup>, T. Zickler<sup>(6)</sup>

(1) INFN, Sezione di Napoli, Napoli, Italy  
 (2) Heidelberg University, Heidelberg, Germany  
 (3) INFN, Sezione di Ferrara, Ferrara, Italy  
 (4) INFN, Laboratori Nazionali di Frascati, Frascati (Rome), Italy,  
 (5) INFN, Sezione di Bologna, Bologna, Italy  
 (6) CERN  
 (7) Karlsruhe Institute of Technology, KIT, Germany  
 (8) INFN, Sezione di Roma III, INFN, Italy  
 (9) University of Freiburg, Freiburg, Germany  
 (10) Charles University, Prague, Czech Republic  
 (11) University of Mainz, Germany  
 (12) Royal Holloway, University of London, UK  
 (13) PARTREC and University of Groningen, Groningen, The Netherland  
 (14) University of Bologna, Bologna, Italy

CERN-SPSC-2022-030 / SPSC-I-254  
04/11/2022

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH (CERN)

**Technical Proposal, 18 August 2023**

SHADOWS


Search for Hidden And Dark Objects With the SPS

*Technical Proposal*

M. Alviggi<sup>(1)</sup>, S. Bachmann<sup>(2)</sup>, W. Baldini<sup>(3)</sup>, A. Balla<sup>(4)</sup>, M. Barth<sup>(2)</sup>, M. Biglietti<sup>(5)</sup>, V. Büscher<sup>(6)</sup>, A. Calcaterra<sup>(4)</sup>, V. Cafaro<sup>(7)</sup>, M. Cavallina<sup>(3)</sup>, A. Ceccucci<sup>(8)</sup>, D. Chouhan<sup>(6)</sup>, V. Cicero<sup>(7)</sup>, P. Ciambrone<sup>(4)</sup>, H. Danielsson<sup>(8)</sup>, M. Della Pietra<sup>(1)</sup>, C. Delogu<sup>(6)</sup>, A. De Roeck<sup>(8)</sup>, L. Dittmann<sup>(2)</sup>, F. Duval<sup>(8)</sup>, G. Felici<sup>(4)</sup>, T. Ferber<sup>(9)</sup>, L. Foggetta<sup>(4)</sup>, E. Gamberini<sup>(8)</sup>, M. Gatta<sup>(4)</sup>, A. Gerbershagen<sup>(10)</sup>, V. Giordano<sup>(7)</sup>, S. Hansmann-Menzemer<sup>(2)</sup>, P. Iengo<sup>(11)</sup>, M. Iodice<sup>(11)</sup>, K. Jakobs<sup>(12)</sup>, J. Kieseler<sup>(9)</sup>, M. Klute<sup>(9)</sup>, K. Köneke<sup>(12)</sup>, M. Koval<sup>(13)</sup>, G. Lanfranchi<sup>(4)</sup>, A. Laudrain<sup>(6)</sup>, I. Lax<sup>(87)</sup>, T. M. Leeftang<sup>(2)</sup>, G. Lehmann Miotto<sup>(8)</sup>, B. Leverington<sup>(2)</sup>, P. Lichard<sup>(8)</sup>, K. Massri<sup>(8)</sup>, A. Montanari<sup>(7)</sup>, T. Napolitano<sup>(4)</sup>, A. Paoloni<sup>(4)</sup>, G. Papalino<sup>(4)</sup>, U. Parzefall<sup>(12)</sup>, B. Ponzio<sup>(4)</sup>, B. Regnery<sup>(9)</sup>, S. Ritter<sup>(6)</sup>, S. Rosati<sup>(15)</sup>, T. Rovelli<sup>(16,7)</sup>, S. Roy<sup>(2)</sup>, A. Saputi<sup>(3)</sup>, B. Schmidt<sup>(8)</sup>, M. Schott<sup>(6)</sup>, D. Schub<sup>(2)</sup>, H.C. Schultz-Coulon<sup>(2)</sup>, G. Sekhniaidze<sup>(1)</sup>, R. Stamen<sup>(2)</sup>, G. Torromeo<sup>(7)</sup>, N. Tosi<sup>(7)</sup>, N. Trevisani<sup>(6)</sup>, U. Uwer<sup>(2)</sup>, A. Vannozzi<sup>(4)</sup>, C. Wang<sup>(6)</sup>, R. Wanke<sup>(6)</sup>, C. Weiser<sup>(12)</sup>, C. Welschoff<sup>(2)</sup>, P. Wertelaers<sup>(8)</sup>


From the Expression of Interest (Jan 22) → Proposal (Aug. 23)  
 the collaboration almost tripled (82 collaborators, 16 institutions)

# From the first idea (Jan 2022)....



**Initial idea of SHADOWS location in TCC8**

.. to a preliminary but complete project...



**Render of SHADOWS in the TCC8 tunnel**


## ... documented in 212 pages of the Technical Proposal

### Contents

<b>1 Introduction</b>	<b>1</b>	5.4.6 Evolution toward the TDR	58	<b>7 Integration</b>	<b>113</b>	<b>10 FIP physics reach</b>	<b>183</b>
<b>2 Beam line and target area</b>	<b>5</b>	5.5 Dipole Magnet	60	7.1 Accessibility	113	10.1 Theoretical framework	183
2.1 Beam line description	6	5.5.1 Warm option	61	7.2 Civil Engineering	115	10.2 Worldwide context	185
2.2 Current intensity and known limitations	8	5.5.2 Superconducting option	69	7.3 Transport and handling	115	10.3 SHADOWS physics reach	188
2.3 Future intensity	8	5.5.3 Preliminary cost estimate for the warm option	74	7.4 Services	116	10.3.1 Computation of the physics reach	188
2.4 The new target complex	10	5.6 Tracker	75	<b>8 Monte Carlo simulation</b>	<b>118</b>	Light dark scalar mixing with the Higgs	189
<b>3 The Magnetized Iron Block (MIB) muon sweeping system</b>	<b>12</b>	5.6.1 Physics requirements	75	8.1 Monte Carlo framework	118	Sensitivity to ALPs with fermion couplings	190
3.1 General idea	13	5.6.2 Straw drift-tube technology	77	8.2 Beamline simulation	118	Sensitivity to Heavy Neutral Leptons	191
3.2 Updates in the Muon Sweeping System with respect to the LoI	14	5.6.3 Straw tubes	77	8.3 Detector simulation	119	10.3.2 Results	195
Stage 1 - Size reduction:	15	5.6.4 Detector concept and optimisation	78	8.4 Reconstruction	124	10.3.3 Impact of SHADOWS on the FIP physics programme	201
Stage 2 - Moving towards an integration friendly design:	15	5.6.5 Readout electronics and data-flow	79	8.4.1 Tracking reconstruction	124	<b>11 Neutrino physics reach</b>	<b>203</b>
Stage 3 - Replacing the detector cover with a MIB:	16	5.6.6 Required R&D	81	Reconstruction methods	124	11.1 Background and Neutrino Fluxes	203
General updates:	16	5.6.7 Preliminary cost estimate	83	Reconstruction performance	124	11.2 Muon Neutrino Physics: Cross Sections, Deep Inelastic Scattering and Charm Production	205
3.3 MIB system working principle	18	5.7 Timing Detector	84	8.4.2 ECAL reconstruction	129	11.3 Tau Neutrino Physics: Discovery of Anti-Tau Neutrinos, Cross Sections, Structure Functions, Anomalous Magnetic Moment	207
3.4 MIB muon background reduction	20	5.7.1 Physics requirements	84	Reconstruction methods	129	11.4 Tests of Lepton Universality and Searches for new physics	207
3.5 MIB design optimization process	21	5.7.2 Technology description	84	Results for benchmark scenarios	130	<b>12 Project Organization</b>	<b>208</b>
3.6 MIB finite element layout	24	5.7.3 Detector concept	84	<b>8.5 Signal studies</b>	<b>134</b>	12.1 Schedule for R&D, construction, installation, commissioning and operation	208
3.7 MIB preliminary cost estimate	26	5.7.4 Required R&D	85	8.5.1 Signal samples	134	12.2 Preliminary detector cost estimate and group interests	208
<b>4 Radiation levels in the SHADOWS area</b>	<b>27</b>	5.7.5 Preliminary cost estimate	85	8.5.2 Signal reconstruction and selection	136	12.3 Present Status, Required R&D	209
4.1 Presentation of the FLUKA geometry	27	5.8 Electromagnetic calorimeter	87	<b>8.6 Background studies</b>	<b>138</b>	<b>13 Conclusions and Outlook</b>	<b>212</b>
4.2 Radiation to electronics considerations	28	5.8.1 Physics and Experimental Requirements	87	8.6.1 Background samples	138	<b>14 Acknowledgements</b>	<b>212</b>
4.3 Radiation Protection	31	5.8.2 Layout Options	87	8.6.2 Muon background	139		
4.3.1 Prompt radiation	32	5.8.3 ECAL Baseline Design	89	The beamline and its magnetic elements	139		
4.3.2 Residual radiation	33	5.8.4 ECAL Performance	90	Muon flux without the MIB	141		
4.3.3 Soil activation	34	5.8.5 Required R&D	90	Muon flux with the MIB	141		
4.3.4 Air activation	36	5.8.6 Preliminary cost estimate	91	Combinatorial muon background	148		
<b>5 The SHADOWS detector</b>	<b>37</b>	5.9 Muon system	93	Inelastic muon interactions	153		
5.1 Detector concept	37	5.9.1 Physics requirements	93	Neutrino background	158		
5.2 Physics requirements	41	5.9.2 Detector layout	93	8.6.4 Summary of the background components	163		
5.3 Decay vessel	42	5.9.3 Technology choice	94	<b>8.7 Validation of the muon background simulation with data</b>	<b>164</b>		
5.3.1 Overview and dimensions	42	5.9.4 Required R&D	98	8.7.1 Summary of the 2021 measurement of the "on-axis" muon flux	164		
5.3.2 Flanges and O-ring compression	46	5.9.5 Preliminary cost estimate	99	8.7.2 Measurement of the 2023 "off-axis" muon flux	167		
5.4 Upstream and Lateral Vetoes	50	5.10 Trigger and Data Acquisition System	101	The setup	167		
5.4.1 Physics requirements	50	5.10.1 Design principles	101	Beam line settings	170		
5.4.2 Resistive Pad Micromegas	51	5.10.2 Preliminary cost estimate	104	Results from simulation	172		
5.4.3 Detector concept	53	<b>6 The Neutrino Subdetector System: NaNu</b>	<b>105</b>	Results from data	176		
5.4.4 Required R&D	55	6.1 Detector Concept	105	<b>9 Evolution towards the TDR</b>	<b>180</b>		
5.4.5 Preliminary cost estimate	57	6.2 Expected Detector Performance	107	9.1 Decay volume filled with Helium	180		
		6.3 Preliminary Detector R&D	109	9.2 Optimisation of the dipole magnet dimensions	180		
		6.4 Preliminary cost estimate	110	9.3 Optimisation of the MIB system	181		
				9.4 Background reduction for Super-NaNu	182		

Submitted to the SPSC the 18 August 2023.


# SHADOWS step-by-step: The beam



Intensity to TCC8 [p/spill]	(PoT/year)	Spill Length [s]	Fastest Repetition Period [s]
up to $2.0 \times 10^{13}$	up to $1.2 \times 10^{19}$	$\geq 4.5$	14.4

Table 1. Experimental requirements for HIKE/SHADOWS.

Up to a factor 7 more than the current intensity




**$5 \times 10^{19}$  protons-on-target (pot), with 4.8 sec long spills, can be delivered to ECN3 in 4 integrated years with a dedicated beam delivery for ECN3, and shared cycles to EHN1 and EHN2. This annual yield is fully compatible with the current North Area operation.**



# The high-intensity K12 beam line

Fully compatible and synergistic with the (already funded) North-Area Consolidation Programme:




## NA-CONS programme:


**Phase 1: 2022–2028 (up to end LS3),**  
primary beam areas TT20, TDC2, TCC2 and initial section  
of NA Transfer Tunnels.

**Phase 2: 2026–2034 (up to end LS4),**  
completing the consolidation of the secondary beam areas.

# SHADOWS step-by-step: The Target station




The instantaneous and integrated beam intensities requested by HIKE/SHADOWS require the installation of a new target complex, associated cooling and ventilation systems, and shielding in TCC8.



- **Significant shielding improvement with respect to the current NA62 target system:** optimised to reduce the prompt radiation above ground to comply with a Non-designated Area.
- **A new TAX Cu-Fe based system needed:** along with upgraded cooling and maintenance capabilities.
- **Change from Kaon mode to beam dump mode very fast:** mostly dominated by different magnet settings

1-MeV equivalent neutron fluence, high energy hadron fluence and thermal neutron fluence evaluated with a detailed FLUKA simulation in SHADOWS area





Quantity	Detector location	R2E safe
Total Ionising Dose (Gy)	0.1-1 per year	< 10
1-MeV neutron equivalent fluence ( $\text{cm}^{-2}$ )	$10^9$ - $10^{10}$ per year	< $10^{11}$
High-Energy Hadron eq. fluence ( $\text{cm}^{-2}\text{year}^{-1}$ )	few $10^9$	< $3 \times 10^6$
Th. neutron eq. fluence ( $\text{cm}^{-2}\text{year}^{-1}$ )	< $5 \times 10^{10}$	< $3 \times 10^7$

**Radiation levels are not a show-stopper in the very-close-to-dump SHADOWS location**

Radiation- tolerant electronics will have to be used in proximity of the SHADOWS detector  
Dedicated alchoves with iron/concrete shielding far from the dump for the off-detector electronics.

# SHADOWS step-by-step: The MIB system

MIB system necessary to sweep the beam-induced muon flux out of the SHADOWS acceptance.



A fully reoptimized design with decreased cost and increased performance wrt to the LoI  
Complete finite element models implemented to study performance






Table 2. New MIB design and dimensions compared with the ones used in the LoI.

		Stage 1	Stage 2	Stage 3
<b>Proposal</b>	size (xyz)	active (1.1×1.6×3.5) m	3×(0.4×2.4×3.0) m	(1.3×1.1×2.0) m
	passive	–	3×(0.4×2.4×1.4) m	–
current		100 A	100 A	100 A
<b>LoI</b>	size (xyz)	active (3.0×4.0×3.5) m	(0.7×4.0×15.2) m	–
	passive	–	–	(1.5×3.5×5.0) m
	current	250 A	250 A	0 A

## The MIB muon sweeping system: Performance


Muon Illumination in front of the first tracking chamber



**Muon flux reduction in SHADOWS acceptance from 150 MHz  $\rightarrow$  2 MHz**


	$\mu^+ + \mu^-$	$\mu^+$	$\mu^-$
rate without MIB	147 MHz	81 MHz	66 MHz
MIB reduction factor	$\sim 70$	$\sim 58$	$\sim 94$
rate with MIB	2.1 MHz	1.4 MHz	0.7 MHz

# SHADOWS step-by-step: The Detector





# The Detector




**SHADOWS must be able to reconstruct and identify most of the visible final states of FIPs decays**

Scalar portal	$l^+l^-, \pi^+\pi^-, K^+K^-$
Pseudo-scalar portal	$l^+l^-, \gamma\gamma, \pi^+\pi^-, K^+K^-$
Vector portal	$l^+l^-, \pi^+\pi^-, K^+K^-$
Fermion (neutrino) portal	$l^\pm\pi^\mp, l^\pm K^\mp, l^\pm\rho^\mp (\rho^\mp \rightarrow \pi^\pm\pi^0), l^+l^-\nu$

Standard spectrometer, with 19m long in-vacuum decay volume, excellent tracking system, high resolution timing layer, ECAL with pointing capability, muon system and efficient vetoes


# The Detector



## Important Remark:

- SHADOWS detectors are based on **well known and established technologies**.
- The detector readiness leverages on the **long-standing expertise of the groups involved**.
- Most of the groups have already **built and operated prototypes or even full-size detectors, mostly at the LHC**.
- **All these elements guarantee the readiness of the detector for data taking in 2030.**


## The Detector: the decay vessel



Fully engineered, modular, transportable, stainless-steel based, in-vacuum decay vessel anchored to the dipole magnet and containing the 4 tracking stations.

# The Detector: Upstream & Lateral Veto

Institutes:  
INFN-Roma3, INFN-Naples  
Expertise:  
ATLAS new small wheels.



## Goal:


veto muons that enter the decay vessel escaping the MIB system

## Technology:


### Double layer of micromegas detectors:

- efficiency  $> 99.8\%$
- space resolution:  $o(1)$  mm
- time resolution:  $o(10)$  ns
- rate capability: up to 10 MHz /cm<sup>2</sup>

Requirements fully satisfied.



Upstream  
Veto



First small prototype funded by INFN in 2023  
A large size prototype will be prepared in 2024

# The Detector: Dipole Magnet

Design by Piet Wertelaers  
CERN EP-DT group




## Requirements:

- field integral  $\sim 1$  Tm (similar to NA62 dipole magnet MNP33)
- low power consumption
- 2.7x2.7 m aperture

## Two solutions:



- *warm option (baseline)*:
  - dissipated power: 287 kW, 10x less than MNP33 NA62 dipole
  - copper-based coil, iron-based yoke
- *superconducting option*
  - compelling & innovative, will be studied for the TDR.

## Dipole magnet: warm option




# The Detector: Dipole Magnet


Design by Piet Wertelaers  
CERN EP-DT group





Field integrals (in Tm) in y



Field integrals (in Tm) in x




Detailed finite-element simulation and corresponding field integrals (in y, and x)



Requirements fully satisfied.

# The Detector: Tracking Stations

Institutes: University of Heidelberg, CERN  
 Expertise: LHCb Outer Tracker,  
 LHCb SciFi Tracker, NA62 straw tracker



Tracking

## Main goal:

Reconstruct signals & reject background  
 with at least 2 tracks

## Requirements:


Vertex resolution  $\sigma(1)$  cm over 20 m  
 IP resolution  $\sigma(1)$  cm from 35 m distance  
 Mass resolution: 1-2% mass

## Baseline technology:

Straw Tubes in vacuum (NA62-like)

(Scintillating fibres technology under consideration)

Four stations, 2 views each, 4 layers per view




## Technology Requirements

Single hit resolution	$< 150 \mu\text{m}$
Single hit efficiency	$> 98\%$
Material per stereo-layer	$< 0.1\% X_0$
Rate capability (hot spot)	$200 \text{ Hz/cm}^2$
Rate capability (total)	4 MHz




# The Detector: Tracking Performance


Institutes: University of Heidelberg, CERN  
 Expertise: LHCb Outer Tracker,  
 LHCb SciFi Tracker, NA62 straw tracker




### Mass resolution



### Vertex resolution



### Impact parameter resolution




Requirements fully satisfied.



# The Detector: Timing layer

Institutes: University of Freiburg  
Expertise: fast timing silicon-based detectors for ATLAS ITk.



## Goals:


reject muon combinatorial background requiring fast time coincidence

## Requirements:

Time resolution of  $\mathcal{O}(100)$  ps

## Baseline solution:


plastic scintillating bars with direct sipm readout  
about 1 cm thickness, 6 cm width, 1.26 m length,  
thereby covering half of the  $2.5 \times 2.5$  m<sup>2</sup> acceptance.  
Proved to reach  $< 100$  ps time resolution.



Requirements fully satisfied.

# The Detector: ECAL

**Groups:** Mainz cluster of excellence, Karlsruhe Institute of Technology; ASIC developed by Heidelberg.  
**Expertise:** NA62 hadron calorimeter, CMS ECAL.



## Requirements:

Moderate energy resolution:

10-15% /  $\sqrt{E(\text{GeV})}$

Particle ID via E/p measurement

Pointing capability for fully neutral

final state (eg: ALP  $\rightarrow$  gg)

Time resolution :  $\mathcal{O}(1)$  ns.

## Baseline solution:


StripCAL: 2.5 m long, 1cm wide, 1 cm thick

strips in x,y directions read out with WLS fibres+sipts

Alternating with iron layers, 9 mm thick.



20  $X_0$  total depth to avoid shower leakage

*Render of the GEANT4 geometry of the SHADOWS ECAL.*



# The Detector: ECAL - performance

**Groups:** Mainz cluster of excellence, Karlsruhe Institute of Technology; ASIC developed by Heidelberg.  
**Expertise:** NA62 hadron calorimeter, CMS ECAL.



Requirements fully satisfied.

# The Detector: Muon System

**Groups:** INFN-LNF, INFN-Bologna, INFN-Ferrara  
**Expertise:** LHCb muon system, CMS muon system



**Goal:**  
 identify muons and reduce muon combinatorial background via timing measurement.

**Technology:**  
 3 stations of [scintillating tiles with direct sipm readout](#)  
 Interleaved by iron filters. Measured 250 ps resolution per station.


**Two full-size modules already funded by INFN in 2023 and used to measure the off-axis muon flux in ECN3 during the June 2023 campaign (see later).**

[Requirements fully satisfied.](#)



# The Detector: Integration

Detector integrated in the area, including civil engineering, transport and handling, and services




# The Detector: TDAQ system



**Trigger-less approach:** data filtering mostly at high-level-trigger level. stream all (zero suppressed) raw data directly from the on-detector electronics towards the TDAQ.

**Use IpGBT** (Low Power Gigabit Transceiver) technology developed for the HL-LHC experiments, which offers data Rates of 8.96 Gb/s.


**Joint project with HIKE** to maximise expertise sharing & development effort.



# The Detector: Full simulation

**SHADOWS full Monte Carlo simulation is part of the general NA62 Monte Carlo framework and is a C++, GEANT4-based code.**


- The beamline simulation is done using the Geant4-based BDSIM package.
- The signals are generated with PYTHIA 8.32
- The background samples with the GEANT4-based BDSIM package.
- The inelastic interactions of neutrinos and muons with the detector material are simulated using GENIE and Pythia6 generators.



## **Advantages of a common HIKE+SHADOWS framework**

- *centralised production of signal and MC samples*
- *possibility of studying the mutual interference*
- *possibility of easily combining physics results*


## Background: The name of the game



**Main background arise from muons and neutrinos emerging from the dump.  
An off-axis setup is much less affected by background than an on-axis one,  
as muons and neutrinos are mostly emitted in the forward direction.**



# Background: The name of the game





## Three important backgrounds to be considered:

1. Muon combinatorial background
2. Muon inelastic interactions with the decay vessel
3. Neutrino inelastic interactions with the decay vessel & residual air in the decay volume

For the first two backgrounds the knowledge of the muon flux is paramount.

## Validation of the simulated *on-axis* muon flux with NA62 data



Monte Carlo simulation has been compared against data collected by NA62 in October 2021, when the experiment was successfully operated in beam-dump mode for about 1 week at about 150% the nominal NA62 beam intensity. In this period NA62 collected about  $1.5 \times 10^{17}$  pot



Excellent agreement in shape, the MC rate is about 3 times less than data as expected.  
MC rates corrected by this factor.




## Validation of the simulated *off-axis* muon flux with SHADOWS prototypes

Measurement performed in June 2023 with NA62 operated in beam-dump mode at nominal beam intensity.  
Effort partially funded via EUROLABS European Grant.



## Validation of the simulated *off-axis* muon flux with SHADOWS prototypes

Measurement performed in June 2023 with NA62 operated in beam-dump mode at nominal beam intensity.  
Effort partially funded via EUROLABS European Grant.





# The SHADOWS teams at work in TCC8



A big thanks to the NA62 Collaboration for the support and help

# Validation of the simulated *off-axis* muon flux with SHADOWS prototypes



**Results from data:** 250-300 counts/ $\text{cm}^2/10^{12}$  pot



**Results from simulation (rescaled):**  $260 \pm 20$  counts/ $\text{cm}^2/10^{12}$  pot

Excellent agreement between the results obtained with (very different) detectors gives reliability of the measurement.

**Off-axis measurements confirmed the on-axis ones. Simulation is now fully validated.**

# 1. Combinatorial background: MIB reduction


Muon Illumination in front of the first tracking chamber



## Muon flux reduction from 150 MHz $\rightarrow$ 2 MHz

	$\mu^+ + \mu^-$	$\mu^+$	$\mu^-$
rate without MIB	147 MHz	81 MHz	66 MHz
MIB reduction factor	$\sim 70$	$\sim 58$	$\sim 94$
rate with MIB	2.1 MHz	1.4 MHz	0.7 MHz

# 1. Combinatorial background: Detector reduction




## Combinatorial background reduction as a function of the selection

	$N_{\mu\mu}/\text{spill}$	requirement
	3000	timing (T)
	$1.2 \cdot 10^{-2}$	Veto (UV)
	$6.0 \cdot 10^{-6}$	vertex requirements (CDA)
	$6.0 \cdot 10^{-10}/3.0 \cdot 10^{-7}$ (fully/partially rec. events)	IP requirements (IP)
	$N_{\mu\mu}/5 \cdot 10^{19}$ pot	
	0.001/0.7 events (fully/partially rec.)	T & UV & CDA & IP

The applied selection has an efficiency of 83% on signals.



## 2. Muon inelastic interactions: Detector reduction




**Muon inelastic interaction reduction as a function of the selection**

$N_{\text{muon inel. inter.}}/\text{spill}$	requirement
0.92	–
$1.8 \cdot 10^{-3}$	veto (V)
$6.5 \cdot 10^{-6}$	kine (K)
$< 1.1 \cdot 10^{-8}$	(IP&VTX, fully reco.)
$< 3.6 \cdot 10^{-7}$	(IP&VTX, partially reco.)
$N_{\text{muon inel. inter.}}/5 \cdot 10^{19}$ pot	
$< 2.5 \cdot 10^{-2}$ events (fully reco.)	V & K & IP & VTX
$< 0.90$ events (partially reco.)	V & K & IP & VTX

The applied selection has an efficiency of 83% on signals.



### 3. Neutrino inelastic interactions: Detector reduction



The probability to have one neutrino inelastic interaction per spill in the decay vessel is about  $8 \times 10^{-5}$ .


The decay products are made of very low-p particles.

Out of 22000 neutrino inelastic interactions, no events survive the requirements to have at least two tracks releasing hits in the first tracking chamber and with  $p > 3$  GeV,

→ probability to have a "visible" neutrino interaction in vessel down to **less than  $4 \times 10^{-9}$  per spill,**  
**and less than 0.01** in the whole SHADOWS lifetime.

→ **No event due to neutrino inelastic interactions with «air» in the decay volume survive the requirements.**


Possible evolution: from an in-vacuum vessel to a Helium-filled balloon...




**Several advantages:**

- recover part of the acceptance while remaining compliant with the integration constraints.
- reduce to zero the inelastic interactions with the vessel
- reduce the cost..

**This layout will be studied in depth for the TDR.**



# Background: Results




**Table 31.** Estimated background events in  $5 \times 10^{19}$  pot. For muon-induced background events the factor 3 difference between data and monte carlo simulation discussed in Section 8.7 has been taken into account.

background type	fully reconstructed	partially reconstructed
combinatorial di-muon	$10^{-3}$	0.7
muon inelastic interactions	$< 2.5 \cdot 10^{-2}$	$< 0.90$
neutrino inelastic interactions	$< 0.01$	$< 0.01$

All the background sources are well under control. Less than 1 event expected in the full data set.

# Physics sensitivity: Light Dark Scalar mixing with the Higgs

(mediator of sub-GeV DM interacting with SM particles; candidate for relaxion mechanism, etc.)



SHADOWS is fully complementary to HIKE-phase 1 in kaon mode, that improves by about one order of magnitude below the kaon mass.


SHADOWS can improve by three orders of magnitude over the existing bounds (mostly from LHCb) between the di-muon threshold and ~4 GeV.

Given the extremely low background, the contour plot can be interpreted as discovery plot.

SHADOWS can cover a larger area in the (still uncharted) parameter space than: DarkQuest; LHCb Run3 & Run 4 (upgrade 1); LHCb Run 6 (upgrade 2); CODEX-b; FASER2 at the Forward Physics Facility.

## Physics sensitivity: ALPs with fermion couplings

*Axions/ALPs in the MeV-GeV range are possible solution to the strong-CP problem*



**SHADOWS is complementary to HIKE-phase 1 in kaon mode,** that can improve by about one order of magnitude the current bound below the K mass, and fill the gap down to BBN.


**SHADOWS can improve by two orders of magnitude over the existing experimental bounds (LHCb)** between the di-muon threshold and  $\sim 4$  GeV.

Given the extremely low background, **the contour plot can be interpreted as discovery plot.**

**SHADOWS can cover a larger area than: FASER2** at the Forward Physics Facility; and is very similar to CODEXb with the full data set at the end of the HL-LHC ( $3 \text{ ab}^{-1}$ ).

# Physics sensitivity: HNL with electron couplings

*Possible solution to the origin of the neutrino masses and matter-antimatter asymmetry*



**Fully complementary with HIKE phase 1 and BD**

Below the K and above the B threshold. Synergistic  
Between K and D thresholds.

**Between D and B thresholds**


**SHADOWS improves by about one order of magnitude** over the existing experimental bounds (Belle & CMS)

Given the extremely low background, **the contour plot can be interpreted as discovery plot.**

**Between K and D thresholds SHADOWS improves by about four orders of magnitude** over the existing experimental bounds (Belle)

# Physics sensitivity: HNL with muon couplings

*Possible solution to the origin of the neutrino masses and matter-antimatter asymmetry*



Fully complementary with HIKE phase 1 and BD below the K and above the B threshold. Synergistic between K and D thresholds.

Between D and B thresholds SHADOWS improves by about one order of magnitude over the existing experimental bounds (CMS)


Between K and D thresholds SHADOWS improves by about one order of magnitude over the existing experimental bounds (NuTeV)

Given the extremely low background, the contour plot can be interpreted as discovery plot.



# Physics sensitivity: HNL with tau couplings

Possible solution to the origin of the neutrino masses and matter-antimatter asymmetry




Between D and B thresholds SHADOWS improves by two orders of magnitude over the existing experimental bounds (DELPHI)

Given the extremely low background, the contour plot can be interpreted as discovery plot.


Up to D threshold SHADOWS improves by two-four orders of magnitude over the existing experimental bounds (ArgoNeut & BaBar) and is better than DarkQuest, CODEX-b and FASER2.

## The North Area Neutrino Detector (NaNu)

NaNu magnet already available at CERN




Tungsten-emulsion detector  
Interleaved with MM active layers



- Magnet-System
- MM Tracking Layers
- Tungsten/Emulsion
- Air/Emulsion
- Scintillator
- Support Structures
- Led Shield

NaNu

Super-NaNu



# Neutrino Physics with NaNu

Expected number of detectable neutrino interactions within the NaNu detector for  $4 \times 10^{19}$  POT for NaNu and Super-NaNu.

Experimental Setup	NaNu	Super-NaNu
$\nu_e$	$4.1 \times 10^3$	$20 \times 10^3$
$\bar{\nu}_e$	$1.0 \times 10^3$	$4.5 \times 10^3$
$\nu_\mu$	$40 \times 10^3$	$40 \times 10^3$
$\bar{\nu}_\mu$	$9 \times 10^3$	$9 \times 10^3$
$\nu_\tau$	$0.12 \times 10^3$	$0.72 \times 10^3$
$\bar{\nu}_\tau$	$0.07 \times 10^3$	$0.41 \times 10^3$

Overview of various **tau decay channels** including their branching ratio (BR) together with the efficiencies of various selection and identification criteria

Decay-Channel	$\tau \rightarrow e$	$\tau \rightarrow \mu$	$\tau \rightarrow h(\pi^\pm)$	$\tau \rightarrow 3h(3\pi^\pm)$
BR	0.17	0.18	0.46	0.12
Geometrical	0.9	0.9	0.9	0.9
Decay search	0.6	0.6	0.6	0.6
PID	1.0	0.9	0.9	0.9
Total Events (NaNu)	10	10	30	10
Total Events (Super-NaNu)	60	60	180	45

**$\nu(\tau)$**

Decay-Channel	$\bar{\tau} \rightarrow e$	$\bar{\tau} \rightarrow \mu$	$\bar{\tau} \rightarrow h(\pi^\pm)$	$\bar{\tau} \rightarrow 3h(3\pi^\pm)$
BR	0.17	0.18	0.46	0.12
Geometrical	0.9	0.9	0.9	0.9
Decay search	0.6	0.6	0.6	0.6
PID	1.0	0.9	0.9	0.9
Total Events (NaNu)	7	7	20	5
Total Events (Super-NaNu)	40	40	115	30

**anti- $\nu(\tau)$**

## Physics programme

- **Deep inelastic scattering of  $\nu(\mu)$  with 5-10% precision**, measurement of charm production sensitive to s-quark content in nucleons (important for W mass measurement).
- **First observation of anti-  $\nu(\tau)$**
- **measurement of  $\nu(\tau)$  and anti-  $\nu(\tau)$  inclusive cross-section** at 10% (5%) at NaNu (superNaNu), with possible observation of F4 and F5 structure function effects.
- **study of  $\nu(\tau)$  (anomalous) magnetic moment**

# Project Schedule

2023	2024	2025	2026	2027	2028	2029	2030	2031
NA62 Run			LS3	LS3	LS3	ECN3/HI Installation/ commissioning	ECN3/HI Installation/ commissioning	ECN3/HI run
Proposal	TDR	TDR	TDR/PRR	Production	Production	Production/ Installation	Installation/ Pilot Run	SHADOWS run
2032	2033	2034	2035	2036	2037	2038	2039	2040
ECN3/HI run	LS4	ECN3/HI Run					LS5	
SHADOWS run	consolidation	SHADOWS run	SHADOWS run	SHADOWS run	SHADOWS run	SHADOWS run	consolidation	SHADOWS run

**If approved, SHADOWS will be ready to start data taking in 2030 and collect  $5 \times 10^{19}$  pot by 2040.**  
(SHADOWS run beyond 2040 will depend on the compatibility with HIKE-phase 2)

# Project Organization: preliminary groups interest

**Table 41.** Preliminary group interests for SHADOWS sub-detectors and activities.

Item	Technology	Interested groups
MIB system	magnetized iron blocks	CERN, LNF-INFN
Upstream Veto	Micromegas	INFN (Rome3, Naples)
Decay Vessel	in-vacuum	CERN
Dipole Magnet	warm	CERN
Tracker	Straws	Heidelberg
Timing Layer	scintillating bars	Freiburg
ECAL	StripCal	Mainz, KIT
Muon	scintillating tiles	INFN (LNF, Ferrara, Bologna)
Software		INFN-Rome 1, Prague
TDAQ		CERN
NaNu		Mainz/Bonn

**All detectors/activities have groups involved. Still a lot of room for new groups/collaborators.**

# Project Organization: preliminary cost estimate

**Table 40.** Preliminary cost estimate of SHADOWS sub-detectors and magnets. The cost of the NaNu experiment is reported in the last row.

Item	Technology	Cost (M€)	
MIB system	magnetized iron blocks	0.992	
Upstream Veto	Micromegas	0.860	
Decay Vessel	in-vacuum	1.0	
Dipole Magnet	warm	2.57	
Tracker	Straws	1.624	
Timing Layer	scintillating bars	0.180	
ECAL	StripCal	0.980	
Muon	scintillating tiles	1.111	
TDAQ		0.250	
<b>Total SHADOWS</b>		<b>9.567</b>	Cost uncertainty C3 class: (-(10-20)%, +(10-30)%)
<b>Total NaNu</b>		<b>2.840</b>	

**The relative small-medium size (and cost) makes SHADOWS feasible and realistic in the short timescale**  
(start production in three years from now, production lasting only two years)

## Conclusions

---


- ✓ **SHADOWS and HIKE** running simultaneously and covering complementary ranges in the FIP parameter space, *above* and *below* the kaon mass, **will become a hot spot for FIP physics in the worldwide landscape.**
- ✓ The possibility of **exploring new light and feebly-interacting phenomena** and, simultaneously, **very high-scale masses through precision measurements** in the kaon sector, **makes the combined SHADOWS + HIKE system unique worldwide.**
- ✓ The upgrade in intensity of the K12 beamline would allow CERN to have **a world-class facility with** several experiments running concurrently and covering **a broad and diverse spectrum of physics topics, which is crucial, we think, for the future of particle physics.**

# The young team who made all this possible

**Florian Stummer:**  
MIB & simulation



**Shreya Roy:**  
Tracking performance



**Nico Tosi & Valentina Cicero:**  
Muon system



**Enrico Gamberini:**  
TDAQ



**Nikos Charitonidis**  
Muon flux



**Stefano Rosati:**  
Software coordination



**L. Dittmann & C. Welschhoff:**  
Muon flux



**Laurie Nevay:**  
BDSIM package



**Brendan Regnery:**  
Muon flux



**Sebastian Ritter:**  
ECAL performance



**Paolo Iengo:**  
Micromegas



**S. Niang and L. Esposito:**  
Radiation levels




**Lukasz Krzempek:**  
integration



**Moritz Barth:**  
Muon-inel. Interactions



**Dhruv Chouan:**  
NaNu



**Torben Leeftang:**  
Nu-inel. interactions






# The young team who made all this possible


**Florian Stummer:**  
MIB & simulation



**Shreya Roy:**  
Tracking performance



**Nico Tosi & Valentina Cicero:**  
Muon system



**Enrico Gamberini:**  
TDAQ



**Nikos Charinotidis:**  
Muon flux



Thanks for your attention

**Stefano:**  
Software




**Laurie Nevay:**  
BDSIM package



**Brendan Regnery:**  
Muon flux



**Sebastian Ritter:**  
ECAL performance




**Paolo Iengo:**  
Micromegas



**S. Niang and L. Esposito:**  
Radiation levels



**Lukasz Krzempek:**  
integration




**Moritz Barth:**  
Muon-inel. Interactions



**Torben Leeflang:**  
Nu-inel. interactions





**Dhruv Chouan:**  
NaNu



SPARES

# The Detector: ECAL - performance



# Full simulation: background samples

---

**sample (a):  $N \sim 5 \times 10^9$  pot with 3 GeV threshold generated using the biasing technique**

- statistically equivalent to  $\sim 3 \times 10^{13}$  pot for the combinatorial component study.

- statistically equivalent to  $\sim 2 \times 10^{21}$  pot for the inelastic interactions study,

(the muons are forced to interact with the material with 100% probability, being probability on average  $1.4 \times 10^{-8}$ ).

**sample (b):  $N \sim 3 \times 10^8$  pot to study the neutrino interactions with the SHADOWS material with GENIE.**

- statistically equivalent to  $N \sim 5 \times 10^{19}$  pot as the neutrinos

neutrinos are forced to interact with the material with 100% probability,

while this probability is on average  $6 \times 10^{-12}$

**- sample (c):  $N \sim 10^8$  pot with 100 MeV threshold to study the muon flux in ECN3,**

generated with the beamline settings used in data and the biasing technique.

This sample is statistically equivalent to  $N \sim 1.7 \times 10^{11}$  pot