


K^*/K ratio and the time between freeze-outs for intermediate-mass Ar+Sc system at the SPS energy range

Bartosz Kozłowski for the NA61/SHINE Collaboration

Warsaw University of Technology



- $K(892)^0$ resonance
 - $\Gamma = 47.3 \text{ MeV} \rightarrow \tau = 4.17 \text{ fm}/c$
 - $K(892)^0 \rightarrow K^+\pi^-$ (BR = 2/3)
- K spectra and yields can be used as input data to Blast-Wave and Hadron Resonance Gas models
- K lifetime is comparable with time between freeze-outs ! some resonances may decay inside the fireball
- Momenta of their decay products can be modified due to elastic scatterings ! problems with experimental reconstruction of resonance via invariant mass ! suppression of the observed K yield




Motivation – time between freeze-outs

- Assuming no regeneration processes (Fig.) time between freeze-outs can be determined from (STAR, Phys.Rev.C 71, 064902, 2005; C. Markert, G. Torrieri, J. Rafelski, AIP Conf.Proc. 631, 533, 2002):

$$\frac{K}{K} j_{kinetic} = \frac{K}{K} j_{chemical} e^{-\frac{\Delta t}{\tau}} \quad (1)$$


- $\frac{K}{K} |_{chemical} - K / K$ ratio in inelastic p+p collisions
- $\frac{K}{K} |_{kinetic} - K / K$ ratio in central Ar+Sc collisions
- $\tau - K$ (892)⁰ lifetime = 4.17 fm/c
- Δt – time between chemical and kinetic freeze-outs (in K rest frame)



The picture assumes that conditions at chemical freeze-out of p+p and Ar+Sc are the same

NA61/SHINE research program:

- Strong interaction physics
 - Study the properties of the onset of deconfinement
 - Search for the critical point of strongly interacting matter
 - Direct measurement of open charm
- Neutrino and cosmic-ray physics
 - Measurements for neutrino programs (J-PARC, Fermilab)
 - Measurements for cosmic-ray physics (Pierre-Auger, KASCADE, satellite experiments)




Detector layout during Ar+Sc data taking
NA61/SHINE, Eur.Phys.J.C 84, 416, 2024


- Fixed-target, multipurpose spectrometer
- 4 TPCs – tracking and PID (by dE/dx)
- PSD – centrality selection

Analyzed data and particle identification



Negative particles



Positive particles



- Analyzed data: 0-10% central Ar+Sc collisions at 40A, 75A, 150A GeV/c ($\sqrt{s_{NN}} = 8.8, 11.9, 16.8$ GeV)
- Candidates for K^+ and π^- were selected based on their energy loss (dE/dx) in TPCs
- Particles were accepted if they were located $2\sigma_\pi$ (for π^-) and $1.5\sigma_K$ (for K^+) around their empirical parametrizations of Bethe-Bloch curves
- $\sigma_\pi = 0.052$ and $\sigma_K = 0.044$
NA61/SHINE, Eur.Phys.J.C 82, 322, 2022





Template method (NA61/SHINE, Eur.Phys.J.C 80, 460, 2020)

$$f(m_{K^+\pi}) = a T_{res}^{MC}(m_{K^+\pi}) + b T_{mix}^{DATA}(m_{K^+\pi}) + c BW(m_{K^+\pi})$$



- $T_{mix}^{DATA}(m_{K^+\pi})$ – background estimated using mixing method
- $T_{res}^{MC}(m_{K^+\pi})$ – resonance background estimated using reconstructed Monte Carlo data (combination of tracks that come from decays of resonances different than $K(892)^0$ and combination of tracks where one comes from the decay of resonance and one comes from direct production in primary interaction)
- $BW(m_{K^+\pi})$ – Breit-Wigner distribution
- a, b, c - normalisation factors

$K^*(892)^0$ p_T and m_T distributions



Transverse momentum and transverse mass spectra of $K^*(892)^0$ measured in 0-10% central Ar+Sc collisions at $0 < y < 1.5$

$K^*(892)^0$ inverse slope parameters



Values of T are larger in A+A due to radial flow

Inverse slope parameters were calculated in rapidity ranges specified below


NA61/SHINE: Ar+Sc: $0 < y < 1.5$, p+p 40 and 80 GeV/c: $0 < y < 1.5$, p+p 158 GeV/c: $0 < y < 0.5$;

NA49: p+p: $0.2 < y < 0.7$, Pb+Pb: $0.43 < y < 1.78$

NA49: Phys.Rev.C 84, 064909, 2011;




NA61/SHINE: Eur.Phys.J.C 80, 460, 2020, Eur.Phys.J.C 82, 322, 2022

$K^*(892)^0$ rapidity distributions at $0 < p_T < 1.5$ GeV/c





- Rapidity spectra of $K^*(892)^0$ were measured in 0-10% central Ar+Sc collisions at $0 < p_T < 1.5$ GeV/c
- Full symbols represent the measurements, open symbols were obtained by reflection around mid-rapidity

$K^*(892)^0$ rapidity distributions – comparison with models



- Both EPOS1.99 and FTFP-BERT do not describe measured rapidity spectra




$K^*(892)^0$ mean multiplicities



NA49: Phys.Rev.C 84, 064909, 2011;

NA61/SHINE: Eur.Phys.J.C 80, 460, 2020, Eur.Phys.J.C 82, 322, 2022

$\langle K^* \rangle / \langle K^\pm \rangle$ ratios



- Suppression of K in Ar+Sc at $\rho_{\overline{s_{NN}}} = 17\text{ GeV}$ similar to Pb+Pb
- No suppression of K observed in Ar+Sc at $\rho_{\overline{s_{NN}}} = 8\text{ GeV}$

NA49:


Phys.Rev.C 84, 064909, 2011, Phys.Rev.C 66, 054902, 2002, Phys.Rev.Lett. 94, 052301, 2005;

NA61/SHINE p+p:

Eur.Phys.J.C 80, 460, 2020, Eur.Phys.J.C 82, 322, 2022, Eur.Phys.J.C 77, 671, 2017;

NA61/SHINE Ar+Sc:

Eur.Phys.J.C 84, 416, 2024



Δt boosted by Lorentz factor $\gamma = \sqrt{1 + (hp_T / m_0 c)^2}$

(see ALICE, Phys.Lett.B 802, 135225, 2020)

$\sqrt{s_{NN}}$ (GeV)

- First results on $K^0(892)$ production in 0-10% central Ar+Sc collisions at 40A, 75A, and 150A GeV/c are presented
- Values of dn/dy for all measured energies are between values obtained from EPOS1.99 and FTTP-BERT models
- hK^0/hK^+ ratios show expected suppression of $K^0(892)$ production in Ar+Sc collisions at 150A and 75A GeV/c. There is no observed suppression in Ar+Sc collisions at 40A GeV/c
- Estimated times between freeze-outs for Ar+Sc collisions at 150A and 75A GeV/c are similar

Thank you for your attention!

This work was supported by the Polish Ministry of Science and Higher Education (WUT ID-UB) and the Polish Minister of Education and Science (contract No. 2021/WK/10)