



## XIII International Conference on New Frontiers in Physics

# Azimuthal transverse single-spin asymmetries of inclusive jets and hadrons within jets from polarized $pp$ collisions at $\sqrt{s} = 510$ GeV at STAR

Yixin Zhang (张宜新),  
for the STAR Collaboration  
Shandong University (山东大学)




Supported in part by  
 U.S. DEPARTMENT OF  
**ENERGY** | Office of  
Science

# Challenges in Transverse Single-Spin Asymmetry (TSSA)



- Anomalously large  $A_N$  in  $pp$  collisions observed for nearly 40 years




$$A_N = \frac{d\sigma^{\uparrow} - d\sigma^{\downarrow}}{d\sigma^{\uparrow} + d\sigma^{\downarrow}}$$

- LO QCD predicts  $A_N \sim 0$

G. Kane, J. Pumplin, W. Repko, Phys. Rev. Lett 41, 1689 (1978).

- Left-right asymmetries of different collaborations at different beam energies



E. C. Aschenauer et al. arXiv:1602.03922

$$x_F = \frac{2p_Z}{\sqrt{s}}$$

- Stable in different C.M. energies
- Interpreted by the twist-3 and transverse-momentum-dependent (TMD) formalisms

# Mechanisms for Transverse Single-Spin Asymmetry




- Transverse Momentum Dependent (TMD) parton distributions and fragmentation functions.

- Need two scales ( $Q$  and  $p_T$ ),  $Q \gg p_T$


✓ **Sivers effect (Sivers'90):**

Parton spin and  $k_{\perp,q}$  correlation in initial state (related to orbital angular momentum)




✓ **Collins effect (Collins'93):**

Quark spin and  $k_{\perp,q}$  correlation in fragmentation process (coupled with transversity)



- Twist-3 mechanism (Efremov-Teryaev'82, Qiu-Sterman'91):
  - Collinear/twist-3 quark-gluon correlation + fragmentation functions
  - Need one scale ( $Q$  or  $p_T$ ),  $Q, p_T \gg \Lambda_{QCD}$
  - Consistent with TMD mechanism in the overlapping kinematics region



X. Ji, J.-W. Qiu, W. Vogelsang, and F. Yuan, Phys. Rev. Lett. 97, 082002 (2006)

# TSSA of $pp$ Collisions




- Transversely polarized proton-proton collision data in recent years at STAR

Year	2011	2012	2015	2017	2022
$\sqrt{s}$ (GeV)	500	200	200	510	508
$L_{int} (pb^{-1})$	25	14	52	350	400
Polarization	53%	57%	57%	55%	52%


- Measurements at RHIC can reach values of  $Q^2$  that are more than two orders of magnitude higher than current SIDIS experiments

STAR, Phys. Lett. B 780 (2018), 332-339



- Collins effect for hadron within jet at STAR


- Separate initial and final state effects
- Jet- $p_T \sim$  hard scale; hadron  $p_T \sim$  soft scale
- Validate factorization and universality with SIDIS and  $e^+e^-$  annihilation



Z. B. Kang, X. Liu, F. Ringer and H. Xing, JHEP 11 (2017), 068


U. D'Alesio, F. Murgia and C. Pisano, Phys. Lett. B 773 (2017), 300-306

# Relativistic Heavy Ion Collider (RHIC)



- RHIC is the world's only machine capable of colliding high-energy polarized protons


# The Solenoidal Tracker At RHIC (STAR)



- Time Projection Chamber (TPC)
  - $|\eta| < 1$  and  $\phi \in [0, 2\pi]$
  - Main detector for tracking and PID
- Time Of Flight (TOF)
  - $|\eta| < 1.0$  and  $\phi \in [0, 2\pi]$
  - Improve PID of tracks
- ElectroMagnetic Calorimeter
  - BEMC:  $|\eta| < 1.0$  and  $\phi \in [0, 2\pi]$ .
  - EEMC:  $1.08 < \eta < 2.0$  and  $\phi \in [0, 2\pi]$
  - Reconstruction of photon, e,  $\pi^0$  and triggering

# Jet Reconstruction

Data jets



MC jets

GEANT

PYTHIA

➤ **Jet reconstruction :**

- Anti- $K_T$  algorithm with  $R = 0.5$
- TPC tracks and EMC energy deposition as input
- Off-axis cone method to estimate underlying event contribution

➤ **Simulation**

- PYTHIA 6.4 with STAR adjustment of Perugia 2012
- Kinematic correction & Systematic uncertainty estimation


# Angle Modulations of TSSA in $pp$ Collisions



- For  $\pi^\pm$  within jets in  $pp$  collisions, the spin dependent cross section can be expressed:

$$\frac{d\sigma^\uparrow(\phi_S, \phi_H) - d\sigma^\downarrow(\phi_S, \phi_H)}{d\sigma^\uparrow(\phi_S, \phi_H) + d\sigma^\downarrow(\phi_S, \phi_H)} \propto \begin{aligned} & A_{UT}^{\sin(\phi_S)} \sin(\phi_S) && \text{related to Sivers effect} \\ & + A_{UT}^{\sin(\phi_S-\phi_H)} \sin(\phi_S - \phi_H) && \text{related to Collins effect} \\ & + A_{UT}^{\sin(\phi_S-2\phi_H)} \sin(\phi_S - 2\phi_H) && \text{related to Collins-like effect} \\ & + A_{UT}^{\sin(\phi_S+\phi_H)} \sin(\phi_S + \phi_H) \\ & + A_{UT}^{\sin(\phi_S+2\phi_H)} \sin(\phi_S + 2\phi_H) \end{aligned}$$

- $\phi_S$ : azimuthal angle between the proton transverse spin polarization vector and jet scattering plane.
- $\phi_H$ : azimuthal angle of pion relative to the jet scattering plane.



STAR, Phys. Rev. D 97, 032004 (2018)

# Extraction of Transverse Single-Spin Asymmetries



- Cross-ratio method to extract the asymmetries of different modulations.



$$A_N \sin(\phi) = \frac{1}{P} \cdot \frac{\sqrt{N^{\uparrow}(\phi)N^{\downarrow}(\phi + \pi)} - \sqrt{N^{\downarrow}(\phi)N^{\uparrow}(\phi + \pi)}}{\sqrt{N^{\uparrow}(\phi)N^{\downarrow}(\phi + \pi)} + \sqrt{N^{\downarrow}(\phi)N^{\uparrow}(\phi + \pi)}}$$

- Cross ratio method can cancel detector efficiencies and spin dependent luminosity.
- $N^{\uparrow}$ (or  $N^{\downarrow}$ ) is the yield for a given spin state.

# Particle Identification



- Particle identification with TOF unmatched (left) and matched (right)



$$n\sigma_{dE/dx} = \frac{1}{\sigma_{\text{exp}}} \ln \left( \frac{dE/dx_{\text{meas}}}{dE/dx_{\text{theo}}} \right)$$


$$n\sigma_{TOF} = \frac{TOF_{\text{meas}} - \frac{L}{c\beta(p)}}{\sigma_{\text{eff}}}$$

- Asymmetries purification through Moore-Penrose inverse.

$$\begin{pmatrix} f_{\pi \text{ rich}}^{\pi \text{ TOF}} & f_{\pi \text{ rich}}^{K \text{ TOF}} & f_{\pi \text{ rich}}^{p \text{ TOF}} \\ f_{K \text{ rich}}^{\pi \text{ TOF}} & f_{K \text{ rich}}^{K \text{ TOF}} & f_{K \text{ rich}}^{p \text{ TOF}} \\ f_{p \text{ rich}}^{\pi \text{ TOF}} & f_{p \text{ rich}}^{K \text{ TOF}} & f_{p \text{ rich}}^{p \text{ TOF}} \\ f_{\pi \text{ rich}}^{dE/dx} & f_{\pi \text{ rich}}^{K dE/dx} & f_{\pi \text{ rich}}^{p dE/dx} \\ f_{K \text{ rich}}^{dE/dx} & f_{K \text{ rich}}^{K dE/dx} & f_{K \text{ rich}}^{p dE/dx} \\ f_{p \text{ rich}}^{dE/dx} & f_{p \text{ rich}}^{K dE/dx} & f_{p \text{ rich}}^{p dE/dx} \end{pmatrix} \begin{pmatrix} A_{\pi \text{ pure}} \\ A_K \text{ pure} \\ A_p \text{ pure} \end{pmatrix} = \begin{pmatrix} A_{\pi \text{ raw}}^{\text{TOF}} \\ A_K^{\text{TOF}} \text{ raw} \\ A_p^{\text{TOF}} \text{ raw} \\ A_{\pi \text{ raw}}^{dE/dx} \\ A_K^{\text{dE/dx}} \text{ raw} \\ A_p^{\text{dE/dx}} \text{ raw} \end{pmatrix}$$

- $f_{i \text{ rich}}^j$  : the fraction of particle type  $j$  in the  $i$ -rich sample.
- Subtract other particles contamination

# Sivers Asymmetry of Inclusive Jet at 200 GeV & 510 GeV





- Sivers asymmetries for inclusive jets are consistent with 0.
- Sensitive to twist-3 correlators associated with the gluon Sivers function
- The forward-upgraded STAR detector may observe non-zero signal

AnDY Collaboration, Phys. Lett. B 750 (2015) 660

# Sivers Asymmetry of Hadron-Tagged Jet at 200 GeV & 510 GeV



M. Abdallah et al. [STAR], Phys. Rev. D 106, 072010 (2022)




- Quark jet fractions are enhanced by tagging  $\pi^\pm$
- Asymmetries are consistent with zero at mid-rapidity
- Theoretical expectations from the KPRY model

Z. B. Kang, A. Prokudin, F. Ringer and F. Yuan, Phys. Lett. B 774 (2017)


# Collins Asymmetry of pion at 510/500 GeV



- Collins results as a function of jet  $p_T$



- Positive for  $\pi^+$  and negative for  $\pi^-$ , and increase with increasing jet  $p_T$  for  $x_F > 0$
- The asymmetries for  $x_F < 0$  are consistent with 0.




- New results are consistent with previous run11 data, but with 14 times more statistics

# Collins Asymmetry of pion at 510/200 GeV: Test the TMD Evolution




➤ As a function of jet- $x_T$



➤ As a function of hadron  $j_T$


$j_T$  : pion's transverse momentum relative to jet axis



- The high precision Collins results of 510 GeV and 200 GeV nicely align with jet  $x_T$  & hadron  $j_T$  scale, giving almost no energy dependence.
- These data provide important constraints on the scale evolution for Collins asymmetry.

# Comparison to Theoretical Calculations


- 500/510GeV, as a function of  $z$   
 $z$  : pion's longitudinal momentum fraction in jet



U. D'Alesio, F. Murgia and C. Pisano, Phys. Lett. B 773 (2017), 300-306  
Z. B. Kang, A. Prokudin, F. Ringer and F. Yuan, Phys. Lett. B 774 (2017)

- Experimental results and theories are in agreement, but model calculations undershoot the observed asymmetries.
- DMP+2013 and KPRY apply the collinear QCD evolution, assume universality and factorization
- KPRY model also apply TMD evolution beyond collinear assumption

- 200GeV, as a function of  $j_T$   
 $j_T$  : pion's transverse momentum relative to jet axis




STAR, Phys. Rev. D 106, 072010 (2022)

# Collins Asymmetry of $K$ & $p$




STAR, Phys. Rev. D 106, 072010 (2022)



- The results for  $K^+$  have a contribution from favored fragmentation of  $u$  quarks, are similar in magnitude to those for  $\pi^+$
- While the results for  $K^-$  can only come from unfavored fragmentation, are consistent with zero within uncertainties
- Fragmentation into protons is not expected to produce Collins asymmetries.


# Collins-like Asymmetry of pion

➤ 200GeV, as a function of  $z$



STAR, Phys. Rev. D 106, 072010 (2022)

➤ 500GeV, as a function of  $z$



STAR, Phys. Rev. D 97, 032004 (2018),

- Sensitive to gluon linear polarization coupled to the “Collins-like” fragmentation function
- No significant asymmetry for either collision energy

# Summary & Outlook




- New preliminary results on transverse single-spin asymmetries of jets and  $\pi^\pm$  within-jets in  $pp$  at  $\sqrt{s} = 510$  GeV with STAR 2017 data
- The high precision Collins asymmetries for  $\pi^+$  and  $\pi^-$  results at 510 GeV, in excellent consistency with 200 GeV data, no energy dependence observed.
- No significant Sivers asymmetry or Collins-like asymmetry observed in  $pp$  collision.
- A large data sample of transverse polarized  $pp$  data taken in 2022 at STAR ( $\sim 400 pb^{-1}$ ), with the forward detectors ( $2.5 < \eta < 4$ ) installed, provides a unique opportunity to study Collins and Sivers effect in the forward region.

# Back up

# Collins Asymmetry from STAR 2017 Data



- Collins results as a function of  $z$  in different jet  $p_T$  regions at 510 GeV:




- These results provide more detailed constraints on the Collins fragmentation function

# Collins Asymmetry from STAR 2017 Data




- Collins results as a function of  $j_T$  in different jet  $p_T$  regions at 510 GeV:



- These results provide more detailed constraints on the Collins fragmentation function

- Recent Collins results Highlighted in the “2023 Long Range Plan for Nuclear Science”



*M. Abdallah et al. [STAR], Phys. Rev. D 106, no.7, 072010 (2022)*

- Significantly larger than the theoretical predictions
- Providing new insight on spin–momentum correlations, challenging some contemporary theoretical models