



Varkarola, Corfu – 7 August 2011

BSM perspective on Future Colliders

Workshop on Standard Model and Beyond - 5 September 2023

Riccardo Torre


INFN Genova

Refs: [my personal view](#), [FCC-CDR](#), [Snowmass 2021](#),
[Corfu program on Future Accelerators](#)




INTRODUCTION

Which BSM?



Which BSM?



Examples:

- SM parameters
- Pole observables
e.g. \hat{S} , \hat{T} , etc.
- Higgs observables


Examples:

- DY (e.g. W , Y , etc.)
- Di-bosons
- Di-jets
- Heavy quarks ($t\bar{t}$, ...)

Examples:

- SUSY
- Top partners
- Resonances

Which BSM?



Examples:

- SM parameters
- Pole observables
e.g. \hat{S} , \hat{T} , etc.
- Higgs observables

Examples:

- DY (e.g. W , Y , etc.)
- Di-bosons
- Di-jets
- Heavy quarks ($t\bar{t}$, ...)

Examples:

- SUSY
- Top partners
- Resonances

On-shell vs off-shell precision

Compare for instance LEP and LHC sensitivity to interactions of the form

Z-pole observable

$$-\frac{\hat{S}}{4m_W^2} (H^\dagger \tau^a H) W_{\mu\nu}^a B^{\mu\nu}$$

LEP

Energy: ~100 GeV

Accuracy: ~‰-%

New physics effects not enhanced by energy

LHC "cannot" compete with LEP

LHC

Energy: ~1 TeV

Accuracy: ~10%

New physics effects not enhanced by energy

off Z-pole observable

$$-\frac{Y}{4m_W^2} (\partial_\rho B_{\mu\nu})^2$$

LEP

Energy: ~100 GeV

Accuracy: ~‰-%

New physics effects not enhanced by energy

LHC comparable with (or better than) LEP

LHC

Energy: ~1 TeV


Accuracy: ~10%

New physics effects enhanced by


$$E_{\text{LHC}}^2 / E_{\text{LEP}}^2 \sim 100$$

This comparison defines two orthogonal directions in the “precision program”


Future colliders




Future colliders in 3D




Future colliders in 3D



Why colliders?




Why colliders?



PROS

- only technique to probe higher energies model independently (agnostically)
- experimenter defined experimental setup
- repeatable (experimental \neq observational)
- exploration of unknown territory
- measurement of the SM in a new energy regime and to unprecedented precision
- answer to well posed BSM questions (aka limits) on EFT, on-shell new physics, etc

Why colliders?




PROS

- only technique to probe higher energies model independently (agnostically)
- experimenter defined experimental setup
- repeatable (experimental \neq observational)
- exploration of unknown territory
- measurement of the SM in a new energy regime and to unprecedented precision
- answer to well posed BSM questions (aka limits) on EFT, on-shell new physics, etc

CONS (difficulties)

- energy increases very slowly with time/money (driven by tech evolution, but not only)
- need to convince funding agencies (the physics case is as simple as above, but it requires understanding of the scientific method and large funds)
- long term planning is becoming harder due to the increasing speed of tech advance
- keep community engaged (or even alive)

Why colliders?



PROS

- only technique to probe higher energies model independently (agnostically)
- experimenter defined experimental setup
- repeatable (experimental \neq observational)
- exploration of unknown territory
- measurement of the SM in a new energy regime and to unprecedented precision
- answer to well posed BSM questions (aka limits) on EFT, on-shell new physics, etc

CONS (difficulties)

- energy increases very slowly with time/money (driven by tech evolution, but not only)
- need to convince funding agencies (the physics case is as simple as above, but it requires understanding of the scientific method and large funds)
- long term planning is becoming harder due to the increasing speed of tech advance
- keep community engaged (or even alive)

We have no guarantee of any discovery, but guarantees cannot be a criterion for fundamental research. There is anyway the guarantee of a spectacular physics program

Are particle colliders observatories?

Some colleagues suggested the analogy of colliders with observatories, defining the former “particle observatories” (a very smart analogy to discuss with funding agencies in the “era of observatories”!)

[Tevong You - Future colliders are particle observatories – CERN Courier](#)


I would argue colliders are much more than observatories: they are both factories and observatories at the same time. Indeed, they respect the basic requirement of experimental science (compared to observational science), that is the possibility for the experimenter to set up the experiment (including the system under study), to repeat it, and to decide when to stop it.

Collider physics is the only
general-purpose experimental
research field in fundamental
science

and the only one who delivered
revolutionary results

Which colliders?

Original from ESG 2020 by UB
Updated July 25, 2022 by MN




Snowmass Energy Frontier summary, 2211.11084

An “outsider” would argue that the most time-efficient strategy is to finalize CepC and ILC while CERN works to make FCC-hh real...

Which colliders?


- Proton collider
- Electron collider
- Muon collider
- Construction/Transformation
- Preparation / R&D

Proposals emerging from Snowmass 2021 for a US based collider




Technically Limited Timeline

To be reviewed considering progress, funding and decisions



...or, even better, while CERN works to make a muon collider real!

Where do (will) we start from?



Main goal: Find signs of New Physics

- directly: probing on-shell new physics
- indirectly: probing the effect of new physics on SM observables

direct searches

precision physics

SOME PERSPECTIVE

Measurements, searches, and global fits: a statistical perspective

Measurement

- What is usually called a “measurement” can be defined as **parameter estimation** within the SM hypothesis
- This quantifies precisely “what you see” (SM), but says nothing about “what you do not see” (NP)
- Used to extract SM inputs to searches and global fits

Search (or direct search)

- This usually refers to “direct searches” where, through a statistical **hypothesis test**, the SM is confronted with a specific alternative hypothesis
- It gives some information on how much your data prefer the SM vs a well-defined alternative model

Fit (or global fit or indirect search or EFT analysis, etc.)

- This consists of either **parameter estimation** beyond the SM or a **hypothesis test** with a general enough alternative hypothesis (e.g. EFT)
- It gives information on “what you see” as well as “what you do not see”
- Notice that usually only BSM parameters are fit, while SM ones are taken from **measurements**

The (HL-)LHC legacy

“BSM measurements” (aka global fit v2.0: SM+EFT)

- It is known that uncertainties on some SM inputs is what limits the extraction of BSM parameters and, conversely, the presence of NP may affect extraction of SM parameters

Examples: PDFs vs DY, multi-jet vs α_S , etc.

- As the knowledge of the SM increases (better predictions and more analyses become available) and the large EFT parameter space gets a “good coverage” (several channels are measured and can be combined with each others) one can build a combined likelihood for SM+EFT
- Analyses that were targeting direct searches need to be turned into “measurements”, which require a higher level of precision (e.g. di-bosons)
- A simple (and interesting) example is given by fitting EFT and PDFs together using DY data (see e.g. [Greljo et al. 2104.02723](#))

The LHC legacy (in ~20 years) is to design and accomplish the final BSM measurement (which includes the SM!)

(New) Challenges

- **Combination and correlation:** combining experimental analyses is still a big issue at the LHC, where uncertainties are parametrized differently, and correlations are not known (there is a slow progress but huge work ahead)
- **Defining observables:** observables related to precision measurements are often targeted on “SM measurements”. It is necessary to extend and optimize them towards multi-differential “BSM measurement” oriented observables (e.g. recent triple differential DY cross section). Multivariate and ML could also provide a solution.
- **Large parameter space:** when the number of parameters $>$ a few, many studies become unfeasible (a lot of work in this direction: MEM, ML techniques, MadMiner, analytic reweighting, etc.)
- **EFT in backgrounds:** EFT effects may be relevant, especially for reducible BGs
- **Theory errors:** a further complication arises when statistical uncertainties become “negligible” and theory errors start to dominate (e.g. PDFs, HO, etc.). Including theory errors in statistical analysis presents conceptual issues that need to be addressed
- **Result presentation:** not only experimental analyses, but also theory results are still shown in an ad-hoc and incomplete way (e.g. 2D contours, etc). For experiments the issue is more severe, but theorists should try to get used to always deliver the full likelihood leading to their fits, that could be used by others and as input to global fits


Still a long way to go, but the path is clear

The EFT direction(s)

EFT for the SM seems like a rather “new” (~10 years) topic for theorists

Many theorists have abandoned model building in favor of EFT

This is not a psychological effect due to the absence of new physics



Absence of new physics (and the existence of precision measurements) is a requirement for EFT to be interesting, relevant, and applicable!

EFT is the simplest and most consistent way of parametrizing the different directions in which deviations from the SM can appear (SM deformations)

It is incredibly powerful at determining what “is possible”, what “is impossible”, what “is likely” and what “is unlikely”

Measurements (and especially precision measurements) in high energy physics have little meaning if one cannot quantify the above in a consistent way

In other words, EFT provides the “alternative hypothesis” necessary for a robust statistical hypothesis test of the SM

SOME BSM

Precision physics (EWPO)


Projected sensitivity to Electroweak Precision Observables at different lepton colliders

Quantity	current	ILC250	ILC-GigaZ	FCC-ee	CEPC	CLIC380
$\Delta\alpha(m_Z)^{-1} (\times 10^3)$	17.8*	17.8*		3.8 (1.2)	17.8*	
Δm_W (MeV)	12*	0.5 (2.4)		0.25 (0.3)	0.35 (0.3)	
Δm_Z (MeV)	2.1*	0.7 (0.2)	0.2	0.004 (0.1)	0.005 (0.1)	2.1*
Δm_H (MeV)	170*	14		2.5 (2)	5.9	78
$\Delta\Gamma_W$ (MeV)	42*	2		1.2 (0.3)	1.8 (0.9)	
$\Delta\Gamma_Z$ (MeV)	2.3*	1.5 (0.2)	0.12	0.004 (0.025)	0.005 (0.025)	2.3*
$\Delta A_e (\times 10^5)$	190*	14 (4.5)	1.5 (8)	0.7 (2)	1.5 (2)	60 (15)
$\Delta A_\mu (\times 10^5)$	1500*	82 (4.5)	3 (8)	2.3 (2.2)	3.0 (1.8)	390 (14)
$\Delta A_\tau (\times 10^5)$	400*	86 (4.5)	3 (8)	0.5 (20)	1.2 (20)	550 (14)
$\Delta A_b (\times 10^5)$	2000*	53 (35)	9 (50)	2.4 (21)	3 (21)	360 (92)
$\Delta A_c (\times 10^5)$	2700*	140 (25)	20 (37)	20 (15)	6 (30)	190 (67)
$\Delta\sigma_{\text{had}}^0$ (pb)	37*			0.035 (4)	0.05 (2)	37*
$\delta R_e (\times 10^3)$	2.4*	0.5 (1.0)	0.2 (0.5)	0.004 (0.3)	0.003 (0.2)	2.5 (1.0)
$\delta R_\mu (\times 10^3)$	1.6*	0.5 (1.0)	0.2 (0.2)	0.003 (0.05)	0.003 (0.1)	2.5 (1.0)
$\delta R_\tau (\times 10^3)$	2.2*	0.6 (1.0)	0.2 (0.4)	0.003 (0.1)	0.003 (0.1)	3.3 (5.0)
$\delta R_b (\times 10^3)$	3.1*	0.4 (1.0)	0.04 (0.7)	0.0014 (< 0.3)	0.005 (0.2)	1.5 (1.0)
$\delta R_c (\times 10^3)$	17*	0.6 (5.0)	0.2 (3.0)	0.015 (1.5)	0.02 (1)	2.4 (5.0)


These are “on-shell” observables and, as expected, profit of lumi more than energy

Precision physics (EWPO)

The indirect fit to Higgs and top masses gives a pictorial feeling of the indirect precision that can be reached by future lepton colliders




Precision physics (EFT)



Precision physics (EFT)

Four fermion operators generally lead to amplitudes that grow with energy and therefore profit of a larger available energy (those involving the electron)




It would be interesting to see muon collider prospects for operators involving the muon

A 10TeV muon collider is expected to have extraordinary performances

Precision physics (EFT)


Again some operators profit of the larger energy of circular colliders



Here it would be useful to include prospects for hadron and muon colliders

Precision physics (EFT)


Operators in the third generation quark sector are separated in two classes, those that “grow with energy” and those that do not



Fantastic example of the complementarity of the different collider options
 Would be interesting to add hadron and muon collider options

Direct vs indirect: universal Z'

Y-Universal Z' , 2σ




Bound coming from the study of Drell-Yan at high invariant mass (Y-parameter)
Direct searches may be more relevant only in a small corner at low masses and

coupling

Direct vs indirect: Composite Higgs


CH models lead to several signatures that can be put into a parameter space corresponding to typical strong sector resonances mass and coupling

Different signatures, from direct production of resonances, to generated HDO set constraints in different directions



SUSY

This plots is likely “unfair” in the way it compares FCC-hh (extrapolation from LHC) with muon collider (just taking energy into account, no analysis)



Universal Z'

Universal Z' models offer a useful benchmark to compare different collider options


Here bounds from direct sensitivity plus indirect EFT sensitivity are combined

Machine	Type	\sqrt{s} (TeV)	$\int L dt$ (ab^{-1})	Source	Z' Model	5σ (TeV)	95% CL (TeV)
HL-LHC	pp	14	3	RH [424]	$Z'_{SSM} \rightarrow \text{dijet}$	4.2	5.2
				ATLAS [425]	$Z'_{SSM} \rightarrow l^+l^-$	6.4	6.5
				CMS [426]	$Z'_{SSM} \rightarrow l^+l^-$	–	6.8
				EPPSU [411]	$Z'_{Univ}(g_{Z'} = 0.2)$	–	6
ILC250, CLIC380 or FCC-ee	e^+e^-	0.25	2	ILC [427]	$Z'_{SSM} \rightarrow f^+f^-$	4.9	7.7
				EPPSU [411]	$Z'_{Univ}(g_{Z'} = 0.2)$	–	7
HE-LHC	pp	27	15	EPPSU [411]	$Z'_{Univ}(g_{Z'} = 0.2)$	–	11
				ATLAS [425]	$Z'_{SSM} \rightarrow e^+e^-$	12.8	12.8
ILC	e^+e^-	0.5	4	ILC [427]	$Z'_{SSM} \rightarrow f^+f^-$	8.3	13
				EPPSU [411]	$Z'_{Univ}(g_{Z'} = 0.2)$	–	13
CLIC	e^+e^-	1.5	2.5	EPPSU [411]	$Z'_{Univ}(g_{Z'} = 0.2)$	–	19
Muon Collider	$\mu^+\mu^-$	3	1	IMCC [421]	$Z'_{Univ}(g_{Z'} = 0.2)$	10	20
ILC	e^+e^-	1	8	ILC [427]	$Z'_{SSM} \rightarrow f^+f^-$	14	22
				EPPSU [411]	$Z'_{Univ}(g_{Z'} = 0.2)$	–	21
CLIC	e^+e^-	3	5	EPPSU [411]	$Z'_{Univ}(g_{Z'} = 0.2)$	–	24
FCC-hh	pp	100	30	RH [424]	$Z'_{SSM} \rightarrow \text{dijet}$	25	32
				EPPSU [411]	$Z'_{Univ}(g_{Z'} = 0.2)$	–	35
				EPPSU [428]	$Z'_{SSM} \rightarrow l^+l^-$	43	43
Muon Collider	$\mu^+\mu^-$	10	10	IMCC [421]	$Z'_{Univ}(g_{Z'} = 0.2)$	42	70

ALPs

Axion like particles coupling to photons are a standard benchmark for the class of feebly interacting particles


They can emerge in a wide range of masses and their parameter space needs several different experiments to be covered efficiently



Long lived (pure higgsino)

Another example of weakly interacting particles is long lived particles, usually constrained through searches of disappearing tracks


The pure higgsino scenario is a useful benchmark to compare different collider options, but the understanding of the detector is crucial for robust estimates



Heavy Neutral Lepton

New neutral leptons (like sterile neutrinos) are another challenging signature

For instance one can consider a HNL with a small coupling to muons as a benchmark signature for collider performance comparison



Conclusions

- Collider physics is the only general-purpose experimental research field in fundamental science and the only one who delivered revolutionary results
- Astroparticle/cosmology cannot replace collider, neither can tens of “smaller” particle physics experiments (they are all complementary to colliders)
- Collider physics established the SM, the best theory of Nature we have so far, and is the only experimental research direction that can guarantee a frontier scientific program (even without guaranteeing any discovery)
- The (HL)-LHC legacy may be given by “BSM measurements” which extend the concepts of SM measurement, NP search, and global fit
- There are several issues to be addressed on top of building the next collider (precise theory predictions, combination of experimental analyses, definition of observables, large parameter space and signal generation, EFT in backgrounds, treatment of TH uncertainties, etc.) but the path is clear
- Optimized schedule would suggest CepC+ILC+FCC-hh (or CepC+ILC+muon-collider) but too many other considerations are in place (political/economical/sociological) and reality is very different

THANK YOU