

Exploring a Novel Dark Hypercharge Symmetry.

Phoenix 2023

Hemant Kumar Prajapati

(In collaboration with Dr. Rahul Srivastava)

IISER, Bhopal

20-12-2023



Table of contents

- 1 Overview
- 2 Introduction
- 3 The uniqueness of SM Hypercharge
- 4 $U(1)_X$ anomaly cancellation
- 5 Dark-Hyper Charge Symmetry

SM Symmetries $\rightarrow SU(3)_C \otimes SU(2)_L \otimes U(1)_Y$

Lepton Doublet $L \rightarrow (1, 2, -1)$	Quark Doublet $Q \rightarrow (3, 2, 1/3)$	Lepton Singlet $l_R \rightarrow (1, 1, -2)$	Up quark $u_R \rightarrow (3, 1, 4/3)$	Down quark $d_R \rightarrow (3, 1, -2/3)$
$\begin{bmatrix} \nu_e \\ e \end{bmatrix}_L$	$\begin{bmatrix} u \\ d \end{bmatrix}_L$	e_R	u_R	d_R
$\begin{bmatrix} \nu_\mu \\ \mu \end{bmatrix}_L$	$\begin{bmatrix} c \\ s \end{bmatrix}_L$	μ_R	c_R	s_R
$\begin{bmatrix} \nu_\tau \\ \tau \end{bmatrix}_L$	$\begin{bmatrix} t \\ b \end{bmatrix}_L$	τ_R	t_R	b_R

Table: SM Particle Content

The Standard Model (SM) has proven to be an extremely successful theory as it effectively accounts for the fundamental forces between particles up to the energy scales examined in recent research.

However, there are sufficient reasons to believe that The SM is not the final theory and it is just an effective theory in the low energy regime.

- Evidence for the existence of Dark Matter.
- Matter–antimatter asymmetry.
- Experimental evidence of Neutrino oscillation.
- Hierarchy problem.
- Muon's anomalous magnetic dipole moment.
- Strong CP problem.

New gauged symmetries beyond the SM (BSM) are motivated by these desire to explain observations that go beyond the SM.

- The simplest and highly motivated one is an extra $U(1)_X$ gauge symmetry. New $U(1)$ symmetry is highly inspired by Grand unified theories (GUT).
- Some symmetries hightly explored in literatures are $B - L$, $L_\mu - L_\tau$, $B - 3L_\mu$, $B - 3L_\tau$, etc. [[arXiv: 2202.11002](https://arxiv.org/abs/2202.11002)]

Gauge Anomalies

- An anomaly is a symmetry of the classical theory which does not survive to the quantum theory.
- Gauge symmetry plays a crucial role in establishing unitarity and renormalizability in gauge theories. An anomaly in the gauge symmetry would have severe consequences, leading to what is termed a gauge anomaly.

$$\begin{aligned} [SU(3)_C]^2 U(1)_Y &= \sum_q Y_{q_L} - \sum_q Y_{q_R} \\ [SU(2)_L]^2 U(1)_Y &= \sum_l Y_{l_L} + 3 \sum_q Y_{q_L} \\ [U(1)_Y]^3 &= \sum_{l,q} (Y_{l_L}^3 + 3Y_{q_L}^3) - \sum_{l,q} (Y_{l_R}^3 + 3Y_{q_R}^3) \\ [G]^2 U(1)_Y &= \sum_{l,q} (Y_{l_L} + 3Y_{q_L}) - \sum_{l,q} (Y_{l_R} + 3Y_{q_R}) \end{aligned} \quad (1)$$




Figure: A triangle diagram.

The uniqueness of SM Hypercharge.



- Hypercharges of all the SM fermions adds up in way to cancel these anomalies. However these anomaly cancellation conditions does not fix the SM hypercharge uniquely. One solution is $Y_L = Y_Q = Y_{e_R} = Y_{u_R} + Y_{d_R} = 0$.

- In SM fermions mass are generated via Yukawa interactios,

$$-\mathcal{L}_Y = Y_e \bar{L} \phi e_R + Y_u \bar{Q} \tilde{\phi} u_R + Y_d \bar{Q} \phi d_R + \text{h.c.} . \quad (2)$$

- These interactions gives us two extra conditions

$$Y_{u_R} = Y_Q + Y_L - Y_{e_R}, \text{ and } Y_{d_R} = Y_Q - Y_L + Y_{e_R}. \quad (3)$$

Q	u_R	d_R	L	e_R	ϕ
$\frac{-Y_L}{3}$	$\frac{-4Y_L}{3}$	$\frac{2Y_L}{3}$	Y_L	$2Y_L$	$-Y_L$

- Hence the SM Yukawa interactions to generate SM fermion masses and the gauge anomaly cancellation conditions plays a crucial role to fix these hypercharges uniquely.

$U(1)_X$ anomaly cancellation

$$[SU(3)_C]^2[U(1)_X] = \sum_q X_{q_L} - \sum_q X_{q_R} \quad (4)$$

$$[SU(2)_L]^2[U(1)_X] = \sum_l X_{l_L} + 3 \sum_q X_{q_L} \quad (5)$$

$$[U(1)_Y]^2[U(1)_X] = \sum_{l,q} (Y_{l_L}^2 X_{l_L} + 3Y_{q_L}^2 X_{q_L}) - \sum_{l,q} (Y_{l_R}^2 X_{l_R} + 3Y_{q_R}^2 X_{q_R}) \quad (6)$$

$$[U(1)_Y][U(1)_X]^2 = \sum_{l,q} (Y_{l_L} X_{l_L}^2 + 3Y_{q_L} X_{q_L}^2) - \sum_{l,q} (Y_{l_R} X_{l_R}^2 + 3Y_{q_R} X_{q_R}^2) \quad (7)$$

$$[U(1)_X]^3 = \sum_{l,q} (X_{l_L}^3 + 3X_{q_L}^3) - \sum_{l,q} (X_{l_R}^3 + 3X_{q_R}^3) \quad (8)$$

$$[G]^2[U(1)_X] = \sum_{l,q} (X_{l_L} + 3X_{q_L}) - \sum_{l,q} (X_{l_R} + 3X_{q_R}) \quad (9)$$

- **Vector Solutions :** In the BSM scenarios, while gauging new $U(1)_X$ symmetries, vector charges are typically assigned to the SM particles. This is done to ensure the invariance of the Yukawa structure.
- $B - L, B - 3L_T, L_\mu - L_\tau$ etc.
- **Chiral Solutions :** Chiral solutions are those in which SM fermions behave non-trivially under $U(1)_X$, meaning that SM fermions are chiral under this symmetry.
- In this study, we explored these chiral solutions. We will show that the induced gauge anomalies can be cancelled by adding right-handed fermions (RHF).



Dark-Hyper Charge Symmetry



$U(1)$	Q	u_R	d_R	L	e_R	f_1	f_2	f_3	φ	χ_0
$U(1)_Y$	$\frac{1}{3}$	$\frac{4}{3}$	$-\frac{2}{3}$	-1	-2	0	0	0	1	0
$U(1)_X$	$-\frac{1}{3}$	$\frac{5}{3}$	$-\frac{7}{3}$	1	-1	10	-18	17	2	-6

■ $\rho = \frac{M_W^2}{M_Z^2 \cos^2 \theta_w}$.


■ $M_{Z'}[g_X, u_\chi] > M_Z[g_X, u_\chi]$





Production and decays of Z'

In hadronic colliders the most efficient process involving Z' production is Drell-Yan. At parton level it can be written as

$$q\bar{q} \rightarrow Z', \quad (10)$$




- In the DHC symmetry, the total branching fraction of invisible decay is approximately 90% when the branching fraction saturates. In contrast, in the $B - L$ symmetry, it is about 38%.
- in the fermionic decay modes, the dileptonic branching fraction, which is maximum in $B - L$ symmetry (25%), is minimum in DHC symmetry (0.5%).





Collider Constraints

We used the ATLAS search for Z' in Dilepton resonance at pp collisions with ($\sqrt{s} = 13$) TeV and an integrated luminosity of 139 fb^{-1} , [arXiv:1903.06248 \[hep-ex\]](https://arxiv.org/abs/1903.06248). Additionally, we incorporated results from CMS, [arXiv:2103.02708 \[hep-ex\]](https://arxiv.org/abs/2103.02708).




Dark Matter Constraints

■ $0.1126 \leq \Omega h^2 \leq 0.1246$.



Feynman diagrams contributing to DM annihilation and nucleon scattering:



Conclusions



- Extensions of the Standard Model with $U(1)$ gauge symmetries hold strong motivations. These new symmetries alter the SM's gauge anomaly conditions, placing constraints on charges of new fermions beyond the SM.
- We have examined such chiral solutions, providing a set of solutions for gauge anomaly cancellation using three new right-handed BSM fermions (RHF's).
- Our presented solution involves RHF's with higher $U(1)_X$ charges compared to SM fermions. These RHNs serve as promising dark matter candidates, with their interactions through Z' being sufficient to fulfil dark matter properties. We confirm that our dark matter candidate, F_1 , with a mass range $M_{F_1} \gtrsim 150$ GeV, satisfies all the DM relevant properties and current constraints.

Thank You

Extras

The covariant derivative is defined as

$$D_\mu = \partial_\mu + ig_s T_g^a G_\mu^a + ig T_w^a W_\mu^a + ig' \frac{Y}{2} B_\mu + ig_x X C_\mu. \quad (11)$$

After the breaking of both the electroweak symmetry and $U(1)_X$, the vev of these scalar fields can be represented as follows:

$$\langle \varphi \rangle = \frac{1}{\sqrt{2}} \begin{bmatrix} 0 \\ v \end{bmatrix}, \quad \langle \chi_i \rangle = \frac{v_i}{\sqrt{2}}. \quad (12)$$

The mass spectrum of the gauge bosons are generated by the expansion of the kinetic terms of the scalars, as given below

$$(D_\mu)^\dagger D^\mu + (D_\mu \chi_i)^\dagger D^\mu \chi_i, \quad (13)$$

here repeated indices are summed over. Replacing the fields and covariant derivatives by the expression defined in Eq. (12) and Eq. (11), we can write the mass matrix of the gauge bosons in the basis (B^μ, W_3^μ, C^μ) as

$$\mathcal{M}_V^2 = \frac{v^2}{4} \begin{pmatrix} g'^2 & -gg' & 2g' X_\varphi g_x \\ -gg' & g^2 & -2g X_\varphi g_x \\ 2g' X_\varphi g_x & -2g X_\varphi g_x & 4u^2 g_x^2 \end{pmatrix}, \quad (14)$$

where $u^2 = X_\varphi^2 + u_X^2/v^2$, and u_X is defined as $u_X = \sqrt{\sum_i (X_{\chi_i}^2 v_i^2)}$.



$$m^2 = \frac{v^2}{8} (A_0 - \sqrt{B_0^2 + C_0^2}), \quad M^2 = \frac{v^2}{8} (A_0 + \sqrt{B_0^2 + C_0^2}),$$

where $A_0 = g^2 + g'^2 + 4u^2 g_x^2$, $B_0 = 4X_\varphi g_x \sqrt{g^2 + g'^2}$, $C_0 = 4u^2 g_x^2 - (g^2 + g'^2)$.
And the W boson mass is given as $M_W^2 = (gv)^2/4$.

$$\mathcal{L}_S = (D^\mu \varphi)^\dagger (D_\mu \varphi) + (D^\mu \chi_0)^\dagger (D_\mu \chi_0) - \mathcal{V}(\varphi, \chi_0), \quad (16)$$

$$\mathcal{V}(\varphi, \chi_0) = m_{\chi_0}^2 (\chi_0^* \chi_0) + \frac{1}{2} \lambda_{\chi_0} (\chi_0^* \chi_0)^2 + m_\varphi^2 (\varphi^\dagger \varphi) + \frac{1}{2} \lambda_\varphi (\varphi^\dagger \varphi)^2 + \lambda_{\varphi \chi_0} (\chi_0^* \chi_0) (\varphi^\dagger \varphi). \quad (17)$$

$$\varphi = \frac{1}{\sqrt{2}} \begin{bmatrix} \sqrt{2} G^+ \\ v + R_\varphi + i I_\varphi \end{bmatrix}, \quad \chi_0 = \frac{1}{\sqrt{2}} (v_0 + R_0 + i I_0). \quad (18)$$

First we solve the minimization equations for the mass parameters, m_{χ_0} and m_φ .

$$\begin{aligned} 2m_\varphi^2 + v^2 \lambda_\varphi + v_0^2 \lambda_{\varphi \chi_0} &= 0, \\ 2m_{\chi_0}^2 + v^2 \lambda_{\varphi \chi_0} + v_0^2 \lambda_{\chi_0} &= 0. \end{aligned} \quad (19)$$

G^\pm are the Goldstone boson corresponding to W^\pm . I_φ and I_0 will mix and give rise to the Goldstone bosons corresponding to the neutral gauge bosons Z and Z' . The mass matrix of CP-even Higgs scalars in the basis (R_1, R_2) reads as

$$\mathcal{M}_R^2 = \begin{pmatrix} A & C \\ C & B \end{pmatrix} = \begin{pmatrix} v^2 \lambda_\varphi & v v_0 \lambda_{\varphi \chi_0} \\ v v_0 \lambda_{\varphi \chi_0} & v_0^2 \lambda_{\chi_0} \end{pmatrix}. \quad (20)$$

The mass eigenvalues of light and heavy mass eigenstates as

$$M_{H_1}^2 = \frac{1}{2} \left[A + B - \sqrt{(A - B)^2 + 4C^2} \right], \quad (21)$$

$$M_{H_2}^2 = \frac{1}{2} \left[A + B + \sqrt{(A - B)^2 + 4C^2} \right]. \quad (22)$$

We follow the convention $M_{H_1} < M_{H_2}$ and have identified H_1 as the SM Higgs, with mass $M_{H_1} = 125$ GeV. The two mass eigenstates H_1, H_2 are related with the (R_1, R_2) fields through the following rotation matrix as

$$\begin{bmatrix} H_1 \\ H_2 \end{bmatrix} = \begin{bmatrix} \cos \beta & -\sin \beta \\ \sin \beta & \cos \beta \end{bmatrix} \begin{bmatrix} R_\phi \\ R_0 \end{bmatrix}, \text{ with } \tan 2\beta = \frac{2C}{B - A}. \quad (23)$$