

Contribution ID: 4405 Type: Oral Competition (Graduate Student) / Compétition orale (Étudiant(e) du 2e ou 3e cycle)

(G*) Probing Hidden Topology with Quantum Detectors

Tuesday 28 May 2024 16:45 (15 minutes)

We consider the transition rate of a static Unruh-DeWitt particle detector in a variety of spacetimes built out of quotients of AdS_3 spacetime. In particular, we contrast the behavior of a Unruh-DeWitt detector interacting with a quantum scalar field in the \mathbb{RP}^2 geon spacetime and a spacetime constructed by Aminneborg et al. The Wightman functions of these spacetimes are obtained using the method of images. We find a number of features that distinguish the two spacetimes, which are identical outside of the black hole's event horizon, most notably, in the response functions of gapless detectors in the sharp-switching limit. This points to a way in which the interior topology of a black hole may be discerned by an external observer.

Keyword-1

Quantum Fields

Keyword-2

Black Hole

Keyword-3

Particle Detector

Primary author: BHATTACHARYA, Dyuman

Co-authors: Prof. MANN, Robert (University of Waterloo); Prof. LOUKO, Jorma (University of Nottingham)

Presenter: BHATTACHARYA, Dyuman

Session Classification: (DTP/DQI) T3-2 Holography and Complexity | Holographie et complexité (DPT/DIQ)

Track Classification: Technical Sessions / Sessions techniques: Theoretical Physics / Physique théorique (DTP-DPT)