

Calculation of Direct photon production in nuclear collisions

High p_T at LHC workshop, Utrecht

Jan Cepila

jan.cepila@fjfi.cvut.cz

CFRJS @ Faculty of Nuclear Sciences and Physical Engineering,
Czech Technical University in Prague,
Czech Republic





Outline of this talk

- ▶ Introduction and motivation
 - ▶ Properties of the nuclear suppression
 - ▶ Effects contributing to the suppression
- ▶ Calculation of the direct photon production
 - ▶ Direct photon production in pp via cda
 - ▶ Coherent scenario of the direct photon production in nuclear collisions
- ▶ Numerical Results
 - ▶ p(d)-A collisions
 - ▶ A-A collisions
- ▶ Conclusions






Introduction and motivation

- ▶ BRAHMS(RHIC) observed a suppression of particles produced at forward rapidity - described by coherence effects
- ▶ Data at lower energies - NA49(SPS), E772(FNAL) - suggest similar suppression where no CGC or shadowing is possible
- ▶ Kinematics
 - ▶ Light front momentum fraction of the projectile and the target


$$x_1 = \frac{m_T}{\sqrt{s}} e^y \quad x_2 = \frac{m_T}{\sqrt{s}} e^{-y}$$
 - ▶ Feynman variable $x_F = x_1 - x_2$
- ▶ High x_1 can be achieved also at midrapidity at high p_T - at RHIC

$$p_T = 30 \text{ GeV} \sim x_1 = 0.3$$



Introduction and motivation

- ▶ The magnitude of observed suppression grows with rapidity (or x_1)
- ▶ We propose energy independent mechanism based on energy sharing restrictions in multiple interactions that lead to x_1 scaling
- ▶ The suppression comes from interplay of several effects - coherence effects (quark and gluon shadowing) and energy sharing restrictions
- ▶ Each of them is dominant in certain kinematic region - all of them has to be included in the calculation.



Coherence effects


- ▶ Controlled by the coherence length

$$l_c = \frac{1}{q_L} = \frac{2E_q\alpha(1-\alpha)}{\alpha^2 m_q^2 + p_T^2}$$

- ▶ p_T and α are transverse momentum and the fraction of the light-cone momentum of the quark carried out by the photon
- ▶ $E_q = x_q \frac{s}{2m_N}$ and m_q are the energy and mass of the projectile quark
- ▶ $q_L = \frac{M_{q\gamma}^2 - m_q^2}{2E_q}$ is the longitudinal momentum transfer

- ▶ Corresponds to lowest Fock component $|\bar{q}q\rangle$ that represents the highest twist shadowing correction

- ▶ But - coherence length drops with increasing p_T and at large quark masses as $1/m_q^2$ - almost no suppression from coherence effects at high p_T





Gluon shadowing

- ▶ Gluon shadowing added via $\sigma_{q\bar{q}}^N(\rho, x) \rightarrow \sigma_{q\bar{q}}^N(\rho, x) \times R_G(x, Q^2, b)$

$$R_G(x, Q^2, b) = \frac{G_A(x, Q^2, b)}{AG_N(x, Q^2, b)}$$

B.Z.Kopeliovich, A.Schaefer and A.V.Tarasov, Phys. Rev. D **62**, 054022 (2000)

- ▶ Represents the leading twist correction to shadowing corresponding to higher Fock components with gluons
- ▶ Dominant at small scales and $x \lesssim 10^{-3}$




B.Z.Kopeliovich, J.Raufeisen, A.V.Tarasov and M.B.Johnson, Phys. Rev. C **67**, 014903 (2003)



Effective energy loss

- ▶ We propose mechanism based on the energy sharing problem at large p_T induced by multiple initial state interactions
- ▶ One can interpret the suppression as a survival probability of the LRG in multiple interactions inside the nucleus
- ▶ Any process $a+b \rightarrow c+X$ at $x_1 \rightarrow 1$ is a LRG process



- ▶ The probability to radiate no gluons in the interval Δy is suppressed by Sudakov form factor $S(\Delta y)$
- ▶ Assuming an uncorrelated Poisson distribution for gluons, the probability to have a rapidity gap Δy is $S(\Delta y) = e^{-\langle n_G(\Delta y) \rangle}$, where the mean number of gluons is $\langle n_G(\Delta y) \rangle = \Delta y \frac{dn_G}{dy}$
- ▶ Using a formula $\Delta y = \ln\left(\frac{1}{1-x_1}\right)$ one has $S(\Delta y) = (1-x_1)^{\frac{dn_G}{dy}}$
- ▶ The height of the plateau in the gluon spectrum was estimated as

$$\frac{dn_G}{dy} = \frac{3\alpha_S}{\pi} \ln\left(\frac{m_\rho^2}{\Lambda_{QCD}^2}\right) \sim 1$$





Energy conservation restrictions

- ▶ Every additional inelastic interaction then contributes an extra suppression factor $S(x_1) \sim 1 - x_1$
- ▶ The probability of an n-fold inelastic collision is related to the Glauber coefficients via AGK cutting rules

$$v_n(b, z) = e^{-\sigma_{eff} T_A(b, z)} \frac{(\sigma_{eff} T_A(b, z))^n}{n!} \quad \sigma_{eff} = 20mb$$

- ▶ Resuming over number of rescatterings leads to

$$f_{q/N}^A(x_1, Q^2, b, z) = \sum_{n=0}^A v_n(b, z) f_{q/N}^n(x_1, Q^2)$$

- ▶ Structure function acquire additional Sudakov factor for each scattering


$$f_{q/N}^n(x_1, Q^2) = C_n f_{q/N}(x_1, Q^2) S^n(x_1)$$

- ▶ Structure function depends on the target → breakdown of the QCD factorization(leading twist effect)



Direct photon production in pp

- ▶ We use the light-cone color dipole approach to calculate the direct photon production cross-section
- ▶ In the target rest frame, the photon production looks like the bremsstrahlung of a real massless photon
- ▶ The quark fluctuates into the coherent state $|q\gamma\rangle$ that is disrupted by the color interaction with a nucleon



- ▶ Using LC wave functions and dipole cross section from DIS

$$\frac{d\sigma(qp \rightarrow \gamma X)}{d\ln\alpha d^2p_T} = \frac{1}{(2\pi)^2} \int d^2\rho_1 d^2\rho_2 e^{i\vec{p}_T \cdot (\rho_1 - \rho_2)} \Psi_{\gamma q}(\alpha, \rho_1) \Psi_{\gamma q}^*(\alpha, \rho_2) \Sigma(\alpha, \rho_1, \rho_2)$$

$$\Sigma(\alpha, \rho_1, \rho_2) = \left(\sigma_{q\bar{q}}^N(\alpha\rho_1) + \sigma_{q\bar{q}}^N(\alpha\rho_2) - \sigma_{q\bar{q}}^N(\alpha|\rho_1 - \rho_2|) \right)$$

$$\frac{d\sigma(pp \rightarrow \gamma X)}{d^2p_T} = \frac{x_1}{x_1 + x_2} \int_{x_1}^1 \frac{d\alpha}{\alpha^2} \sum_q Z_q^2 \left(f_q\left(\frac{x_1}{\alpha}\right) + f_{\bar{q}}\left(\frac{x_1}{\alpha}\right) \right) \frac{d\sigma(qp \rightarrow \gamma X)}{d\ln\alpha d^2p_T}$$



Direct photon production in pp


- Essential components for the calculation:
 - Dipole cross sections - revised GBW parametrization

$$\sigma_{q\bar{q}}^N(x, r) = 23.9 \text{mb} \left(1 - e^{-\frac{r^2 Q_0^2}{4r_0(x)}} \right)$$

$$Q_0^2 = 1 \text{GeV}^2 \quad r_0(x) = \left(\frac{x}{x_0} \right)^\lambda$$

$$x_0 = 0.000111 \quad \lambda = 0.287$$

H.Kowalski, L. Motyka and G. Watt, Phys. Rev. D 74(2006)



- Light-cone wave functions

$$\Psi_{\gamma q}(\alpha, \rho_1) \Psi_{\gamma q}^*(\alpha, \rho_2) = \frac{\alpha_{em}}{\pi^2} \left[(m_q^2 \alpha^4) K_0(\epsilon \rho_1) K_0(\epsilon \rho_2) + (1 + (1 - \alpha)^2) \epsilon^2 K_1(\epsilon \rho_1) K_1(\epsilon \rho_2) \right]$$

B.Z.Kopeliovich, A.Schaefer and A.V.Tarasov, Phys. Rev. D **62**, 054022 (2000)


- Parton distribution functions - GRV98 LO parametrizations

M. Gluck, E. Reya and A. Vogt, Eur. Phys. J. C **5** (1998) 461



The direct photon production in pp

- ▶ Calculated cross section is in good agreement with data
- ▶ RHIC 200GeV at midrapidity




S. S. Adler *et al.* (PHENIX Collaboration), Phys. Rev. Lett. **98** (2007) 012002

High p_T workshop 2011 – p. 11/18



The direct photon production in pp

- ▶ No data yet available ...
- ▶ LHC 5.5TeV at midrapidity





Coherent scenario of $p(d)$ - A collisions

- ▶ Long coherence length limit - $\langle l_c \rangle \gg R_A$
 - ▶ High energy limit
 - ▶ Fluctuation arises long before the quark enters the nucleus
 - ▶ Transverse separations of the fluctuation are “frozen” through the propagation - they form eigenstates of interaction
 - ▶ Interaction with the nucleons is coherent - maximal quark shadowing
 - ▶ Glauber eikonalization can be used to evaluate the σ^{NA}


$$\sigma_{q\bar{q}}^N(\rho, x) \rightarrow \sigma_{q\bar{q}}^A(\rho, x) = 2 \int d^2b \left(1 - \left(1 - \frac{1}{2A} \sigma_{q\bar{q}}^N(\rho, x) T_A(b) \right)^A \right)$$

B.Z.Kopeliovich, J.Raufeisen, A.V.Tarasov and M.B.Johnson, Phys. Rev. C **67**, 014903 (2003)




Numerical results - d+Au@RHIC(200GeV)

- ▶ The onset of isospin effects $R \rightarrow 0.8$ at high p_T
- ▶ Effective energy loss start to manifest themselves at $p_T \gtrsim 30$ GeV
- ▶ Gluon shadowing is negligible due to high x_2 at midrapidity




Numerical results - d+Au@RHIC(200GeV)

- ▶ Magnitude of energy loss effects rises with rapidity - dominates at high p_T
- ▶ Gluon shadowing rises from almost 0% at $\eta = 0$ to $\sim 10\%$ at $\eta = 3$ and gradually decreases with p_T




Numerical results - $p+Pb@LHC(5.5TeV)$

- ▶ The QCD factorization predicts $R \rightarrow 1$ at high p_T
- ▶ Effective energy loss negligible at this p_T range - manifest themselves at much higher p_T
- ▶ Gluon shadowing $\sim 20\%$ at low p_T





Numerical results - $p+Pb@LHC(5.5TeV)$

- ▶ Effects of energy conservation rise with rapidity and are clearly observable at $p_T > 30\text{GeV}$ at $\eta = 3 - 4$ - they can be verified at LHC
- ▶ Gluon shadowing rises from $\sim 20\%$ at $\eta = 0$ to $\sim 50\%$ at $\eta = 4$ at low p_T



Numerical results - Au+Au@RHIC(200GeV)

- ▶ The onset of isospin effects - $R \rightarrow 0.8$ at high p_T
- ▶ Effective energy loss mechanism gives a stronger suppression at high p_T - in a better agreement with data



T. Sakaguchi (PHENIX Collaboration), Nucl. Phys. A **805** (2008) 355

K. Reygers (PHENIX Collaboration), J. Phys. G **35** (2008) 104045



Conclusions

- ▶ Direct photon production cross-sections were calculated within the color dipole approach in the RHIC and LHC kinematic regions
- ▶ We included coherence effects (quark and gluon shadowing) and corrections for energy conservation in our calculations to evaluate nuclear effects.
- ▶ At RHIC energy
 - ▶ Calculations of the dAu/pp production rate show $\sim 20\%$ effect of isospin
 - ▶ The suppression driven by energy sharing problem in multiple initial state interactions is weak at midrapidity at high p_T but rapidly rises with rapidity
 - ▶ The effect of the gluon shadowing is $\sim 5-10\%$






Conclusions

- ▶ At LHC energy
 - ▶ Calculations of the pPb/pp production rate show no isospin effects and consequently one should expect $R \rightarrow 1$ in accord with QCD factorization.
 - ▶ The suppression driven by energy sharing problem in multiple initial state interactions is very weak at midrapidity at high p_T and starts to play role at very forward rapidity $\eta \sim 3 - 4$
 - ▶ The effect of the gluon shadowing is $\sim 20-50\%$
- ▶ In central Au-Au collisions at RHIC the effective energy loss mechanism represents a significant effect and describes well available data also at high p_T
- ▶ Data from RHIC and LHC at forward rapidity needed (FoCal@ALICE) for better comparison





Thank you for your attention!

