



Cornering BSMs with Positivity



Alex Pomarol, IFAE & UAB (Barcelona)

based on 2211.12488 [hep-th] with C. Fernandez, F. Riva and F. Sciotti
2307.04729 [hep-th] with T. Ma and F. Sciotti

Motivation



Effective Field Theory (EFT)

good tool to describe exp. data!

Motivation




UV completion?

more ambitious goal!

Effective Field Theory (EFT)

good tool to describe exp. data!

Motivation




☞ theory of **gravitons** (GR) → strings, ...

☞ theory of **Goldstones** (Chiral Lagrangian) → Higgs mechanism, QCD, ...


☞ **BSMs** {
Dark Matter
Axions
Composite Higgs
→ UV completion?

Motivation




👉 It has been shown in many recent examples that they can provide very **powerful** constraints


UV completion for a theory of pions



UV completion for a theory of pions





UV completion for a theory of pions




Also aiming **strongly-coupled gauge theories** (QCD)
in the **large- N_c** limit:

quarks, gluons
 $SU(N_c)$



mesons ($q\bar{q}$ states), glueballs



Positivity bounds

N. Arkani-Hamed, T.-C. Huang, and Y.-T. Huang, arXiv: 2012.15849

C. de Rham, S. Melville, A. J. Tolley, and S.-Y. Zhou, arXiv: 1702.06134

B. Bellazzini, J. Elias Miro', R. Rattazzi, M. Riembau, and F. Riva, arXiv: 2011.00037

A. Sinha and A. Zahed, arXiv: 2012.04877

A. J. Tolley, Z.-Y. Wang, and S.-Y. Zhou, arXiv: 2011.02400

S. Caron-Huot and V. Van Duong, arXiv: 2011.02957

S. Caron-Huot, D. Mazac, L. Rastelli, and D. Simmons-Duffin, arXiv: 2102.08951

and much more...

- **Generalizations of the optical theorem**

forward limit:

$$2\text{Im} \left(\begin{array}{c} k_2 \\ \bullet \\ k_1 \end{array} \right) = \sum_f \int d\Pi_f \left(\begin{array}{c} k_2 \\ \bullet \\ k_1 \end{array} \right) \left(\begin{array}{c} k_2 \\ \bullet \\ k_1 \end{array} \right) \geq 0$$

Positivity bounds

N. Arkani-Hamed, T.-C. Huang, and Y.-T. Huang, arXiv: 2012.15849

C. de Rham, S. Melville, A. J. Tolley, and S.-Y. Zhou, arXiv: 1702.06134

B. Bellazzini, J. Elias Miro', R. Rattazzi, M. Riembau, and F. Riva, arXiv: 2011.00037

A. Sinha and A. Zahed, arXiv: 2012.04877

A. J. Tolley, Z.-Y. Wang, and S.-Y. Zhou, arXiv: 2011.02400

S. Caron-Huot and V. Van Duong, arXiv: 2011.02957

S. Caron-Huot, D. Mazac, L. Rastelli, and D. Simmons-Duffin, arXiv: 2102.08951

and much more...

- **Generalizations of the optical theorem:**



forward limit (tree-level):

$$\propto |g_{abi}|^2 \geq 0$$

Positivity bounds on (tree-level mediated) amplitudes

Analytical structure of amplitudes:

$\mathcal{M}(s, t)$





Positivity bounds on (tree-level mediated) amplitudes

Analytical structure of amplitudes:


$\mathcal{M}(s, t)$

complex s-plane




Positivity bounds on (tree-level mediated) amplitudes

This simple structure allows to get dispersion relations:




Positivity bounds on (tree-level mediated) amplitudes

This simple structure allows to get dispersion relations:



Positivity bounds on (tree-level mediated) amplitudes

This simple structure allows to get dispersion relations:




$$\oint \frac{\mathcal{M}(s, t)}{s^{k+1}} = 0$$

Positivity bounds on (tree-level mediated) amplitudes

This simple structure allows to get dispersion relations:

Cauchy at work:




$$\text{residue at the origin} + \text{sum of residues at the mass poles} = 0$$

(low-energy EFT parameters **related to** masses and couplings of mesons)

pion-pion scattering

J. Albert and L. Rastelli, arXiv: 2203.11950


Lets assume an **SU(2)** (isospin) global symmetry



$\pi^a \in \mathbf{3}$ massless

Goldstones from
 $SU(2) \otimes SU(2) \rightarrow SU(2)$


Extra condition from large- N_c :

Mesons = 

Isospin = $I = 1/2 \otimes 1/2 = 0, 1$


 **no $I = 2$ states**

Extra condition from large- N_c :

Mesons = 


Isospin = $I = 1/2 \otimes 1/2 = 0, 1$

 **no $I = 2$ states**




$\mathcal{M}_s^{I=2}$

cannot have poles in s


 **$I = 2$**

Extra condition from large- N_c :

Mesons = 

Isospin = $I = 1/2 \otimes 1/2 = 0, 1$

 **no $I = 2$ states**




$\mathcal{M}_t^{I=2}$

cannot have poles in t

$I = 2$


Working with $\mathcal{M}_t^{I=2}(s, u)$
(that cannot have poles in the t-channel)

crossing $s \leftrightarrow u$ invariant



Working with $\mathcal{M}_t^{I=2}(s, u)$
 (that cannot have poles in the t-channel)


crossing $s \leftrightarrow u$ invariant



$$\text{Res} \frac{\mathcal{M}_t^{I=2}(s, u)}{s^{k+1}} = \sum_i \frac{|g_{\pi\pi i}|^2}{m_i^{2k}} P_{J_i} \left(1 + \frac{2u}{m_i^2} \right)$$

Working with $\mathcal{M}_t^{I=2}(s, u)$
 (that cannot have poles in the t-channel)

crossing $s \leftrightarrow u$ invariant



$$\text{Res} \frac{\mathcal{M}_t^{I=2}(s, u)}{s^{k+1}} = \sum_i \frac{|g_{\pi\pi i}|^2}{m_i^{2k}} P_{J_i} \left(1 + \frac{2u}{m_i^2} \right)$$

Wilson coefficients

$$\mathcal{M}_t^{I=2}(s, u) \xrightarrow{s, u \rightarrow 0} \boxed{g_{1,0}}(s + u) + \boxed{g_{2,0}}(s^2 + u^2) + \boxed{g_{2,1}}su + \dots$$

Legendre pol. and derivatives (all positive!)

small u expansion:

$$k = 1 : \quad g_{1,0} + g_{2,1}u + g_{3,1}u^2 + \dots = \sum_i |g_{\pi\pi i}|^2 \left(\frac{P_{J_i}(1)}{m_i^2} + 2 \frac{P'_{J_i}(1)}{m_i^4} u + 2 \frac{P''_{J_i}(1)}{m_i^6} u^2 + \dots \right),$$

$$k = 2 : \quad g_{2,0} + g_{3,1}u + g_{4,2}u^2 + \dots = \sum_i |g_{\pi\pi i}|^2 \left(\frac{P_{J_i}(1)}{m_i^4} + 2 \frac{P'_{J_i}(1)}{m_i^6} u + 2 \frac{P''_{J_i}(1)}{m_i^8} u^2 + \dots \right),$$

$$k = 3 : \quad g_{3,0} + g_{4,1}u + g_{5,2}u^2 + \dots = \sum_i |g_{\pi\pi i}|^2 \left(\frac{P_{J_i}(1)}{m_i^6} + 2 \frac{P'_{J_i}(1)}{m_i^8} u + 2 \frac{P''_{J_i}(1)}{m_i^{10}} u^2 + \dots \right),$$

⋮

Legendre pol. and derivatives (all positive!)

small u expansion:

$$k = 1 : \quad g_{1,0} + g_{2,1}u + g_{3,1}u^2 + \dots = \sum_i |g_{\pi\pi i}|^2 \left(\frac{P_{J_i}(1)}{m_i^2} + 2 \frac{P'_{J_i}(1)}{m_i^4} u + 2 \frac{P''_{J_i}(1)}{m_i^6} u^2 + \dots \right),$$

$$k = 2 : \quad g_{2,0} + g_{3,1}u + g_{4,2}u^2 + \dots = \sum_i |g_{\pi\pi i}|^2 \left(\frac{P_{J_i}(1)}{m_i^4} + 2 \frac{P'_{J_i}(1)}{m_i^6} u + 2 \frac{P''_{J_i}(1)}{m_i^8} u^2 + \dots \right),$$


$$k = 3 : \quad g_{3,0} + g_{4,1}u + g_{5,2}u^2 + \dots = \sum_i |g_{\pi\pi i}|^2 \left(\frac{P_{J_i}(1)}{m_i^6} + 2 \frac{P'_{J_i}(1)}{m_i^8} u + 2 \frac{P''_{J_i}(1)}{m_i^{10}} u^2 + \dots \right),$$

⋮

$$g_{n,0} = \sum_i \frac{g_{i\pi\pi}^2}{m_i^{2n}}$$

$$g_{n+1,1} = \sum_i \frac{g_{i\pi\pi}^2 J_i(J_i + 1)}{m_i^{2(n+1)}}$$

all states
contribute
positively!



small u expansion:

$$\begin{aligned} k = 1 : \quad & g_{1,0} + g_{2,1}u + \boxed{g_{3,1}u^2} + \dots = \sum_i |g_{\pi\pi i}|^2 \left(\frac{P_{J_i}(1)}{m_i^2} + 2\frac{P'_{J_i}(1)}{m_i^4}u + 2\frac{P''_{J_i}(1)}{m_i^6}u^2 + \dots \right), \\ k = 2 : \quad & g_{2,0} + \boxed{g_{3,1}u} + g_{4,2}u^2 + \dots = \sum_i |g_{\pi\pi i}|^2 \left(\frac{P_{J_i}(1)}{m_i^4} + 2\frac{P'_{J_i}(1)}{m_i^6}u + 2\frac{P''_{J_i}(1)}{m_i^8}u^2 + \dots \right), \\ k = 3 : \quad & g_{3,0} + g_{4,1}u + g_{5,2}u^2 + \dots = \sum_i |g_{\pi\pi i}|^2 \left(\frac{P_{J_i}(1)}{m_i^6} + 2\frac{P'_{J_i}(1)}{m_i^8}u + 2\frac{P''_{J_i}(1)}{m_i^{10}}u^2 + \dots \right), \\ & \vdots \end{aligned}$$

due to crossing, overconstrained system!

👉 **infinite constraints in the spectrum and couplings**

small u expansion:

$$\begin{aligned} k = 1 : \quad & g_{1,0} + g_{2,1}u + \boxed{g_{3,1}u^2} + \dots = \sum_i |g_{\pi\pi i}|^2 \left(\frac{P_{J_i}(1)}{m_i^2} + 2\frac{P'_{J_i}(1)}{m_i^4}u + 2\frac{P''_{J_i}(1)}{m_i^6}u^2 + \dots \right), \\ k = 2 : \quad & g_{2,0} + \boxed{g_{3,1}u} + g_{4,2}u^2 + \dots = \sum_i |g_{\pi\pi i}|^2 \left(\frac{P_{J_i}(1)}{m_i^4} + 2\frac{P'_{J_i}(1)}{m_i^6}u + 2\frac{P''_{J_i}(1)}{m_i^8}u^2 + \dots \right), \\ k = 3 : \quad & g_{3,0} + g_{4,1}u + g_{5,2}u^2 + \dots = \sum_i |g_{\pi\pi i}|^2 \left(\frac{P_{J_i}(1)}{m_i^6} + 2\frac{P'_{J_i}(1)}{m_i^8}u + 2\frac{P''_{J_i}(1)}{m_i^{10}}u^2 + \dots \right), \\ & \vdots \end{aligned}$$

due to crossing, overconstrained system!

 **infinite constraints in the spectrum and couplings**

$$\text{e.g.} \quad \sum_i \frac{|g_{\pi\pi i}|^2}{m_i^6} J_i(J_i + 1)(J_i - 2)(J_i + 3) = 0$$

Implications of Positivity bounds

Lets assume at $|s| \rightarrow \infty$ & either t or u fixed:

$$\frac{\mathcal{M}_t^{I=2}(s, u)}{s} \xrightarrow[k=1]{} 0$$

Infinite set of Sum Rules:

$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^6} J_i(J_i + 1)(J_i - 2)(J_i + 3) = 0$$

$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^{10}} J_i(J_i - 1)(J_i + 1)(J_i + 2)(J_i^2 + J_i - 15) = 0$$

$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^{14}} J_i(J_i - 2)(J_i - 1)(J_i + 1)(J_i + 2)(J_i + 3)(J_i^2 + J_i - 28) = 0$$

⋮

Infinite set of Sum Rules:

$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^6} J_i (J_i + 1)(J_i - 2)(J_i + 3) = 0$$

$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^{10}} J_i (J_i - 1)(J_i + 1)(J_i + 2)(J_i^2 + J_i - 15) = 0$$


$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^{14}} J_i (J_i - 2)(J_i - 1)(J_i + 1)(J_i + 2)(J_i + 3)(J_i^2 + J_i - 28) = 0$$

⋮

No constraints for $J_i = 0$ states

➡ possible UV completion:

Theory of Scalars (*Higgs mechanism*)




Infinite set of Sum Rules:

$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^6} J_i(J_i + 1)(J_i - 2)(J_i + 3) = 0$$

$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^{10}} J_i(J_i - 1)(J_i + 1)(J_i + 2)(J_i^2 + J_i - 15) = 0$$

$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^{14}} J_i(J_i - 2)(J_i - 1)(J_i + 1)(J_i + 2)(J_i + 3)(J_i^2 + J_i - 28) = 0$$


$$\frac{|g_{\pi\pi 1}|^2}{m_{J=1}^6} = 9 \frac{|g_{\pi\pi 3}|^2}{m_{J=3}^6} + 35 \frac{|g_{\pi\pi 4}|^2}{m_{J=4}^6} + \dots$$


spin-1 must be in the spectrum with the largest coupling

Infinite set of Sum Rules:

$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^6} J_i(J_i + 1)(J_i - 2)(J_i + 3) = 0$$

$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^{10}} J_i(J_i - 1)(J_i + 1)(J_i + 2)(J_i^2 + J_i - 15) = 0$$

$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^{14}} J_i(J_i - 2)(J_i - 1)(J_i + 1)(J_i + 2)(J_i + 3)(J_i^2 + J_i - 28) = 0$$


$$\frac{|g_{\pi\pi 1}|^2}{m_{J=1}^6} = 9 \frac{|g_{\pi\pi 3}|^2}{m_{J=3}^6} + 35 \frac{|g_{\pi\pi 4}|^2}{m_{J=4}^6} + \dots$$

spin-1 must be in the spectrum with the largest coupling



Vector Meson Dominance (VMD),

assumed in the past to explain QCD experimental data

Infinite set of Sum Rules:

$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^6} J_i(J_i + 1)(J_i - 2)(J_i + 3) = 0$$

$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^{10}} J_i(J_i - 1)(J_i + 1)(J_i + 2)(J_i^2 + J_i - 15) = 0$$

$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^{14}} J_i(J_i - 2)(J_i - 1)(J_i + 1)(J_i + 2)(J_i + 3)(J_i^2 + J_i - 28) = 0$$

⋮

spin-2 must be in the spectrum

Infinite set of Sum Rules:

$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^6} J_i(J_i + 1)(J_i - 2)(J_i + 3) = 0$$

$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^{10}} J_i(J_i - 1)(J_i + 1)(J_i + 2)(J_i^2 + J_i - 15) = 0$$

$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^{14}} J_i(J_i - 2)(J_i - 1)(J_i + 1)(J_i + 2)(J_i + 3)(J_i^2 + J_i - 28) = 0$$

⋮



spin-3 must be in the spectrum

Infinite set of Sum Rules:

$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^6} J_i(J_i + 1)(J_i - 2)(J_i + 3) = 0$$

$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^{10}} J_i(J_i - 1)(J_i + 1)(J_i + 2)(J_i^2 + J_i - 15) = 0$$

$$\sum_i \frac{|g_{\pi\pi i}|^2}{m_i^{14}} J_i(J_i - 2)(J_i - 1)(J_i + 1)(J_i + 2)(J_i + 3)(J_i^2 + J_i - 28) = 0$$

⋮



non-scalar UV completions require **all spin states**
with couplings to pions decreasing with J

From the constraints, we find numerically (~ 50 constraint, $J_{\max} \sim 1000$):

Upper bound on couplings

(normalized to m_i^2 / F_π^2)

J	$ g_{\pi\pi i} ^2$
1	0.78
2	0.18
3	0.03

$g_{1,0}$



From the constraints, we find numerically (~ 50 constraint, $J_{\max} \sim 1000$):


Upper bound on couplings

(normalized to m_i^2 / F_π^2)

J	$ g_{\pi\pi i} ^2$	Exp. QCD	
1	0.78	\longrightarrow	0.5
2	0.18	\longrightarrow	0.18
3	0.03		

Constraints on Wilson coefficients


$$\mathcal{L} = \frac{F_\pi^2}{4} \text{Tr} (\partial_\mu U^\dagger \partial^\mu U) + L_1 \text{Tr}^2 (\partial_\mu U^\dagger \partial^\mu U) + L_2 \text{Tr} (\partial_\mu U^\dagger \partial_\nu U) \text{Tr} (\partial^\mu U^\dagger \partial^\nu U) + L_3 \text{Tr} (\partial_\mu U^\dagger \partial^\mu U \partial_\nu U^\dagger \partial^\nu U)$$


$$e^{i \sigma^a \pi^a / F_\pi}$$

Constraints on Wilson coefficients

$$\mathcal{L} = \frac{F_\pi^2}{4} \text{Tr} (\partial_\mu U^\dagger \partial^\mu U) + L_1 \text{Tr}^2 (\partial_\mu U^\dagger \partial^\mu U) + L_2 \text{Tr} (\partial_\mu U^\dagger \partial_\nu U) \text{Tr} (\partial^\mu U^\dagger \partial^\nu U) + L_3 \text{Tr} (\partial_\mu U^\dagger \partial^\mu U \partial_\nu U^\dagger \partial^\nu U)$$

$O(s^2)$: $\tilde{g}_{2,0} = 4(2L_1 + 3L_2 + L_3) \frac{M^2}{F_\pi^2}$, $\tilde{g}_{2,1} = 16L_2 \frac{M^2}{F_\pi^2}$ ← mass of the 1st meson



“Polyhedral”
bounds


↓

EFTs are
“**EFT-hedron**”

Constraints on Wilson coefficients

$$\mathcal{L} = \frac{F_\pi^2}{4} \text{Tr} (\partial_\mu U^\dagger \partial^\mu U) + L_1 \text{Tr}^2 (\partial_\mu U^\dagger \partial^\mu U) + L_2 \text{Tr} (\partial_\mu U^\dagger \partial_\nu U) \text{Tr} (\partial^\mu U^\dagger \partial^\nu U) + L_3 \text{Tr} (\partial_\mu U^\dagger \partial^\mu U \partial_\nu U^\dagger \partial^\nu U)$$


$\mathcal{O}(s^2)$: $\tilde{g}_{2,0} = 4(2L_1 + 3L_2 + L_3) \frac{M^2}{F_\pi^2}$, $\tilde{g}_{2,1} = 16L_2 \frac{M^2}{F_\pi^2}$ ← mass of the 1st meson



Constraints on Wilson coefficients

$$\mathcal{L} = \frac{F_\pi^2}{4} \text{Tr} (\partial_\mu U^\dagger \partial^\mu U) + L_1 \text{Tr}^2 (\partial_\mu U^\dagger \partial^\mu U) + L_2 \text{Tr} (\partial_\mu U^\dagger \partial_\nu U) \text{Tr} (\partial^\mu U^\dagger \partial^\nu U) + L_3 \text{Tr} (\partial_\mu U^\dagger \partial^\mu U \partial_\nu U^\dagger \partial^\nu U)$$


$\mathcal{O}(s^2)$: $\tilde{g}_{2,0} = 4(2L_1 + 3L_2 + L_3) \frac{M^2}{F_\pi^2}$, $\tilde{g}_{2,1} = 16L_2 \frac{M^2}{F_\pi^2}$ ← mass of the 1st meson



Constraints on Wilson coefficients

$$\mathcal{L} = \frac{F_\pi^2}{4} \text{Tr} (\partial_\mu U^\dagger \partial^\mu U) + L_1 \text{Tr}^2 (\partial_\mu U^\dagger \partial^\mu U) + L_2 \text{Tr} (\partial_\mu U^\dagger \partial_\nu U) \text{Tr} (\partial^\mu U^\dagger \partial^\nu U) + L_3 \text{Tr} (\partial_\mu U^\dagger \partial^\mu U \partial_\nu U^\dagger \partial^\nu U)$$


$O(s^2)$: $\tilde{g}_{2,0} = 4(2L_1 + 3L_2 + L_3) \frac{M^2}{F_\pi^2}$, $\tilde{g}_{2,1} = 16L_2 \frac{M^2}{F_\pi^2}$ ← mass of the 1st meson



Constraints on Wilson coefficients

$$\mathcal{L} = \frac{F_\pi^2}{4} \text{Tr} (\partial_\mu U^\dagger \partial^\mu U) + L_1 \text{Tr}^2 (\partial_\mu U^\dagger \partial^\mu U) + L_2 \text{Tr} (\partial_\mu U^\dagger \partial_\nu U) \text{Tr} (\partial^\mu U^\dagger \partial^\nu U) + L_3 \text{Tr} (\partial_\mu U^\dagger \partial^\mu U \partial_\nu U^\dagger \partial^\nu U)$$


O(s²): $\tilde{g}_{2,0} = 4(2L_1 + 3L_2 + L_3) \frac{M^2}{F_\pi^2}$, $\tilde{g}_{2,1} = 16L_2 \frac{M^2}{F_\pi^2}$ ← mass of the 1st meson



Constraints on Wilson coefficients

$$\mathcal{L} = \frac{F_\pi^2}{4} \text{Tr} (\partial_\mu U^\dagger \partial^\mu U) + L_1 \text{Tr}^2 (\partial_\mu U^\dagger \partial^\mu U) + L_2 \text{Tr} (\partial_\mu U^\dagger \partial_\nu U) \text{Tr} (\partial^\mu U^\dagger \partial^\nu U) + L_3 \text{Tr} (\partial_\mu U^\dagger \partial^\mu U \partial_\nu U^\dagger \partial^\nu U)$$


O(s²): $\tilde{g}_{2,0} = 4(2L_1 + 3L_2 + L_3) \frac{M^2}{F_\pi^2}$, $\tilde{g}_{2,1} = 16L_2 \frac{M^2}{F_\pi^2}$ ← mass of the 1st meson



Constraints on Wilson coefficients

$$\mathcal{L} = \frac{F_\pi^2}{4} \text{Tr} (\partial_\mu U^\dagger \partial^\mu U) + L_1 \text{Tr}^2 (\partial_\mu U^\dagger \partial^\mu U) + L_2 \text{Tr} (\partial_\mu U^\dagger \partial_\nu U) \text{Tr} (\partial^\mu U^\dagger \partial^\nu U) + L_3 \text{Tr} (\partial_\mu U^\dagger \partial^\mu U \partial_\nu U^\dagger \partial^\nu U)$$

$O(s^2)$: $\tilde{g}_{2,0} = 4(2L_1 + 3L_2 + L_3) \frac{M^2}{F_\pi^2}$, $\tilde{g}_{2,1} = 16L_2 \frac{M^2}{F_\pi^2}$ ← mass of the 1st meson




■ $J = 0$
■ $J \geq 1$

Lovelace-Shapiro:

$$\mathcal{M}^{(LS)}(s, u) = \frac{\Gamma\left(\frac{1}{2} - \frac{s}{2m_\rho^2}\right) \Gamma\left(\frac{1}{2} - \frac{u}{2m_\rho^2}\right)}{\Gamma\left(\frac{t}{2m_\rho^2}\right)}$$


$$m_n^2 = m_\rho^2(2n + 1), \quad n = 0, 1, 2, \dots$$

**stronger bounds if we assume that,
as in QCD, $J > I$ mesons are heavier**



**stronger bounds if we assume that,
as in QCD, $J > I$ mesons are heavier**


**+ the spin-1 meson, the ρ ,
has a non-zero coupling to $\pi\pi$**



**stronger bounds if we assume that,
as in QCD, $J>I$ mesons are heavier**

**+ the spin-1 meson, the ρ ,
has a non-zero coupling to $\pi\pi$**


$$\tilde{g}_\rho^2 \equiv \frac{g_{\rho\pi\pi}^2 F_\pi^2}{m_\rho^2}$$



**stronger bounds if we assume that,
as in QCD, $J>I$ mesons are heavier**

**+ the spin-1 meson, the ρ ,
has a non-zero coupling to $\pi\pi$**

$$\tilde{g}_\rho^2 \equiv \frac{g_{\rho\pi\pi}^2 F_\pi^2}{m_\rho^2}$$



**Experimental
QCD data**

Explaining the success of holography

AdS/QCD:

5D model for QCD mesons (spin=0,1):

$SU(2)_L \times SU(2)_R$ model:

Erlich+Katz+Son+Stephanov 05
Da Rold+Pomarol 05

$$\mathcal{L}_5 = \frac{M_5}{2} \text{Tr} \left[-L_{MN} L^{MN} - R_{MN} R^{MN} + |D_M \Phi|^2 + 3|\Phi|^2 \right]$$

	Experiment	AdS ₅	Deviation
m_ρ	775	824	+6%
m_{a_1}	1230	1347	+10%
m_ω	782	824	+5%
F_ρ	153	169	+11%
F_ω/F_ρ	0.88	0.94	+7%
F_π	87	88	+1%
$g_{\rho\pi\pi}$	6.0	5.4	-10%
L_9	$6.9 \cdot 10^{-3}$	$6.2 \cdot 10^{-3}$	-10%
L_{10}	$-5.2 \cdot 10^{-3}$	$-6.2 \cdot 10^{-3}$	-12%
$\Gamma(\omega \rightarrow \pi\gamma)$	0.75	0.81	+8%
$\Gamma(\omega \rightarrow 3\pi)$	7.5	6.7	-11%
$\Gamma(\rho \rightarrow \pi\gamma)$	0.068	0.077	+13%
$\Gamma(\omega \rightarrow \pi\mu\mu)$	$8.2 \cdot 10^{-4}$	$7.3 \cdot 10^{-4}$	-10%
$\Gamma(\omega \rightarrow \pi ee)$	$6.5 \cdot 10^{-3}$	$7.3 \cdot 10^{-3}$	+12%

Explaining the success of holography

AdS/QCD:

5D model for QCD mesons (spin=0,1):

$SU(2)_L \times SU(2)_R$ model:

Erlich+Katz+Son+Stephanov 05
Da Rold+Pomarol 05

$$\mathcal{L}_5 = \frac{M_5}{2} \text{Tr} \left[-L_{MN} L^{MN} - R_{MN} R^{MN} + |D_M \Phi|^2 + 3|\Phi|^2 \right]$$


	Experiment	AdS ₅	Deviation
m_ρ	775	824	+6%
m_{a_1}	1230	1347	+10%
m_ω	782	824	+5%
F_ρ	153	169	+11%
F_ω/F_ρ	0.88	0.94	+7%
F_π	87	88	+1%
$g_{\rho\pi\pi}$	6.0	5.4	-10%
L_9	$6.9 \cdot 10^{-3}$	$6.2 \cdot 10^{-3}$	-10%
L_{10}	$-5.2 \cdot 10^{-3}$	$-6.2 \cdot 10^{-3}$	-12%
$\Gamma(\omega \rightarrow \pi\gamma)$	0.75	0.81	+8%
$\Gamma(\omega \rightarrow 3\pi)$	7.5	6.7	-11%
$\Gamma(\rho \rightarrow \pi\gamma)$	0.068	0.077	+13%
$\Gamma(\omega \rightarrow \pi\mu\mu)$	$8.2 \cdot 10^{-4}$	$7.3 \cdot 10^{-4}$	-10%
$\Gamma(\omega \rightarrow \pi ee)$	$6.5 \cdot 10^{-3}$	$7.3 \cdot 10^{-3}$	+12%

Success can be understood from positivity bounds that restrict $J > 1$ mesons to contribute little to low-energy observables

Impact on BSM searches at the LHC


Higgs as a Pseudo-Goldstone boson:

Indirect probes:



deviations in
Higgs coupling


Direct probes:



Impact on BSM searches at the LHC



Higgs as a Pseudo-Goldstone boson:

Indirect probes:



deviations in Higgs coupling

Direct probes:




e.g. 1502.01701 [hep-th]

Impact on BSM searches at the LHC



Higgs as a Pseudo-Goldstone boson:

Indirect probes:



deviations in Higgs coupling

Direct probes:




$J > 1$ must **at least** contribute a **23%** to the Wilson coeff.

e.g. 1502.01701 [hep-th]

Similar structure for higher-order Wilson coeff.

$\mathcal{O}(s^3)$:



$U(1)_A$ axial anomaly

Introducing the η' (Goldstone of an anomalous symmetry):

$$U(2) \otimes U(2) \rightarrow U(2)$$


$$\hookrightarrow SU(2) \otimes SU(2) \otimes U(1)_A \otimes U(1)$$

- WZW term: *5-goldstone int.*
 - Adding external gauge-bosons:
- $\pi \rightarrow \gamma\gamma$


} $\propto \kappa$

$$\kappa = \frac{N_c}{12\pi^2 F_\pi^3}$$

but also




two q_L, q_R model



a) It cannot be mediated by scalars

☞ axial anomaly **discards** theories with **only** scalar resonances

b) Bounded by




☞ bound on the anomaly:

$$\frac{\kappa}{\sqrt{\mathcal{P}/F_\pi^2}} \leq \frac{1}{\sqrt{2}}$$

↪ pion polarizabilities

Conclusions

- Positivity bounds from **Crossing + Analyticity + Unitarity** shows the “**EFT-hedron**” structure of the Chiral Lagrangian at large- N_c



👉 **Allows to get information on possible UV completions for a theory of Goldstones!**

Two possibilities {

- scalars (Higgs mechanism)
- Higher-spin (states with all J needed)

- Higher-spin ($J > 1$) states are strongly constrained, giving a possible explanation for VMD & the success of holographic QCD
- **Axial anomaly** can distinguish between the two possibilities

Bounded from above:

$$\frac{\kappa}{\sqrt{\mathcal{P}/F_\pi^2}} \leq \frac{1}{\sqrt{2}}$$

what theory saturates it?

➡ potential interest to constrain DM scenarios (e.g. SIMPs)

RESTRICTED AREA


**MONITORED
BY VIDEO
CAMERA**




$$0 = \sum_i \frac{|g_{\pi\pi i}|^2}{m_i^{2n}} \left(\frac{2^{n-1}}{(n-1)!} P_{J_i}^{(n-1)}(1) - \mathcal{J}_i^2 \right) \quad n=2,3,4,\dots$$

$\underbrace{\hspace{10em}}$
 $\mathcal{X}_{n,1}$


$$\mathcal{J}^2 \equiv J(J+1)$$





Lets assume at $s \rightarrow \infty$ and either t or u fixed:

$$\frac{\mathcal{M}_t^{I=2}(s, u)}{s^2} \rightarrow 0$$



C The su -models

Let us consider the most general theory of a degenerate spectrum that contributes to the four-pion amplitude $\mathcal{M}(s, u)$ [7, 8]. This means that all states have equal mass m , and therefore the denominator of this amplitude is fixed to be $\mathcal{M}(s, u) \propto 1/((s - m^2)(u - m^2))$. If we further demand that Eq. (6a) and Eq. (6b) are satisfied for $k_{\min} = 1$, we are led to

$$\mathcal{M}(s, u) = \frac{a_1 m^4 + a_2 m^2 (s + u) + a_3 s u}{(s - m^2)(u - m^2)}, \quad (91)$$

where a_i are constants. The Adler's zero condition fixes $a_1 = 0$. Then, aside from a global multiplicative factor, the amplitude has only one free parameter. We can write it as

$$\mathcal{M}_1^{(su)}(s, u) = \frac{m^2 (s + u) + \lambda s u}{(s - m^2)(u - m^2)}, \quad (92)$$

where the possible values of λ are determined by unitarity. Indeed, imposing the positivity of the residues of Eq. (92), we obtain

$$-2 \leq \lambda \leq \frac{2 \ln 2 - 1}{1 - \ln 2}. \quad (93)$$

In the limiting case $\lambda = -2$, the residues of all $J > 0$ states are zero, and we are left with the scalar amplitude Eq. (22). In the other limit,


$$\lambda = \frac{2 \ln 2 - 1}{1 - \ln 2} \simeq 1.26, \quad (94)$$

$g_{\pi\pi i}^2$

0.01

10^{-5}

$m_i = m$



J

D The Lovelace-Shapiro amplitude

The Lovelace-Shapiro (LS) amplitude for the scattering of four pions is defined as [26, 27]

$$\mathcal{M}^{(\text{LS})}(s, u) = \frac{\Gamma(1 - \alpha(s))\Gamma(1 - \alpha(u))}{\Gamma(1 - \alpha(s) - \alpha(u))}, \quad (105)$$

where $\alpha(s) = \alpha_0 + \alpha's$ is referred as the Regge trajectory. We will fix the values of α_0 and α' by requiring that Eq. (106) satisfies the Adler zero condition, $\mathcal{M}^{(\text{LS})}(s, u) \rightarrow 0$ for $s, u \rightarrow 0$, and that the first pole of Eq. (106) occurs for $s = m_\rho^2$. These two conditions lead to $\alpha_0 = 1/2$ and $\alpha' = 1/(2m_\rho^2)$ [66] and then we can write

$$\mathcal{M}^{(\text{LS})}(s, u) = \frac{\Gamma\left(\frac{1}{2} - \frac{s}{2m_\rho^2}\right)\Gamma\left(\frac{1}{2} - \frac{u}{2m_\rho^2}\right)}{\Gamma\left(\frac{t}{2m_\rho^2}\right)}. \quad (106)$$

By looking at the poles of Eq. (106), one can see that the LS amplitude corresponds to a theory of higher-spin states with masses

$$m_n^2 = m_\rho^2(2n + 1), \quad n = 0, 1, 2, \dots \quad (107)$$

For a given n , there are at most $n + 1$ states with spin $J = 0, 1, \dots, n + 1$. Furthermore, Eq. (106) satisfies the condition Eq. (6a) and Eq. (6b) with $k_{\min} = 1$.

E The Coon amplitude

The Lovelace-Shapiro amplitude presented in Appendix D can be generalized to a larger class of amplitudes depending on an additional parameter q . This is the so-called Coon amplitude, which was first proposed in [28]¹¹:

$$\mathcal{M}_q(s, u) = C(\sigma, \tau, q) \prod_{n=0}^{\infty} \frac{(1 - q^{n+1})(\sigma\tau - q^{n+1})}{(\sigma - q^{n+1})(\tau - q^{n+1})}, \quad (118)$$

where $\sigma = 1 + (q - 1)(\alpha_0 + \alpha' s)$ and $\tau = 1 + (q - 1)(\alpha_0 + \alpha' u)$. As explained in Appendix D, we take $\alpha_0 = 1/2$ and $\alpha' = 1/(2m_\rho^2)$. The parameter q takes values between 0 and 1, and in the limit $q \rightarrow 1$ we recover the LS amplitude Eq. (106). There is some freedom in the choice of the prefactor C , as long as it satisfies $\lim_{q \rightarrow 1} C(\sigma, \tau, q) = 1$.

The Coon amplitude has an infinite number of simple poles at

$$s_n = m_\rho^2 \frac{1 + q - 2q^{n+1}}{1 - q}, \quad n = 0, 1, 2, \dots . \quad (119)$$