

A bottom-up approach to nucleon decay. RGEs and correlations

Arnau Bas i Beneito
4th of June, 2024
Planck 2024, Lisboa

Based on work in collaboration with J. Gargalionis, J. Herrero-García,
M. A. Schmidt. A. Santamaría [2312.13361] (Accepted for publication in JHEP)



GENERALITAT
VALENCIANA



CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS

Standard story

\mathcal{L}_{SM}

$SU(3)_C \times SU(2)_L \times U(1)_Y$

+

$H, Q_L^i, u_R^i, d_R^i, L_L^i, e_R^i, i = 1, 2, 3$

B and L accidentally conserved
(B + L violated in 3 units
by sphaleron transitions)

Individual Flavour Symmetries



Yukawa couplings

$U(1)_{L_e} \times U(1)_{L_\mu} \times U(1)_{L_\tau} \times U(1)_B$



ν oscillations

[*Super-K 1999,
KamLAND 2003...*]


$U(1)_L \times U(1)_B$



Proton stable



Experimental perspectives



Sensitivity of $n \rightarrow K^+ e^-$!
increased by 10^3 in HK

Hyper-K design report

BNV nucleon decay could be the next big discovery

Then... why nucleon decay?


- There is **no fundamental reason** to have **B** and **L** conserved (Leptoquarks, Seesaw particles, SUSY, GUTs...)
- Experimental probes of **BNV** and **LNV** constitute one of the **strongest** evidence for physics beyond SM (BSM) → PD will be looked for in future experiments (HK, DUNE...)

Grand Unified Theories (GUTs)

[Georgi et. al. 1973, Pati et al. 1974]

Baryon Number violation (BNV)

E.g. proton decay (PD), neutron-antineutron oscillations



Large number of UV theories predicting PD

Systematic study of PD in a model-independent way (bottom-up)

BNV within the SMEFT

Parametrization of new physics through Effective operators ($d > 4$)
SM Effective Field Theory (SMEFT)

Bounds on SMEFT WCs serve as a bridge to specific UV models

[S. Weinberg 1979 ,
B. Grzadkowski et al. 2010,
W. Buchmuller et al. 1986,
Brivio et al. 2019,
B. Henning et al. 2016,
De Gouvea et al. 2014]

$$\Delta L = 2$$

 \downarrow

$$\mathcal{L} = \mathcal{L}_{SM} + \frac{c^{d=5}}{\Lambda} \mathcal{O}_W + \left[\frac{c^{d=6}}{\Lambda^2} \mathcal{O}^{d=6} + \frac{c^{d=7}}{\Lambda^3} \mathcal{O}^{d=7} \right] + \dots$$

$$\Delta(B - L) = 0$$

 \downarrow


$$\Delta(B - L) = 2$$

 \downarrow

Different phenomenology \rightarrow

$$\begin{array}{ll} \Lambda \gtrsim 10^{15} \text{ GeV} & \Lambda \gtrsim 10^{10} \text{ GeV} \\ p \rightarrow \pi^0 e^+, \quad p \rightarrow K^+ \bar{\nu} & n \rightarrow \pi^+ e^-, \quad p \rightarrow K^+ \nu \end{array}$$

BNV within the SMEFT



- Assumptions: Energy Desert and no SUSY in the TeV scale/RpV

[S. Antusch et al. 2021,
H. Dreiner et al. 2020]

SMEFT

$d = 6 \rightarrow 4$ (273) operators

[B. Grzadkowski et al. 2010, L. F. Abbott et al. 1980]

$$\begin{aligned}\mathcal{O}_{qqql,pqrs} &= (Q_p^i Q_q^j)(Q_r^l L_s^k)\epsilon_{ik}\epsilon_{jl}, & \mathcal{O}_{qque,pqrs} &= (Q_p^i Q_q^j)(\bar{u}_r^\dagger \bar{e}_s^\dagger)\epsilon_{ij}, \\ \mathcal{O}_{duue,pqrs} &= (\bar{d}_p^\dagger \bar{u}_q^\dagger)(\bar{u}_r^\dagger \bar{e}_s^\dagger), & \mathcal{O}_{duql,pqrs} &= (\bar{d}_p^\dagger \bar{u}_q^\dagger)(Q_r^i L_s^j)\epsilon_{ij},\end{aligned}$$

$d = 7 \rightarrow 6$ (297) operators

[L. Lehman 2014, Yi Liao et al. 2016]

$$\begin{aligned}\mathcal{O}_{\bar{l}dddH,pqrs} &= (L_p^\dagger \bar{d}_q^\dagger)(\bar{d}_r^\dagger \bar{d}_s^\dagger)H, & \mathcal{O}_{\bar{l}dqq\tilde{H},pqrs} &= (L_p^\dagger \bar{d}_q^\dagger)(Q_r Q_s^i)\tilde{H}^j\epsilon_{ij}, \\ \mathcal{O}_{\bar{e}qdd\tilde{H},pqrs} &= (\bar{e}_p Q_q^i)(\bar{d}_r^\dagger \bar{d}_s^\dagger)\tilde{H}^j\epsilon_{ij}, & \mathcal{O}_{\bar{l}dud\tilde{H},pqrs} &= (L_p^\dagger \bar{d}_q^\dagger)(\bar{u}_r^\dagger \bar{d}_s^\dagger)\tilde{H}, \\ \cancel{\mathcal{O}_{\bar{l}qdDd,pqrs}} &= (L_p^\dagger \bar{\sigma}^\mu Q_q)(\bar{d}_r^\dagger i \cancel{D}_\mu \bar{d}_s^\dagger), & \cancel{\mathcal{O}_{\bar{e}dddD,pqrs}} &= (\bar{e}_p \sigma^\mu \bar{d}_q^\dagger)(\bar{d}_r^\dagger i \cancel{D}_\mu \bar{d}_s^\dagger),\end{aligned}$$

$$\left. \begin{aligned}c^{d=6}(m_W) &\sim (2-4) c^{d=6}(10^{15} \text{ GeV}) \\ c^{d=7}(m_W) &\sim (1-2) c^{d=7}(10^{11} \text{ GeV})\end{aligned}\right\}$$

From gauge interactions and y_t
(Operator mixing subdominant)

- RGEs for $d = 6$ SMEFT [Manohar et al. 2014]
- RGEs for $d = 7$ SMEFT [Yi Liao et al. 2016]

LEFT

$$\left. \begin{array}{l} 288 \Delta(B-L) = 0 \text{ operators} \rightarrow 14 \text{ operators} \\ 228 \Delta(B+L) = 0 \text{ operators} \rightarrow 9 \text{ operators} \end{array} \right\} \text{LEFT operators involved in nucleon decay at tree-level}$$

Name [52]	SMEFT matching
$[\mathcal{O}_{udd}^{S,LL}]_{pqrs}$	$V_{q'q} V_{r'r} (C_{qqql,r'q'ps} - C_{qqql,q'r'ps} + C_{qqql,q'pr's})$
$[\mathcal{O}_{duu}^{S,LL}]_{pqrs}$	$V_{p'p} (C_{qqql,rqp's} - C_{qqql,qrp's} + C_{qqql,qp'r's})$
$[\mathcal{O}_{duu}^{S,LR}]_{pqrs}$	$-V_{p'p} (C_{qque,p'qrs} + C_{qque,qp'r's})$
$[\mathcal{O}_{duu}^{S,RL}]_{pqrs}$	$C_{duql,pqrs}$
$[\mathcal{O}_{dud}^{S,RL}]_{pqrs}$	$-V_{r'r} C_{duql,pqr's}$
$[\mathcal{O}_{duu}^{S,RR}]_{pqrs}$	$C_{duue,pqrs}$
$[\mathcal{O}_{udd}^{S,LR}]_{pqrs}$	$-V_{q'q} C_{\bar{l}dqq\tilde{H},rspq'} \frac{v}{\sqrt{2}\Lambda}$
$[\mathcal{O}_{ddd}^{S,LR}]_{pqrs}$	$V_{p'p} V_{q'q} (C_{\bar{l}dqq\tilde{H},rsq'p'} - C_{\bar{l}dqq\tilde{H},rsp'q'}) \frac{v}{2\sqrt{2}\Lambda}$
$[\mathcal{O}_{ddd}^{S,RL}]_{pqrs}$	$V_{s's} (C_{\bar{e}qdd\tilde{H},rs'qp} - C_{\bar{e}qdd\tilde{H},rs'pq}) \frac{v}{\sqrt{2}\Lambda}$
$[\mathcal{O}_{udd}^{S,RR}]_{pqrs}$	$C_{\bar{l}dud\tilde{H},rspq} \frac{v}{\sqrt{2}\Lambda}$
$[\mathcal{O}_{ddd}^{S,RR}]_{pqrs}$	$C_{\bar{l}ddd\tilde{H},rspq} \frac{v}{\sqrt{2}\Lambda}$

Name	Operator	Flavour
$[\mathcal{O}_{udd}^{S,LL}]_{111r}$	(O_{LL}^ν)	$(\mathbf{8}, \mathbf{1})$
$[\mathcal{O}_{udd}^{S,LL}]_{121r}$	(\tilde{O}_{LL1}^ν)	$(us)(d\nu_r)$
$[\mathcal{O}_{udd}^{S,LL}]_{112r}$	(\tilde{O}_{LL2}^ν)	$(ud)(s\nu_r)$
$[\mathcal{O}_{duu}^{S,LL}]_{111r}$	(O_{LL}^e)	$(du)(ue_r)$
$[\mathcal{O}_{duu}^{S,LL}]_{211r}$	(\tilde{O}_{LL}^e)	$(su)(ue_r)$
$[\mathcal{O}_{duu}^{S,LR}]_{111r}$	(O_{LR}^e)	$(du)(\bar{u}^\dagger \bar{e}_r^\dagger)$
$[\mathcal{O}_{duu}^{S,LR}]_{211r}$	(\tilde{O}_{LR}^e)	$(su)(\bar{u}^\dagger \bar{e}_r^\dagger)$
$[\mathcal{O}_{duu}^{S,RL}]_{111r}$	(O_{RL}^e)	$(\bar{d}^\dagger \bar{u}^\dagger)(ue_r)$
$[\mathcal{O}_{duu}^{S,RL}]_{211r}$	(\tilde{O}_{RL}^e)	$(\bar{s}^\dagger \bar{u}^\dagger)(ue_r)$
$[\mathcal{O}_{dud}^{S,RL}]_{111r}$	(O_{RL}^ν)	$(\bar{d}^\dagger \bar{u}^\dagger)(d\nu_r)$
$[\mathcal{O}_{dud}^{S,RL}]_{211r}$	(\tilde{O}_{RL1}^ν)	$(\bar{s}^\dagger \bar{u}^\dagger)(d\nu_r)$
$[\mathcal{O}_{dud}^{S,RL}]_{112r}$	(\tilde{O}_{RL2}^ν)	$(\bar{d}^\dagger \bar{u}^\dagger)(s\nu_r)$
$[\mathcal{O}_{duu}^{S,RR}]_{111r}$	(O_{RR}^e)	$(\bar{d}^\dagger \bar{u}^\dagger)(\bar{u}^\dagger \bar{e}_r^\dagger)$
$[\mathcal{O}_{duu}^{S,RR}]_{211r}$	(\tilde{O}_{RR}^e)	$(\bar{s}^\dagger \bar{u}^\dagger)(\bar{u}^\dagger \bar{e}_r^\dagger)$

- RG effects universal in the LEFT

— Not generated by D = 6, 7 SMEFT ops.

$$c(2 \text{ GeV}) \sim 1.26 c(m_W)$$

B χ PT

$$M = \sum_{a=1}^8 M_a \frac{\lambda_a}{\sqrt{2}} = \begin{pmatrix} \frac{\pi^0}{\sqrt{2}} + \frac{\eta}{\sqrt{6}} & \pi^+ & K^+ \\ \pi^- & -\frac{\pi^0}{\sqrt{2}} + \frac{\eta}{\sqrt{6}} & K^0 \\ K^- & \bar{K}^0 & -\sqrt{\frac{2}{3}}\eta \end{pmatrix}$$

$$B = \sum_{a=1}^8 B_a \frac{\lambda_a}{\sqrt{2}} = \begin{pmatrix} \frac{\Sigma^0}{\sqrt{2}} + \frac{\Lambda^0}{\sqrt{6}} & \Sigma^+ & p \\ \Sigma^- & -\frac{\Sigma^0}{\sqrt{2}} + \frac{\Lambda^0}{\sqrt{6}} & n \\ \Xi^- & \Xi^0 & -\sqrt{\frac{2}{3}}\Lambda^0 \end{pmatrix}$$

Flavour group $U(3)_L \times U(3)_R$


BNC interactions

↓

$$\mathcal{L} = g_{MB}^N \bar{B} \gamma^\mu \gamma_5 N \partial_\mu M + m_{B\alpha,X} \bar{\ell}_\alpha P_{\bar{X}} B + i y_{M\alpha,X}^N \bar{\ell}_\alpha P_{\bar{X}} NM$$

BNV interactions

[M. Claudson et al. 1981,
P. Nath et al. 2007]



$\Gamma(N \rightarrow M \ell_\alpha)$


2 inputs from lattice α, β

[JLQCD 2000,
Y. Aoki et al. 2017,]

(First-time computation of $|\Delta(B - L)| = 2$ two-body decays in the B χ PT formalism)

D = 6 limits


Bounds on other flavor components can
be found in [J. Gargalionis et al. 2024]



$$\Lambda/\sqrt{c} \sim (1 \sim 10) \times 10^{15} \text{ GeV}$$


D = 7 limits

Bounds on other flavor components can
be found in [J. Gargalionis et al. 2024]




$$\Lambda/\sqrt{3c} \sim (2 \sim 12) \times 10^{10} \text{ GeV}$$

$D = 6$ pairs of WCs



- Different search channels provide complementary constraints
- No flat directions

$D = 7$ pairs of WCs





- Different search channels provide complementary constraints
- No flat directions

$$\text{Bound Saturation} \equiv \frac{\Gamma_{i,\text{th}}/\Gamma_{\max,\text{th}}}{\Gamma_{i,\text{exp}}/\Gamma_{\max,\text{exp}}}$$

$\tau_{p \rightarrow \pi^0 e^+} > 24 \cdot 10^{33}$ years

$\tau_{p \rightarrow K^+ \nu} > 6.61 \cdot 10^{33}$ years

$\tau_{n \rightarrow K^+ e^-} > 0.032 \cdot 10^{33}$ years



Phenomenological matrices



Numerical κ -matrices
available online

$$\Gamma_{(i)}^{\Delta(B-L)=0} \equiv 10^{-4} c_j^* \boxed{\kappa_{(i)}^{jk}} c_k \frac{m_p^5}{\Lambda^4} \quad \text{for} \quad i = p \rightarrow \pi^0 e^+, p \rightarrow K^+ \nu \dots \quad (9 \text{ matrices})$$


$$\Gamma_{(i)}^{|\Delta(B-L)|=2} \equiv 10^{-4} c_j^* \boxed{\kappa_{(i)}^{jk}} c_k \frac{m_p^7}{\Lambda^6} \quad \text{for} \quad i = p \rightarrow K^+ \nu, n \rightarrow K^+ e^- \dots \quad (6 \text{ matrices})$$

$$c_{qque,1111} \sim 4.1 \ c_{duue,1111}$$

$$c_{qque,1111} \sim -0.44 \ c_{duue,1111}$$



Example UV model



SM enhanced by a scalar LQ ω_2 and a VLF Q_1


$$\omega_2 \sim (3, 1, 2/3), Q_1 + \bar{Q}_1^\dagger \sim (3, 2, 1/6)$$

$$\mathcal{L}_{\text{int}} = y_{1,ij} \omega_2 \bar{d}^{\dagger i} \bar{d}^{\dagger j} + y_{2,k} H^\dagger Q_1 \bar{d}^k + y_{3,k} Q_1 \epsilon H \bar{u}^k + y_{4,l} \omega_2 \bar{Q}_1 L^l + \text{h.c.}$$

$$\mathcal{L}_{\text{eff}} \supset C_{\bar{l}dddH,pqrs} \mathcal{O}_{\bar{l}dddH,pqrs} + C_{\bar{l}dud\tilde{H},pqrs} \mathcal{O}_{\bar{l}dud\tilde{H},pqrs} + \text{h.c.}$$


y_1 antisymmetric

$$\left\{ \begin{array}{lll} p \rightarrow K^+ \nu & n \rightarrow K^0 \nu & n \rightarrow K^+ e^- \\ C_{\bar{l}dud\tilde{H},1211} = -C_{\bar{l}dud\tilde{H},1112} \end{array} \right.$$



κ -matrices for the 3 processes above and compute Γ \longrightarrow **$p \rightarrow K^+ \nu$ the most constraining**

Example UV model



$$M_{\text{eff}}^3 = \frac{M_\omega^2 M_Q}{y_{1,12} y_{4,1}^*}$$

Main results of this work

- Model-independent analysis on nucleon decay
- RG effects important: limits enhanced by 30% - 130% ($d=6$) , and 20 - 30% ($d=7$)
- Complementary analysis → Correlations and flat directions
- κ -matrices: SMEFT WC at $\Lambda \leftrightarrow$ observables at m_p
- Positive signals in 2-3 channels → SMEFT operators → GUT/Models

Main source of uncertainty: nuclear matrix elements α, β



Muito obrigado!

Backup slides

RGEs

$$\left(\dot{C}_i \equiv 16\pi^2 \mu \frac{dC_i}{d\mu} = \sum_j \gamma_{ij} C_j \right)$$

$$\dot{C}_{duue,prst} = (-4g_3^2 - 2g_1^2) C_{duue,prst} - \frac{20}{3} g_1^2 C_{duue,psrt}$$

$$\dot{C}_{duq\ell,prst} = \left(-4g_3^2 - \frac{9}{2}g_2^2 - \frac{11}{6}g_1^2 \right) C_{duq\ell,prst}$$

$$\dot{C}_{qque,prst} = \left(-4g_3^2 - \frac{9}{2}g_2^2 - \frac{23}{6}g_1^2 \right) C_{qque,prst}$$

$$\dot{C}_{qqq\ell,prst} = \left(-4g_3^2 - 3g_2^2 - \frac{1}{3}g_1^2 \right) C_{qqq\ell,prst} - 4g_2^2 \left(C_{qqq\ell,rpst} + C_{qqq\ell,srpt} + C_{qqq\ell,psrt} \right)$$

$$\dot{C}_{\bar{l}dud\tilde{H},prst} = \left(-4g_3^2 - \frac{9}{4}g_2^2 - \frac{17}{12}g_1^2 + y_t^2 \right) C_{\bar{l}dud\tilde{H},prst} - \frac{10}{3}g_1^2 C_{\bar{l}dud\tilde{H},ptsr},$$

$$\dot{C}_{\bar{l}dddH,prst} = \left(-4g_3^2 - \frac{9}{4}g_2^2 - \frac{13}{12}g_1^2 + y_t^2 \right) C_{\bar{l}dddH,prst},$$

$$\dot{C}_{\bar{e}qdd\tilde{H},prst} = \left(-4g_3^2 - \frac{9}{4}g_2^2 + \frac{11}{12}g_1^2 + y_t^2 \right) C_{\bar{e}qdd\tilde{H},prst},$$

$$\dot{C}_{\bar{l}dq\bar{q}\tilde{H},prst} = \left(-4g_3^2 - \frac{15}{4}g_2^2 - \frac{19}{12}g_1^2 + y_t^2 \right) C_{\bar{l}dq\bar{q}\tilde{H},prst} - 3g_2^2 C_{\bar{l}dq\bar{q}\tilde{H},prts}.$$

Direct VS Indirect method



Direct VS Indirect method

$$\Gamma(N \rightarrow M + \ell) = \frac{m_N}{32\pi} \left(1 - \frac{m_M^2}{m_N^2}\right)^2 \left| \sum_I C_I W_0^I(N \rightarrow M) \right|^2$$

$W_0^I(N \rightarrow M)$ computed in the lattice
(Several parameters)

$$\Gamma(p \rightarrow \pi^+ \nu_r) = (32\pi f_\pi^2 m_p^3)^{-1} (m_p^2 - m_\pi^2)^2 \left| \alpha \left[L_{udd}^{S,LR} \right]_{11r1} + \beta \left[L_{udd}^{S,RR} \right]_{11r1} \right|^2 (1 + D + F)^2$$

$$\begin{aligned} \Gamma(n \rightarrow K^+ e_r^-) &= (32\pi f_\pi^2 m_n^3)^{-1} (m_n^2 - m_K^2)^2 \times \\ &\left\{ \left| \beta \left[L_{ddd}^{S,LL} \right]_{12r1} - \alpha \left[L_{ddd}^{S,RL} \right]_{12r1} + \frac{m_n}{m_\Sigma} \left(\alpha \left[L_{ddd}^{S,RL} \right]_{12r1} + \beta \left[L_{ddd}^{S,LL} \right]_{12r1} \right) (D - F) \right|^2 \right. \\ &\quad \left. + \left| \beta \left[L_{ddd}^{S,RR} \right]_{12r1} - \alpha \left[L_{ddd}^{S,LR} \right]_{12r1} + \frac{m_n}{m_\Sigma} \left(\alpha \left[L_{ddd}^{S,LR} \right]_{12r1} + \beta \left[L_{ddd}^{S,RR} \right]_{12r1} \right) (D - F) \right|^2 \right\} \end{aligned}$$

D, F, f_π
Low-energy B χ PT constants






$$\begin{aligned} \Gamma(n \rightarrow K^0 \nu_r) &= (32\pi f_\pi^2 m_n^3)^{-1} (m_n^2 - m_K^2)^2 \times \\ &\left| -\alpha \left[L_{udd}^{S,LR} \right]_{12r1} + \beta \left[L_{udd}^{S,RR} \right]_{12r1} + \alpha \left[L_{udd}^{S,LR} \right]_{11r2} + \beta \left[L_{udd}^{S,RR} \right]_{11r2} \right. \\ &\quad - \frac{m_n}{2m_\Sigma} \left(\alpha \left[L_{udd}^{S,LR} \right]_{12r1} + \beta \left[L_{udd}^{S,RR} \right]_{12r1} \right) (D - F) \\ &\quad \left. + \frac{m_n}{6m_\Lambda} \left(\alpha \left[L_{udd}^{S,LR} \right]_{12r1} + \beta \left[L_{udd}^{S,RR} \right]_{12r1} + 2\alpha \left[L_{udd}^{S,LR} \right]_{11r2} + 2\beta \left[L_{udd}^{S,RR} \right]_{11r2} \right) (D + 3F) \right|^2 \end{aligned}$$

α, β
computed in the lattice

K-matrices

$$\Gamma(p \rightarrow \pi^+ \nu) = 2 \Gamma(n \rightarrow \pi^0 \nu)$$

$$\Gamma(n \rightarrow \pi^- e^+) = 3 \Gamma(p \rightarrow \pi^0 e^+)$$



BNC Lagrangian

$$\begin{aligned}
\mathcal{L}_0 \supset & \left(\frac{D-F}{f_\pi} \overline{\Sigma^+} \gamma^\mu \gamma_5 p - \frac{D+3F}{\sqrt{6}f_\pi} \overline{\Lambda^0} \gamma^\mu \gamma_5 n - \frac{D-F}{\sqrt{2}f_\pi} \overline{\Sigma^0} \gamma^\mu \gamma_5 n \right) \partial_\mu \bar{K}^0 \\
& + \left(\frac{D-F}{\sqrt{2}f_\pi} \overline{\Sigma^0} \gamma^\mu \gamma_5 p - \frac{D+3F}{\sqrt{6}f_\pi} \overline{\Lambda^0} \gamma^\mu \gamma_5 p + \frac{D-F}{f_\pi} \overline{\Sigma^-} \gamma^\mu \gamma_5 n \right) \partial_\mu K^- \\
& + \frac{3F-D}{2\sqrt{6}f_\pi} (\bar{p} \gamma^\mu \gamma_5 p + \bar{n} \gamma^\mu \gamma_5 n) \partial_\mu \eta \\
& + \frac{D+F}{f_\pi} \bar{p} \gamma^\mu \gamma_5 n \partial_\mu \pi^+ \\
& + \frac{D+F}{2\sqrt{2}f_\pi} (\bar{p} \gamma^\mu \gamma_5 p - \bar{n} \gamma^\mu \gamma_5 n) \partial_\mu \pi^0 + \text{h.c.} .
\end{aligned}$$

$$\begin{aligned}
\xi B \xi & \rightarrow L \xi B \xi R^\dagger \\
\xi B \xi^\dagger & \rightarrow L \xi B \xi^\dagger L^\dagger
\end{aligned}$$

$$\xi B \xi \sim (\mathbf{3}, \bar{\mathbf{3}}), \quad \xi^\dagger B \xi^\dagger \sim (\bar{\mathbf{3}}, \mathbf{3}), \quad \xi B \xi^\dagger \sim (\mathbf{8}, \mathbf{1}), \quad \xi^\dagger B \xi \sim (\mathbf{1}, \mathbf{8})$$

$$\alpha \cdot \nu \text{ tr}(\xi B \xi^\dagger P_{32}) = -(du)(d\nu) = [\mathcal{O}_{udd}]_{1111}^{S,LL}$$

$$\langle 0 | \epsilon^{abc} (\bar{u}_a^\dagger \bar{d}_b^\dagger) u_c | p^{(s)} \rangle = \alpha P_L u_p^{(s)}$$

$$\langle 0 | \epsilon^{abc} (u_a d_b) u_c | p^{(s)} \rangle = \beta P_L u_p^{(s)}$$

BNV Lagrangian

Name	LEFT	Flavour/B χ PT
$[\mathcal{O}_{udd}^{S,LL}]_{rstu}$	$(u_r d_s)(d_t \nu_u)$	$(\mathbf{8}, \mathbf{1})$
$[\mathcal{O}_{udd}^{S,LL}]_{111r}$	$(ud)(d\nu_r)$	$-\beta \overline{\nu_{Lr}^c} \text{tr}(\xi B \xi^\dagger P_{32}) \supset -\beta \overline{\nu_{Lr}^c} n - \frac{i\beta}{f_\pi} \overline{\nu_{Lr}^c} \left(\sqrt{\frac{3}{2}} n\eta - \frac{1}{\sqrt{2}} n\pi^0 + p\pi^- \right)$
$[\mathcal{O}_{udd}^{S,LL}]_{121r}$	$(us)(d\nu_r)$	$-\beta \overline{\nu_{Lr}^c} \text{tr}(\xi B \xi^\dagger \tilde{P}_{22}) \supset -\beta \overline{\nu_{Lr}^c} \left(-\frac{\Lambda^0}{\sqrt{6}} + \frac{\Sigma^0}{\sqrt{2}} \right) - \frac{i\beta}{f_\pi} \overline{\nu_{Lr}^c} n \bar{K}^0$
$[\mathcal{O}_{udd}^{S,LL}]_{112r}$	$(ud)(s\nu_r)$	$-\beta \overline{\nu_{Lr}^c} \text{tr}(\xi B \xi^\dagger P_{33}) \supset \beta \sqrt{\frac{2}{3}} \overline{\nu_{Lr}^c} \Lambda^0 - \frac{i\beta}{f_\pi} \overline{\nu_{Lr}^c} (n \bar{K}^0 + p K^-)$
$[\mathcal{O}_{duu}^{S,LL}]_{rstu}$	$(d_r u_s)(u_t e_u)$	$(\mathbf{8}, \mathbf{1})$
$[\mathcal{O}_{duu}^{S,LL}]_{111r}$	$(du)(u e_r)$	$-\beta \overline{e_{Lr}^c} \text{tr}(\xi B \xi^\dagger \tilde{P}_{31}) \supset \beta \overline{e_{Lr}^c} p + \frac{i\beta}{f_\pi} \overline{e_{Lr}^c} \left(\sqrt{\frac{3}{2}} p\eta + \frac{1}{\sqrt{2}} p\pi^0 + n\pi^+ \right)$
$[\mathcal{O}_{duu}^{S,LL}]_{211r}$	$(su)(u e_r)$	$-\beta \overline{e_{Lr}^c} \text{tr}(\xi B \xi^\dagger P_{21}) \supset -\beta \overline{e_{Lr}^c} \Sigma^+ + \frac{i\beta}{f_\pi} \overline{e_{Lr}^c} p \bar{K}^0$
$[\mathcal{O}_{uud}^{S,LR}]_{[rs]tu}$	$(u_r u_s)(\bar{d}_t^\dagger \bar{e}_u^\dagger)$	—
$[\mathcal{O}_{duu}^{S,LR}]_{rstu}$	$(d_r u_s)(\bar{u}_t^\dagger \bar{e}_u^\dagger)$	$(\bar{\mathbf{3}}, \mathbf{3})$
$[\mathcal{O}_{duu}^{S,LR}]_{111r}$	$(du)(\bar{u}^\dagger \bar{e}_r^\dagger)$	$\alpha \overline{e_{Rr}^c} \text{tr}(\xi^\dagger B \xi^\dagger \tilde{P}_{31}) \supset -\alpha \overline{e_{Rr}^c} p + \frac{i\alpha}{f_\pi} \overline{e_{Rr}^c} \left(-\frac{1}{\sqrt{6}} p\eta + \frac{1}{\sqrt{2}} p\pi^0 + n\pi^+ \right)$
$[\mathcal{O}_{duu}^{S,LR}]_{211r}$	$(su)(\bar{u}^\dagger \bar{e}_r^\dagger)$	$\alpha \overline{e_{Rr}^c} \text{tr}(\xi^\dagger B \xi^\dagger P_{21}) \supset \alpha \overline{e_{Rr}^c} \Sigma^+ - \frac{i\alpha}{f_\pi} \overline{e_{Rr}^c} p \bar{K}^0$
$[\mathcal{O}_{uud}^{S,RL}]_{[rs]tu}$	$(\bar{u}_r^\dagger \bar{u}_s^\dagger)(d_t e_u)$	—
$[\mathcal{O}_{duu}^{S,RL}]_{rstu}$	$(\bar{d}_r^\dagger \bar{u}_s^\dagger)(u_t e_u)$	$(\mathbf{3}, \bar{\mathbf{3}})$
$[\mathcal{O}_{duu}^{S,RL}]_{111r}$	$(\bar{d}^\dagger \bar{u}^\dagger)(u e_r)$	$-\alpha \overline{e_{Lr}^c} \text{tr}(\xi B \xi \tilde{P}_{31}) \supset \alpha \overline{e_{Lr}^c} p + \frac{i\alpha}{f_\pi} \overline{e_{Lr}^c} \left(-\frac{1}{\sqrt{6}} p\eta + \frac{1}{\sqrt{2}} p\pi^0 + n\pi^+ \right)$
$[\mathcal{O}_{duu}^{S,RL}]_{211r}$	$(\bar{s}^\dagger \bar{u}^\dagger)(u e_r)$	$-\alpha \overline{e_{Lr}^c} \text{tr}(\xi B \xi P_{21}) \supset -\alpha \overline{e_{Lr}^c} \Sigma^+ - \frac{i\alpha}{f_\pi} \overline{e_{Lr}^c} p \bar{K}^0$
$[\mathcal{O}_{dud}^{S,RL}]_{rstu}$	$(\bar{d}_r^\dagger \bar{u}_s^\dagger)(d_t \nu_u)$	$(\mathbf{3}, \bar{\mathbf{3}})$
$[\mathcal{O}_{dud}^{S,RL}]_{111r}$	$(\bar{d}^\dagger \bar{u}^\dagger)(d \nu_r)$	$\alpha \overline{\nu_{Lr}^c} \text{tr}(\xi B \xi \tilde{P}_{32}) \supset -\alpha \overline{\nu_{Lr}^c} n + \frac{i\alpha}{f_\pi} \overline{\nu_{Lr}^c} \left(\frac{1}{\sqrt{6}} n\eta + \frac{1}{\sqrt{2}} n\pi^0 - p\pi^- \right)$
$[\mathcal{O}_{dud}^{S,RL}]_{211r}$	$(\bar{s}^\dagger \bar{u}^\dagger)(d \nu_r)$	$\alpha \overline{\nu_{Lr}^c} \text{tr}(\xi B \xi P_{22}) \supset \alpha \overline{\nu_{Lr}^c} \left(\frac{\Lambda^0}{\sqrt{6}} - \frac{\Sigma^0}{\sqrt{2}} \right) + \frac{i\alpha}{f_\pi} \overline{\nu_{Lr}^c} n \bar{K}^0$
$[\mathcal{O}_{dud}^{S,RL}]_{112r}$	$(\bar{d}^\dagger \bar{u}^\dagger)(s \nu_r)$	$\alpha \overline{\nu_{Lr}^c} \text{tr}(\xi B \xi \tilde{P}_{33}) \supset \alpha \overline{\nu_{Lr}^c} \sqrt{\frac{2}{3}} \Lambda^0 - \frac{i\alpha}{f_\pi} \overline{\nu_{Lr}^c} (n \bar{K}^0 + p K^-)$
$[\mathcal{O}_{dud}^{S,RL}]_{212r}$	$(\bar{s}^\dagger \bar{u}^\dagger)(s \nu_r)$	$\alpha \overline{\nu_{Lr}^c} \text{tr}(\xi B \xi P_{23}) \supset \alpha \overline{\nu_{Lr}^c} \Xi^0$
$[\mathcal{O}_{ddu}^{S,RL}]_{[rs]tu}$	$(\bar{d}_r^\dagger \bar{d}_s^\dagger)(u_t \nu_u)$	$(\mathbf{3}, \bar{\mathbf{3}})$
$[\mathcal{O}_{ddu}^{S,RL}]_{[12]1r}$	$(\bar{d}^\dagger \bar{s}^\dagger)(u \nu_r)$	$-\alpha \overline{\nu_{Lr}^c} \text{tr}(\xi B \xi P_{11}) \supset \alpha \overline{\nu_{Lr}^c} \left(\frac{\Lambda^0}{\sqrt{6}} + \frac{\Sigma^0}{\sqrt{2}} \right) - \frac{i\alpha}{f_\pi} \overline{\nu_{Lr}^c} p K^-$
$[\mathcal{O}_{duu}^{S,RR}]_{rstu}$	$(\bar{d}_r^\dagger \bar{u}_s^\dagger)(\bar{u}_t^\dagger \bar{e}_u^\dagger)$	$(\mathbf{1}, \mathbf{8})$
$[\mathcal{O}_{duu}^{S,RR}]_{111r}$	$(\bar{d}^\dagger \bar{u}^\dagger)(\bar{u}^\dagger \bar{e}_r^\dagger)$	$\beta \overline{e_{Rr}^c} \text{tr}(\xi^\dagger B \xi^\dagger \tilde{P}_{31}) \supset -\beta \overline{e_{Rr}^c} p + \frac{i\beta}{f_\pi} \overline{e_{Rr}^c} \left(\sqrt{\frac{3}{2}} p\eta + \frac{1}{\sqrt{2}} p\pi^0 + n\pi^+ \right)$
$[\mathcal{O}_{duu}^{S,RR}]_{211r}$	$(\bar{s}^\dagger \bar{u}^\dagger)(\bar{u}^\dagger \bar{e}_r^\dagger)$	$\beta \overline{e_{Rr}^c} \text{tr}(\xi^\dagger B \xi^\dagger P_{21}) \supset \beta \overline{e_{Rr}^c} \Sigma^+ + \frac{i\beta}{f_\pi} \overline{e_{Rr}^c} p \bar{K}^0$

Name	LEFT	Flavour/B χ PT
$[\mathcal{O}_{udd}^{S,LL}]_{rstu}$	$(u_r d_s)(d_t \nu_u)$	$(\mathbf{8}, \mathbf{1})$
$[\mathcal{O}_{udd}^{S,LL}]_{111r}$	$(ud)(d\nu_r)$	$-\beta \overline{\nu_{Lr}^c} \text{tr}(\xi B \xi^\dagger P_{32}) \supset -\beta \overline{\nu_{Lr}^c} n - \frac{i\beta}{f_\pi} \overline{\nu_{Lr}^c} \left(\sqrt{\frac{3}{2}} n\eta - \frac{1}{\sqrt{2}} n\pi^0 + p\pi^- \right)$
$[\mathcal{O}_{udd}^{S,LL}]_{121r}$	$(us)(d\nu_r)$	$-\beta \overline{\nu_{Lr}^c} \text{tr}(\xi B \xi^\dagger \tilde{P}_{22}) \supset -\beta \overline{\nu_{Lr}^c} \left(-\frac{\Lambda^0}{\sqrt{6}} + \frac{\Sigma^0}{\sqrt{2}} \right) - \frac{i\beta}{f_\pi} \overline{\nu_{Lr}^c} n \bar{K}^0$
$[\mathcal{O}_{udd}^{S,LL}]_{112r}$	$(ud)(s\nu_r)$	$-\beta \overline{\nu_{Lr}^c} \text{tr}(\xi B \xi^\dagger P_{33}) \supset \beta \sqrt{\frac{2}{3}} \overline{\nu_{Lr}^c} \Lambda^0 - \frac{i\beta}{f_\pi} \overline{\nu_{Lr}^c} (n \bar{K}^0 + p K^-)$
$[\mathcal{O}_{duu}^{S,LL}]_{rstu}$	$(d_r u_s)(u_t e_u)$	$(\mathbf{8}, \mathbf{1})$
$[\mathcal{O}_{duu}^{S,LL}]_{111r}$	$(du)(u e_r)$	$-\beta \overline{e_{Lr}^c} \text{tr}(\xi B \xi^\dagger \tilde{P}_{31}) \supset \beta \overline{e_{Lr}^c} p + \frac{i\beta}{f_\pi} \overline{e_{Lr}^c} \left(\sqrt{\frac{3}{2}} p\eta + \frac{1}{\sqrt{2}} p\pi^0 + n\pi^+ \right)$
$[\mathcal{O}_{duu}^{S,LL}]_{211r}$	$(su)(u e_r)$	$-\beta \overline{e_{Lr}^c} \text{tr}(\xi B \xi^\dagger P_{21}) \supset -\beta \overline{e_{Lr}^c} \Sigma^+ + \frac{i\beta}{f_\pi} \overline{e_{Lr}^c} p \bar{K}^0$
$[\mathcal{O}_{uud}^{S,LR}]_{[rs]tu}$	$(u_r u_s)(\bar{d}_t^\dagger \bar{e}_u^\dagger)$	—
$[\mathcal{O}_{duu}^{S,LR}]_{rstu}$	$(d_r u_s)(\bar{u}_t^\dagger \bar{e}_u^\dagger)$	$(\bar{\mathbf{3}}, \mathbf{3})$
$[\mathcal{O}_{duu}^{S,LR}]_{111r}$	$(du)(\bar{u}^\dagger \bar{e}_r^\dagger)$	$\alpha \overline{e_{Rr}^c} \text{tr}(\xi^\dagger B \xi^\dagger \tilde{P}_{31}) \supset -\alpha \overline{e_{Rr}^c} p + \frac{i\alpha}{f_\pi} \overline{e_{Rr}^c} \left(-\frac{1}{\sqrt{6}} p\eta + \frac{1}{\sqrt{2}} p\pi^0 + n\pi^+ \right)$
$[\mathcal{O}_{duu}^{S,LR}]_{211r}$	$(su)(\bar{u}^\dagger \bar{e}_r^\dagger)$	$\alpha \overline{e_{Rr}^c} \text{tr}(\xi^\dagger B \xi^\dagger P_{21}) \supset \alpha \overline{e_{Rr}^c} \Sigma^+ - \frac{i\alpha}{f_\pi} \overline{e_{Rr}^c} p \bar{K}^0$
$[\mathcal{O}_{uud}^{S,RL}]_{[rs]tu}$	$(\bar{u}_r^\dagger \bar{u}_s^\dagger)(d_t e_u)$	—
$[\mathcal{O}_{duu}^{S,RL}]_{rstu}$	$(\bar{d}_r^\dagger \bar{u}_s^\dagger)(u_t e_u)$	$(\mathbf{3}, \bar{\mathbf{3}})$
$[\mathcal{O}_{duu}^{S,RL}]_{111r}$	$(\bar{d}^\dagger \bar{u}^\dagger)(u e_r)$	$-\alpha \overline{e_{Lr}^c} \text{tr}(\xi B \xi^\dagger \tilde{P}_{31}) \supset \alpha \overline{e_{Lr}^c} p + \frac{i\alpha}{f_\pi} \overline{e_{Lr}^c} \left(-\frac{1}{\sqrt{6}} p\eta + \frac{1}{\sqrt{2}} p\pi^0 + n\pi^+ \right)$
$[\mathcal{O}_{duu}^{S,RL}]_{211r}$	$(\bar{s}^\dagger \bar{u}^\dagger)(u e_r)$	$-\alpha \overline{e_{Lr}^c} \text{tr}(\xi B \xi^\dagger P_{21}) \supset -\alpha \overline{e_{Lr}^c} \Sigma^+ - \frac{i\alpha}{f_\pi} \overline{e_{Lr}^c} p \bar{K}^0$

Nucleon decay channels

$$\Gamma(N \rightarrow M\ell_\alpha) \quad \left\{ \begin{array}{l} n \rightarrow \eta^0 \nu \\ n \rightarrow \pi^0 \nu \\ p \rightarrow \pi^+ \nu \\ n \rightarrow \pi^- e^+ \\ p \rightarrow \eta^0 e^+ \\ p \rightarrow \pi^0 e^+ \\ p \rightarrow K^0 e^+ \\ n \rightarrow K^0 \nu \\ p \rightarrow K^+ \nu \end{array} \right. \quad \Delta(B - L) = 0$$

$$\Gamma(N \rightarrow M\ell_\alpha) \quad \left\{ \begin{array}{l} n \rightarrow \eta^0 \nu \\ n \rightarrow \pi^0 \nu \\ p \rightarrow \pi^+ \nu \\ n \rightarrow K^0 \nu \\ p \rightarrow K^+ \nu \\ n \rightarrow K^+ e^- \end{array} \right. \quad |\Delta(B - L)| = 2$$

- All 2-body PS decays except for $p \rightarrow \bar{K}^0 e^+$ $n \rightarrow \bar{K}^0 \nu$ $n \rightarrow K^- e^+$ $n \rightarrow \pi^+ e^-$

(No $B\chi PT$ formalism developed for PD into vector mesons, e.g. $p \rightarrow \rho^0 e^+$)

SMEFT

- RGEs dominated by the SM gauge couplings → enhancement
- QCD contributions universal and dominant, BUT electroweak contributions are relevant for $\mathcal{O}_{qqql,1111}$
- top-loop contributions universal and suppressive for $d = 7$ WCs
- Operator Mixing subdominant for proton decay

$$c(m_W) \sim (2 - 4) c(10^{15} \text{ GeV})$$

- RGEs for $d = 6$ SMEFT Manohar et. al. [2014]
- RGEs for $d = 7$ SMEFT Yi Liao et. al. [2016]

Phenomenological matrices

$$\Gamma_{(i)}^{\Delta(B-L)=0} \equiv 10^{-4} c_j^* \kappa_{(i)}^{jk} c_k \frac{m_p^5}{\Lambda^4} \quad \text{for} \quad i = p \rightarrow \pi^0 e^+, p \rightarrow K^+ \nu \dots \quad (9 \text{ matrices})$$

$$\Gamma_{(i)}^{|\Delta(B-L)|=2} \equiv 10^{-4} c_j^* \kappa_{(i)}^{jk} c_k \frac{m_p^7}{\Lambda^6} \quad \text{for} \quad i = p \rightarrow K^+ \nu, n \rightarrow K^+ e^- \dots \quad (6 \text{ matrices})$$

(Numerical κ -matrices available online)

