

Neutrinoless double beta decay searches with the LEGEND experiment

Thursday 6 June 2024 14:50 (20 minutes)

Neutrinoless double beta ($0\nu\beta\beta$) decay is an ultra-rare process which could take place only if neutrinos were Majorana fermions, namely if neutrinos were their own antiparticle: if observed, this decay would shed light on neutrinos' nature and would be an unambiguous evidence for the existence of some Beyond Standard Model Physics, as it entails a violation of the lepton number by two units. Also, from the study of this decay it would be possible to give an explanation of the matter-antimatter asymmetry observed in the Universe and to extract information about neutrino masses.

The LEGEND Experiment is designed to search for the neutrinoless double beta decay of ^{76}Ge employing active ^{76}Ge -enriched HPGe detectors; these detectors are operated bare in Liquid Argon (LAr), serving both as a refrigerant and as a veto for background events; the LAr cryostat itself is immersed in a large volume of water, serving as muon veto.

The first phase of the experiment, LEGEND-200, started taking data in March 2023 at Laboratori Nazionali del Gran Sasso (LNGS) in Italy and is now running in a stable physics data taking regime. With an exposure of 1 ton yr and a target background index of $2 \cdot 10^{-4}$ cts/(keV kg yr) at $Q_{\beta\beta} = 2039$ keV, LEGEND-200 is planned to reach a 3σ discovery sensitivity of 1027 yr. The second phase, LEGEND-1000, will operate 1000 kg of Germanium and is planned to achieve a 3σ discovery sensitivity beyond 1028 yr with its target background index of $1 \cdot 10^{-5}$ cts/(keV kg yr) at $Q_{\beta\beta}$. LEGEND-1000 sensitivity will allow to cover the full inverted mass ordering region.

In this contribution LEGEND's physics program will be presented, with a focus on the current status and results of the ongoing experimental campaign.

Primary author: SALEH, Giovanna

Presenter: SALEH, Giovanna

Session Classification: Parallel Session PII.4