

# Plasma Wakefield Acceleration

**AWAKE Experiment at CERN**

Marlene Turner

# Outline

- What are **plasma wakefields** and why are they interesting?
- How to accelerate **charged particles** using plasma wakefields?
  - Underlying physics concepts, state-of-the-art results
- What is the **AWAKE experiment**, and why is it important?
- The AWAKE experimental setup
- Latest AWAKE results
  - Ideas and plans for the future


# AWAKE

## Advanced Proton-Driven Plasma Wakefield Acceleration Experiment



- Plasma ?
- Proton-driven ?
- Wakefield acceleration ?
- Acceleration ?

# Charged Particle Acceleration



- Acceleration of charged particles requires an electric field
- Charged particle will accelerate as long as it experiences the field

1 TV =  $10^{12}$  V

1 GV =  $10^9$  V

1 MV =  $10^6$  V

1 kV =  $10^3$  V


1.5V with battery length of ~3 cm  $\rightarrow$  50 V/m

To reach 1 TeV  $\rightarrow$  ~20 000 million km


Distance Earth-Sun ~ 152 million km



# Charged Particle Acceleration




- Acceleration of charged particles requires an electric field
- Charged particle will accelerate as long as it experiences the field



- Even better:
  - Field travels together with the beam

# Definition of Plasma and Plasma Wakefield

## Plasma




### Plasma: ionised gas (4<sup>th</sup> state of matter)

- **Quasi-neutrality:** the overall charge of a plasma is about zero.
- **Collective effects:** Charged particles must be close enough together that each particle influences many nearby charged particles.
- **Electrostatic interactions dominate** over collisions or ordinary gas kinetics.

# Definition of Plasma and Plasma Wakefield


## Plasma



## Plasma: ionised gas (4<sup>th</sup> state of matter)


- **Quasi-neutrality:** the overall charge of a plasma is about zero.
- **Collective effects:** Charged particles must be close enough together that each particle influences many nearby charged particles.
- **Electrostatic interactions dominate** over collisions or ordinary gas kinetics.

## Plasma Wakefields



## Plasma Wakefields:

- are the **fields** created/sustained by collective motion of plasma particles.




# Why use Plasmas for Charged Particle Acceleration?

**Conventional technology:**  
metallic radiofrequency (RF) cavities



LHC cavities

**New concept:**  
plasma wakefields acceleration  
→ transient structures in plasma






# Accelerating Gradient

## RF cavities


Limited to  $\sim 100$  MV/m due to electric breakdowns (ionization).



## Plasma Wakefields

Plasma is already ionized or “broken-down” and can sustain electric fields  $\sim 100$  GV/m.


$$eE_{max} = 1 \left[ \frac{eV}{cm} \right] \cdot n^{1/2} [cm^{-3}]$$



# Accelerating Gradient

## RF cavities

Limited to  $\sim 100$  MV/m due to electric breakdowns (ionization).




→ Plasma wakefields can sustain order of magnitude higher fields

## Plasma Wakefields

Plasma is already ionized or “broken-down” and can sustain electric fields  $\sim 100$  GV/m.

$$eE_{max} = 1 \left[ \frac{eV}{cm} \right] \cdot n^{1/2} [cm^{-3}]$$




Structure exists only for a very short amount of time!

# Circular and Linear Accelerators

## Circular accelerators

Image from P. Arpaia et al.,  
NIMA 985, 164652 (2021)




- Advantage: beam passes accelerating section many times
- Disadvantage: synchrotron radiation losses

LHC tunnel:  
 $p+p \rightarrow 14 \text{ TeV}$   
 $e+e^- \rightarrow 209 \text{ GeV}$

- Beam passes acceleration section multiple times.
- Max. energy (E) limited by synchrotron radiation losses  
 $\propto E^4/(r^2m^4)$

# Synchrotron Radiation



Synchrotron radiation is caused by leaving part of fields behind when the beam moves along the curve.

$$\propto E^4 / (r^2 m^4)$$

Particle energy      Bending radius      Mass of the particle


→ Needs to be taken into account when accelerated charged particles are deflected in the radial direction.



# Circular and Linear Accelerators


## Circular accelerators

Image from P. Arpaia et al.,  
NIMA 985, 164652 (2021)



- Beam passes acceleration section multiple times.
- Max. energy (E) limited by synchrotron radiation losses  
 $\propto E^4/(r^2m^4)$

## Linear accelerators



- Beam passes acceleration section multiple times.
- Negligible synchrotron radiation losses
- Accelerator length and accelerating gradient define final beam energy.

e.g. to accelerate electrons to 1 TeV ( $10^{12}$  eV):  
 100 MeV/m x 10000 m or  
 100 GeV/m x 10 m


# Let Us Repeat...

- Plasma wakefields allow to accelerate charged particles with  $\sim 1\text{-}100$  GeV/m
- High gradients are important when using linear accelerators (e.g. for light particles) to minimize synchrotron radiation losses
  - For linear accelerators, their length defines the final beam energy

# How to Create Plasma Wakefields?

Two ingredients

Plasma  
(Medium)



Relativistic charged particle beam or  
Short intense laser pulse  
(Energy source)









Image from  
<https://revbalance.com/improving-balance-for-wakesurfing/>




# Energy Source: The Driver





- Relativistic charged particle bunches or laser pulses
  - → Relativistic charged particle bunches carry almost purely **transverse electric** fields
- What we need → longitudinal electric field to accelerate charged particles

## Trick:

- Use plasma to convert the transverse electric field of the proton bunch into a longitudinal electric field in the plasma.
- The more energy is available, the longer (distance-wise) these plasma wakefields can be sustained




# How to Drive a Plasma Wave




## Important to understand

- Plasma electron motion is mostly **transverse**
- Electrons do not move significantly longitudinally
- Rb ions are heavy and do not move significantly on the timescale of the electrons

# How to Drive a Plasma Wave




Charge separation  $\rightarrow$  electric field  
(longitudinal and transverse)



Where should we place an  
electron bunch to be  
accelerated?



# Plasma Wakefields



- ← Accelerating for negatively charged particles
- Decelerating for negatively charged particles
- ↓ Focusing for negatively charged particles
- ↑ Defocusing for negatively charged particles

# Plasma Acts As a Transformer

Driver deposits energy, witness gains energy



Acceleration distance typically limited by either



**Depletion:** The drive bunch/pulse running out of energy. Solution: couple in a new driver or use a more energetic driver



**Dephasing:** The witness bunch outruns the driver. Solution: couple in a new driver or shape the plasma density




**Diffraction:** Drive beam evolution. Solution: Guiding or use a more rigid driver

# Let Us Repeat...

- Plasma wakefields require: plasma, energy source (driver)
  - Place a particle beam (witness) to be accelerated
- Plasma acts as a transformer
  - Drive beam energy is transferred to the witness bunch
- What limits the energy gain:
  - Depletion: Driver runs out of energy
  - Dephasing: Accelerating bunch outruns the driver
  - Diffraction: Driver no longer intense enough


# State-of-the-Art Results





# State-of-the-Art Results


## BELLA (Berkeley, California)



Petawatt Laser Guiding and Electron Beam Acceleration to 8 GeV in a Laser-Heated Capillary Discharge Waveguide

A. J. Gonsalves et al., Phys. Rev. Lett. 122, 084801









# State-of-the-Art Results

## SLAC (Stanford, California)




**42 GeV energy gain  
in 85 cm of  
plasma**



Blumenfeld, I., Clayton, C., Decker, F.J. *et al.* Energy doubling of 42 GeV electrons in a metre-scale plasma wakefield accelerator. *Nature* **445**, 741–744 (2007).

# First Demonstration of a Free Electron Laser Driven by a Plasma Wakefield Accelerator



Free-electron lasing at 27 nanometres based on a laser wakefield accelerator  
Wentao Wang et al., *Nature* volume 595, pages 516–520 (2021)


# Let Us Repeat...

- Plasma wakefields acceleration has been demonstrated experimentally:
  - ~8 GeV in 20 cm of plasma
  - 42 GeV in 85 cm of plasma
  - First laser-plasma wakefield driven free electron laser

# The AWAKE Experiment @CERN

# Plasma Wakefield Physics @ CERN

The CERN accelerator complex  
*Complexe des accélérateurs du CERN*



▶  $H^-$  (hydrogen anions) ▶ p (protons) ▶ ions ▶ RIBs (Radioactive Ion Beams) ▶ n (neutrons) ▶  $\bar{p}$  (antiprotons) ▶  $e^-$  (electrons) ▶  $\mu$  (muons)

LHC - Large Hadron Collider // SPS - Super Proton Synchrotron // PS - Proton Synchrotron // AD - Antiproton Decelerator // CLEAR - CERN Linear Electron Accelerator for Research // AWAKE - Advanced WAKEfield Experiment // ISOLDE - Isotope Separator OnLine // REX/HIE-ISOLDE - Radioactive Experiment/High Intensity and Energy ISOLDE // MEDICIS // LEIR - Low Energy Ion Ring // LINAC - LINear ACcelerator // n\_TOF - Neutrons Time Of Flight // HiRadMat - High-Radiation to Materials // Neutrino Platform

- CERN has very high energetic proton bunches available.
- Idea: use energy stored in the proton bunches to accelerate lighter particles e.g. electrons

19 kJ  
400 GeV

→ however, they are too long to excite wakefields



# AWAKE Requires Microbunching of p<sup>+</sup> Bunch



To effectively excite wakefields:

- The drive bunch length has to be on the order of the plasma wavelength

**For AWAKE → mm-scale bunch length**

CERN SPS proton bunch length is ~6 cm


⇒ Plasma process: Self-Modulation Instability (SMI), can be seeded (SSM)




- Wakefields driven resonantly to large amplitude
- Self-modulation necessary to drive ~GV/m accelerating fields in  $10^{14} \text{ cm}^{-3}$  density plasma



# Self-Modulation in Plasma Wakefields




# Simulation Result




# Self-Modulation Diagnostics

## Streak camera measurement



Foil emits waves up to the plasma wavelength of the foil including:


- radiation in the optical range (OTR).
- Coherent radiation (CTR) for wavelengths bigger than the structure of the micro-bunches






# Self-Modulation Measurement Results

Plasma off:



Plasma on:



Front

$\sim \sigma_z / c \sim 200 \text{ ps}$

Defocused p<sup>+</sup>


P. Muggli  
F. Batsch





**Shortly after we  
have observed  
the Seeded-Self  
Modulation for the  
first time!**

# Experimental Realization @CERN → AWAKE Experiment



From a concept and an idea to reality !



# AWAKE Components

## Plasma

- Laser
- Rubidium vapor

## Drive Bunch


- Proton beam (400 GeV/c)

## Witness Bunch


- Electron beam (10-20 MeV)

## Diagnostics:

- Proton
- Laser
- Electron



# Protons Delivered by CERN SPS



- Proton bunch momentum: 400 GeV/c
- $3 \times 10^{11}$  protons/bunch
- Bunch length:  $\sigma_z = \sim 10$  cm
- Radial bunch size at plasma entrance:  $\sigma_r = 0.2$  mm





# The AWAKE Plasma

**Rubidium vapour** cell.

The laser **ionizes** the outermost electron of each rubidium atom.

Desired **plasma density**:  $\sim 1-10 \times 10^{14}$  electrons/cm<sup>3</sup>.









# Accelerated Electron Energy Measurement




- Electrons will be injected with an energy around 10-20 MeV.
- Accelerated electrons are sent through a **dipole magnet** and deposit energy on a scintillating screen which is imaged by a camera.




8.5 ton, 1.2 T, 1.3 Tm, L=1.6 m, W=1.3 m




Typical final energy distribution of the accelerated electron beam after 10 m plasma:



# First Electron Acceleration




No electrons accelerated.



Accelerated electrons.



# Electron Acceleration Results



AWAKE Collaboration, *Nature* volume 561, pages 363–367 (2018)




**Shortly after we  
have observed  
electron  
acceleration for  
the first time!**

# Let Us Repeat...

- To realize the AWAKE experiment at CERN, we need:
  - Plasma (vapor source + laser)
  - Proton bunch (wakefield driver)
  - Electron bunch (witness – to be accelerated)
  
- Diagnostics are key to a successful measurement
  - AWAKE diagnostics include:
    - Screens + Streak camera (to know beam positions and verify that SSM was successful)
    - Electron spectrometer (energy of the accelerated witness bunch)




# Most Recently: New Plasma Source Technology






Xe






Ar






He




# Motion of Ions Leads to Decoherence of Plasma Electron Motion






# Experimental Observation of Motion of Ions


Simulation



Experiment





# Now in AWAKE: New Plasma Source




- Allows to adjust the plasma density along the 10 m
- More stable SSM, → higher wakefield amplitudes







# AWAKE until ~2030






# First AWAKE Particle Physics Applications

## Example I: Dark Matter Experiment

These experiments use the collisions of an electron beam with a fixed-target or a dump to generate the dark photon via Bremsstrahlung (electron and proton beams) or meson production.

The products of the collisions are mostly absorbed in the dump and the dark photon is searched for as a displaced vertex with two opposite charged tracks in the decay volume of the experiment.



From Caldwell: <https://arxiv.org/pdf/1812.11164.pdf>


# First AWAKE Particle Physics Applications

## Example II: Electron-Proton Collisions

➤ Collide:


- 50 GeV electrons with 7 TeV LHC protons
- ~TeV electrons with 7 TeV LHC protons

### Plasma-based collider design



### Physics cases:

- Study of the sub-structure and spin structure of the proton and photon
- Determine if partons are fundamental point-like objects
- Clarifying the underlying physics leading to the energy dependence of cross sections
- Leptoquark production: hypothetical particles that would interact with quarks and leptons



Diagrams of neutral-current (top) and charged-current (bottom) deep-inelastic electron–proton scattering processes. Image credit: DESY.

Caldwell, A., Wing, M. VHEeP: a very high energy electron–proton collider. *Eur. Phys. J. C* **76**, 463 (2016). <https://doi.org/10.1140/epjc/s10052-016-4316-1>

# Summary and Conclusions

- **Plasma wakefield acceleration** is a novel technique to accelerate charged particles
  - Advantage: Very **high accelerating gradient**, compact accelerators
  - Proof of principle acceleration has been demonstrated
  - **Next step**: aim for high beam quality in long plasmas → First applications
- AWAKE is an accelerator R&D experiment at CERN:
  - Only proton-driven wakefield acceleration experiment worldwide
  - The experiment opens a pathway towards particle physics applications
- **Final Goal**: Design high quality & high energy electron accelerator based on acquired knowledge.





[home.cern](http://home.cern)