

Quantum Collider probes of the fermionic Higgs Portal

Thursday 29 February 2024 19:00 (10 minutes)

Solutions to the dark matter puzzle and the hierarchy problem can include a BSM fermion ψ that is a SM singlet and couples to the visible sector via a Higgs portal interaction: $\mathcal{L}_{H\psi} = c_\psi/f |H|^2 \bar{\psi}\psi$. Despite its potential of solving these tensions with the SM being well-established, the study of $\mathcal{L}_{H\psi}$ at hadron colliders remains challenging as the Higgs boson is the only portal between the ψ -particles and the SM sector. We study the interaction $\mathcal{L}_{H\psi}$ with several off-shell Higgs probes, namely the contribution of ψ to the Higgs self-energy in $pp \rightarrow ZZ \rightarrow 4\ell$ production, Di-Higgs production, and the direct production of ψ -pairs in off-shell Higgs decays. Finally, we present the combination of these three analyses providing complementary results and give an outlook on the collider reach of HL-LHC and other beyond-LHC pp -colliders in constraining the $\mathcal{L}_{H\psi}$ interaction.

Primary author: SCHMID, Konstantin (University and INFN Padova)

Presenter: SCHMID, Konstantin (University and INFN Padova)

Session Classification: Short Talks Session