



Contribution ID: 175

Type: Oral

Heavy Quark Diffusion Coefficient from Lattice QCD

We present new lattice results for the heavy quark diffusion coefficient in 2+1 flavor QCD in the temperature range from 163 MeV up to few GeV. Compared to previous lattice calculations with unphysical light quark masses, we consider near-physical values and a much wider temperature range. Our results for the spatial heavy diffusion coefficient near the crossover temperature are considerably smaller than the estimates obtained by comparing phenomenological models with experimental data and by T-matrix calculations, and are close to the AdS/CFT limit. At high temperatures, however, the spatial heavy diffusion coefficient increases and approaches the NLO weak coupling prediction within the estimated errors. We also find that the dependence of the spatial heavy quark diffusion coefficient on the heavy quark mass M_Q on the lattice is weaker compared to the model calculations.

Category

Theory

Collaboration (if applicable)

HotQCD Collaboration

Primary authors: SHU, Hai-Tao (Central China Normal University); Dr WEBER, Johannes Heinrich (Humboldt University of Berlin); DASILVA GOLAN, Jorge Luis (Brookhaven National Laboratory); KACZMAREK, Olaf; PETRECZKY, Peter; LARSEN, Rasmus (University of Stavanger); MUKHERJEE, Swagato

Presenter: DASILVA GOLAN, Jorge Luis (Brookhaven National Laboratory)

Track Classification: Heavy flavor & quarkonia