Quark Matter 2025



Contribution ID: 636

Type: Oral

## Finite density lattice QCD without extrapolations

Finite density lattice QCD usually relies on extrapolations in baryon chemical potential ( $\mu_B$ ), be it Taylor expansion, T' expansion or analytical continuation. However, their range of validity is difficult to control. In the canonical formulation, the baryon density is the parameter of the system, not  $\mu_B$ . Here we demonstrate that we can access finite density QCD in the canonical formulation with physical quark masses along the strangeness neutral line. We present first results with both the strangeness ( $n_S$ ) and baryon ( $n_B$ ) densities as parameters. Specifically, we compute the QCD pressure and chemical potentials as functions of  $n_B$  and  $n_S$  and construct an equation of state. Our computations rely on high-statistics simulations with 2+1 4HEX-staggered fermions.

## Category

Theory

## **Collaboration (if applicable)**

**Primary authors:** BORSANYI, Szabolcs Istvan; FODOR, Zoltan; GUENTHER, Jana N. (University of Wuppertal); PAROTTO, Paolo; PASZTOR, Attila (Eötvös University); PIRELLI, Ludovica (Bergische Universität Wuppertal); Dr WONG, Chik Him

Presenter: PIRELLI, Ludovica (Bergische Universität Wuppertal)

Track Classification: New theoretical developments