



Contribution ID: 54

Type: Poster

## The effect of extra dimensions on compact star observables

Kaluza and Klein proposed a theory with a compactified extra dimension, which may appear in high-energy reactions or in the presence of superdense matter. In this work, I show how astrophysical observables will be modified in the presence of extra compactified dimensions.

The interior of a compact star is modeled as a multidimensional interacting degenerate Fermi gas, embedded in a static, spherically symmetric spacetime with extra compactified spatial dimensions. The equation of state of this extreme medium is given and compared to the standard models of superdense matter. The modification of the mass-radius relation of compact stars is calculated and compared to realistic star models and astrophysical observation data. The interaction strength has been determined for this extraordinary matter. Constraints on the size of the extra dimension have been estimated based on pulsar measurements [1,2].

[1] A. Horváth, E. Forgács-Dajka, G.G. Barnaföldi: “Application of Kaluza-Klein Theory in Modeling Compact Stars: Exploring Extra Dimensions”, Submitted to MNRAS, arXiv:2408.16497

[2] A. Horváth, E. Forgács-Dajka, G.G. Barnaföldi: “The effect of multiple extra dimensions on the maximal mass of compact stars in Kaluza-Klein space-time”, Submitted to International Journal of Modern Physics A, arXiv:2410.10920

### Category

Theory

### Collaboration (if applicable)

**Authors:** HORVATH, Anna; FORGÁCS-DAJKA, Emese (Eötvös Loránd University); Dr BARNAFOLDI, Gergely Gabor (HUN-REN Wigner Research Centre for Physics (HU))

**Presenter:** HORVATH, Anna

**Session Classification:** Poster session 1

**Track Classification:** QCD matter in astrophysics