



Contribution ID: 253

Type: Poster

Preliminary results on the transverse flow of strange hadrons emitted from Ag+Ag collisions at beam energy of 1.58 AGeV measured with HADES

The High Acceptance Di-Electron Spectrometer (HADES) [1] installed at the GSI institute in Darmstadt registers products of heavy ion collisions at energies of a few GeV per nucleon. The nuclear matter produced in such collisions reaches extreme densities and temperatures [2, 3, 4], comparable to those found in neutron stars, especially during their mergers [5]. By measuring the kinematic distributions of emitted particles and comparing them to transport model calculations –many aspects of the nuclear matter can be understood.

Particles containing the (anti)strange quark are of particular interest, as they are produced sparsely in this energy regime and act as good probes of their interaction with the medium. For charged K mesons it is predicted that due to their interaction in nuclear matter, their effective mass and decay constant should change [6]. This phenomenon was investigated by other experiments, but so far without a definitive conclusion [7]. Another interesting particle species is the Λ baryon, due to its predicted role in the so-called hyperon puzzle [8], where its interaction potential with nucleonic matter is an important piece of the “puzzle”.

It is predicted that the transverse flow –anisotropies in particle emission in the azimuthal angle (around the beam axis) –of strange hadrons should be sensitive to their interaction with the nuclear medium surrounding them during the collision [7, 9]. As such, studying the transverse flow of these particles may bring a significant advantage in our understanding of hadronic matter in the studied energy range.

This contribution will contain preliminary maps of v_n coefficients in the momentum phase space of strange hadrons – K^\pm mesons and Λ baryons –emitted from Ag+Ag collisions at a beam energy of 1.58 GeV/nucleon. The work is in progress, however, the results may already provide insight into the kaon-nucleon interaction. The transverse flow of Λ baryons is the first such result in this energy regime.

References

- [1] G. Agakichiev et al., Eur. Phys. J. A41, 243-277 (2009)
- [2] C. Hartnack et al., Phys. Rept.510, 119-200 (2012)
- [3] P. Danielewicz et al., Science298, 1592-1596 (2002)
- [4] W. Reisdorf et al., Nucl. Phys. A848, 366 (2010)
- [5] E. R. Mostet et al., Phys. Rev. D107, 043034 (2023)
- [6] T. Song et al., Phys. Rev. C103, 044901 (2021)
- [7] V. Zinyuk et al., Phys. Rev. C90, 025210(2014)
- [8] A. Ohnishi et al., EPJ Web of Conferences271, 08006 (2022)
- [9] Y. Nara et al., Phys. Rev. C106, 044902 (2022)

Category

Experiment

Collaboration (if applicable)

HADES

Author: ORLINSKI, Jan

Presenter: ORLINSKI, Jan

Session Classification: Poster session 2

Track Classification: Light and strange flavor physics & nuclei