

Poster: Colour breaking in the early universe: a minimal leptoquark model

The electroweak phase transition is a promising explanation for the origin of baryon asymmetry in the universe, a core problem in cosmology and particle physics.

An extension of the Standard Model is necessary to generate a strong first-order phase transition. Besides representing a target for several future-generation colliders, such Beyond the Standard Model (BSM) theories can generate - through a thermal phase transition - gravitational waves (GWs) potentially detectable by future space-based detectors, such as LISA 1, DECIGO, and BBO.

As a result, the interplay between BSM phenomenology and GWs is among the most active areas in the field of high-energy physics. Of particular interest are leptoquark (LQ) models, offering an alternative to conventional seesaw scenarios for the generation of Majorana neutrino masses at TeV scale. The presence of LQs can induce first order phase transitions with a temporary colour-breaking phase in the early universe.

With this poster, I intend to present results from the analysis of a minimal leptoquark model. In a dimensionally reduced effective theory approach 3, the model presents strong first order transitions, producing - in some scenarios - gravitational waves detectable by LISA. To our knowledge, these results provide the first evidence for the potential detection of color-breaking features in the above mentioned detectors.

The poster will be organized in 3 sections:

1. Introduction of the leptoquark model introduction and its features
2. Sketch of the pipeline implemented, and the adopted tools: (a) Computation of an effective potential with `DRalgo` 3. (b) Derivation of the phase structure and phase transition parameters by means of `CosmoTransitions`. (c) Derivation of the GW parameters with a dedicated python routine.
3. Illustration of the results (a) comparison of GW spectral peaks with detector sensitivities. (b) Interpretation of the various vacuum expectation value (vev) configurations identified in the two phases (c) correlation between collider observables.

1 Amaro-Seoane, P., Audley, H., Babak, S., Baker, J., Barausse, E., Bender, P., ... & Zweifel, P. (2017). *Laser interferometer space antenna*. arXiv preprint arXiv:1702.00786.

2 Felipe F. Freitas, João Gonçalves, António P. Morais, Roman Pasechnik, Werner Porod. *On interplay between flavour anomalies and neutrino properties*. Phys.Rev.D 108 (2023) 11, 115002.

3 Andreas Ekstedt, Philipp Schicho, Tuomas V.I. Tenkanen. *DRalgo: A package for effective field theory approach for thermal phase transitions*. Comput.Phys.Commun. 288 (2023) 108725.

Would you be interested in presenting a poster? (this will not impact the decision on your talk)

yes

Authors: EKSTEDT, Andreas; MORAIS, António (University of Aveiro); RATHSMAN, Johan (Lund University); FINETTI, Marco (Aveiro University); BERTENSTAM, Mårten (Lund University); PASECHNIK, Roman (Lund university)

Presenter: FINETTI, Marco (Aveiro University)

Session Classification: Reception and Poster Session