

First results on  
 $0\nu\beta\beta$  decay with  
the LEGEND  
experiment

Sofia Calgari – on behalf of  
the LEGEND Collaboration






[sofia.calgari@pd.infn.it](mailto:sofia.calgari@pd.infn.it)  
35th Rencontres de Blois

# Searching for $0\nu 2\beta$

## Signature

- $2\nu 2\beta$  (SM): continuous, broad spectrum  $\rightarrow$  observed ( $10^{18-21}$  yr)
- $0\nu 2\beta$  (BSM): peak at  $Q_{\beta\beta}$  (electrons energy)  $\rightarrow$  not observed ( $> 10^{21-26}$  yr)



## Physics Implications

- Nature of neutrinos (Dirac/Majorana) - *golden channel*
- Matter-antimatter asymmetry in the Universe
- Neutrino mass scale & ordering

Half-life

$$\frac{1}{T_{1/2}^{0\nu}} = G^{0\nu}(Q_{\beta\beta}, Z) |M^{0\nu}|^2 \left(\frac{m_{\beta\beta}}{m_e}\right)^2$$

Phase-space integral  
Nuclear Matrix Element (NME)

$$m_{\beta\beta} = \left| \sum_i U_{ei}^2 m_i \right| \rightarrow \text{to compare results obtained with different isotopes: } ^{76}\text{Ge}, ^{136}\text{Xe}, ^{130}\text{Te}, ^{100}\text{Mo}, ^{82}\text{Se}$$

Effective Majorana neutrino mass

Sensitivity

$$T_{1/2}^{0\nu} \propto \begin{cases} \epsilon \cdot f \cdot \sqrt{\frac{\epsilon}{BI \cdot \Delta E}} & \text{with bkg} \\ \epsilon \cdot f \cdot \epsilon & \text{without bkg} \end{cases}$$

$\epsilon$ : efficiency  
 $f$ : isotopic fraction  
 $\epsilon = \text{mass} \cdot \text{time}$ : exposure  
 $\Delta E$ : energetic resolution at  $Q_{\beta\beta}$   
 $BI$ : background index

<https://legend-exp.org/>

<https://github.com/legend-exp/>

# LEGEND

Large Enriched Germanium  
Experiment for Neutrinoless  $\beta\beta$  Decay



new groups

GERDA and MAJORANA DEMONSTRATOR  
set most stringent constraints for  $0\nu\beta\beta$  Ge-76



FNSNF



INFN



DFG



*"The collaboration aims to develop a phased,  $^{76}\text{Ge}$ -based  $0\nu\beta\beta$  decay experimental program with **discovery potential at a half-life beyond  $10^{28}$  yr**, using existing resources as appropriate to expedite physics results"*

LEGEND-200 [5 yr]

$\varepsilon = 1 \text{ t} \cdot \text{yr}$

$\text{BI} = 2 \times 10^{-4} \text{ cts}/(\text{keV} \cdot \text{kg} \cdot \text{yr})$

$T_{1/2}^{0\nu} > 10^{27} \text{ yr}$

$m_{\beta\beta} < 34 - 78 \text{ meV}$

LEGEND-1000 [10 yr]

$\varepsilon = 10 \text{ t} \cdot \text{yr}$

$\text{BI} = 10^{-5} \text{ cts}/(\text{keV} \cdot \text{kg} \cdot \text{yr})$

$T_{1/2}^{0\nu} > 1.3 \times 10^{28} \text{ yr}$

$m_{\beta\beta} < 9.4 - 21.4 \text{ meV}$



GERDA (PRL 125 252502, 2020)

$\varepsilon = 127.2 \text{ kg} \cdot \text{yr}$

$\text{BI} = 5.2 \times 10^{-4} \text{ cts}/(\text{keV} \cdot \text{kg} \cdot \text{yr})$



$T_{1/2}^{0\nu} > 1.8 \times 10^{26} \text{ yr}$

$m_{\beta\beta} < 79 - 180 \text{ meV}$



“By combining the lowest background levels and the best energy resolution in the field, LEGEND-1000 will perform a quasi-background-free search and can make an


**unambiguous discovery with just a handful of counts at the  $Q_{\beta\beta}$ ”**



# Why Germanium?


- No theoretical preferences
- Experimental preferences
  - costs
  - energy resolution
  - background level
  - scalability (liquids, gas, crystals)
- ${}^{76}_{32}\text{Ge} \rightarrow {}^{76}_{34}\text{Se} + 2e^{-} (+2\bar{\nu}_e)$  with  $Q_{\beta\beta}=2039.06(7)$  keV
- $Q_{\beta\beta} > 2$  MeV: less processes can mimic the  $0\nu 2\beta$  signal
- Natural abundance is low ( $\sim 8\%$ ): enrichment up to  $\sim 92\%$  is possible

$$\frac{1}{T_{1/2}^{0\nu}} = G^{0\nu}(Q_{\beta\beta}, Z) |M^{0\nu}|^2 \left(\frac{m_{\beta\beta}}{m_e}\right)^2$$




High Purity Germanium (HPGe) detector

# LEGEND-200 HPGe diodes






- p+ (implanted B), n+ (diffused Li), passivated groove
- Fully depleted crystals
- Different geometries - mass: 0.7-4 kg
- 130 kg operational (12 kg OFF for hardware issues)





# LEGEND-200

- @ Hall A of *Laboratori Nazionali del Gran Sasso (LNGS)*, Italy
- Same infrastructure of GERDA
- Rock overburden of 3500 m.w.e.
- Muon flux reduced to  $1.25 / (\text{m}^2\text{h})$  - reduction factor:  $\mathcal{O}(10^6)$






Array of HPGe diodes

Liquid argon (LAr) system

- 58 read-out modules of SiPMs coupled to WLS fibers



Outer Barrel (20+20)  
Inner Barrel (9+9)









Cryostat

- $\varnothing$  4m, H 5.88m
- LAr: 64 m<sup>3</sup>
- LLAMA for monitoring LAr properties


Ultrapure water tank

- Shields n,  $\gamma$
- 66 PMTs (Cherenkov) + plastic scintillators for  $\mu$
- $\varnothing$  10m, H 8.5 m, V 590 m<sup>3</sup>
- Clean room at 9.7 m





Pulse shape discrimination (PSD): signal discrimination based on the signal risetime and amplitude



## SILVER DATA SET - 76.2 kg·yr




- Background characterization
- All active detectors
- Extra 10.2/5.9 kg·yr of special background characterization runs without outer barrel/mini-shrouds


## GOLDEN DATA SET - 48.3 kg·yr

- Final neutrinoless double-beta decay data set
- All active detectors excluding those for which additional PSD cut studies are ongoing


(\* ) Test or HV scan runs - duty cycle of 91.4% (242 days in total)

- Weekly energy calibrations between physics runs using Th-228 sources
- Overall resolution of **0.1% FWHM at  $Q_{\beta\beta}$**  (including large IC detectors)
- Very stable energy scale - **energy bias  $0.3 \pm 0.2$  keV at  $Q_{\beta\beta}$**
- Data partitioned according to stability of energy observables





- Blinded analysis in  $Q_{\beta\beta} \pm 25$  keV
- Spectrum after:
  - data cleaning  $\rightarrow$  95-99% survival after removal of unphysical events
  - muon veto  $\rightarrow$  2 events removed at  $Q_{\beta\beta}$
  - multiplicity cut  $\rightarrow$  26% events removed at  $Q_{\beta\beta}$




LEGEND-200 · 09-2024




$$T_{1/2}^{2\nu} = 2.022 \pm 0.018_{stat} \pm 0.038_{syst} \times 10^{21} \text{ yr}$$

PRL 131, 142501 (2023)





- Decomposition of the full-range energy spectrum
- Bayesian background model using silver data set + 10.2 kg·yr of special runs
- Silver data .vs. simulations and material radioassay: Th-228 underprediction in physics data
  - this background is efficiently suppressed by analysis cuts
  - tested different Th-228 locations via the background model: no hotspots or asymmetries










- 60% suppression of Compton MSE at  $Q_{\beta\beta}$
- Cut acting on  $A/E = \max(\text{current}) / \text{energy}$ 
  - Late charge cut for PPC (large passivated surface)
  - Neural-network methods under development for Coaxial
- Data partitioned according to stability of PSD observables






- Strong anti-correlation of PSD & LAr cuts
- Characterized via special runs: 1 p.e. per 10 keV
- $\beta\beta$  decay signal acceptance of  $\sim 93\%$





5 surviving events  
in the analysis  
window (1930-  
2190 keV) before  
unblinding

- 7 events after unblinding (1 event at  $1.4\sigma$  from  $Q_{\beta\beta}$ )
- $BI = 5.3 \pm 2.2 \times 10^{-4}$  cts/(keV·kg·yr)
  - world-leading result
  - goal:  $2 \times 10^{-4}$  cts/(keV · kg · yr)
- Unbinned fit of GERDA<sup>(\*)</sup> + MJD<sup>(\*\*)</sup> + LEGEND-200
  - $p$ -value of background-only = 26%
  - Observed  $T_{1/2} > 1.9 \times 10^{26}$  yr @ 90% CL
  - Sensitivity  $T_{1/2} = 2.8 \times 10^{26}$  yr @ 90% CL  
+30% thanks to LEGEND-200



(\*) [PRL 125 252502 \(2020\)](#) (\*\*) [PRL 130 062501 \(2023\)](#)

# Summary

- LEGEND-200
  - fully operative at LNGS
  - first unblinded results over 1 yr of data taking
  - +30% in  $0\nu 2\beta$  median sensitivity
  - allows for a prompt investigation of issues
  - powerful LEGEND-1000 test-bench
  - ongoing analysis for testing background hypotheses with different setups
  - ongoing radioassay campaign
  - maintenance work (gain in background rejection)
  - install new 35 kg of enr-Ge + restart data taking at the end of 2024
- LEGEND-1000
  - preparations underway at LNGS following Borexino decommissioning
  - funding sought from U.S. (DOE and NSF) and EU
  - funding already in hand from several EU institutions
  - data taking will start in the next decade

Thank you for your attention!

