

Energy-energy correlators in jets across collision systems with ALICE

Anjali Nambrath for the ALICE collaboration

University of California, Berkeley

nambrath@berkeley.edu

Hard Probes 2024
September 22-27, Nagasaki




In this talk: differential measurements of EECs

1. EECs in pp: final results
universal shape independent of jet p_T
2. First measurement of EECs in D^0 -tagged jets
probing flavor effects
3. First measurement of EECs in p-Pb
probing EECs in high-multiplicity environments


The two-point EEC is calculated from pairs of tracks.

EEC definition:

$$\frac{d\sigma_{EEC}}{dR_L} = \sum_{i,j} \int d\sigma(R'_L) \frac{p_{T,i} p_{T,j}}{p_{T,jet}^2} \delta(R'_L - R_{L,ij})$$




1. For each pair of tracks inside the jet, calculate the **energy weight**.
2. Count the number of weighted track pairs as a function of R_L .



EECs show a clear transition region.

- Transition between **perturbative** (large R_L , partonic) and **non-perturbative** regimes (hadronic, small R_L)
- Time evolution of jet formation is imprinted onto the EEC angular scaling
- EECs let us probe jet formation and confinement!



EECs in pp



Clear separation of perturbative and non-perturbative regions.

EEC peak is visibly dependent on jet p_T .

Universal transition region


after rescaling the x-axis to $\langle p_T^{\text{ch jet}} \rangle R_L$ (common peak position and height).

NEW on arXiv! [arXiv:2409.12687](https://arxiv.org/abs/2409.12687) [hep-ex]




How does the shower depend on flavor?

- At the largest R_L , the scaling behavior in heavy-flavor jets is identical to light quark jets.
- **Turnover exhibits a mass dependence!**
 - Flavor effects can be probed with ratios to inclusive jets.
- Change of shape at small angles is a consequence of the dead cone.




gluon jets




$$\frac{C_A}{C_F} = \frac{9}{4}$$



quark jets



heavy quark jets



Comparing D⁰-tagged jet to inclusive jet EECs

ALI-PREL-579241



Upper panel:

- p_T cut on leading track in inclusive jets to study fragmentation bias
- slopes at large R_L seem different, significant suppression at all R_L
- peak positions are similar due to gluon contribution to inclusive

From the ratios:

- D⁰/inclusive → mass + Casimir
- D⁰/LF → isolated mass effects

Clear mass effect in D⁰ jets!




See poster by P. Dankher for more details on the D⁰ EEC measurement!
 See poster by B. Liang-Gilman for D⁰ Monte Carlo studies!


Talk by J. Klein tomorrow at 12:10pm on more HF jets from ALICE!

Probing hadronization with pp EECs

Lund string models: PYTHIA 8, Sherpa Lund
Cluster models: Herwig 7, Sherpa AHADIC



arXiv:2409.12687



Inclusive jets:

- PYTHIA & Herwig perform well, *Herwig captures peak position*
- Sherpa Lund does well, AHADIC does not
- *both cluster models peak at smaller R_L*

Heavy flavor jets:


- *data favors PYTHIA's implementation*
- Herwig overpredicts inclusive jet EECs; underpredicts in HF
- Sherpa Lund underpredicts the data
- AHADIC fails to describe the peak

Cluster hadronization for higher p_T jets, string-breaking models for D^0 jets?


Are EECs modified in p-Pb?

Differences from pp:


- initial state (nPDF, isospin)
- final-state interactions?
- comovers? collectivity??



EECs in p-Pb are a window into interactions in small systems.



EECs in p-Pb have the same overall shape as pp.



About the data:

- $R = 0.4$ charged-particle jets
- $p_T^{\text{ch jet}}$ in [20, 80] GeV/c
- threshold cut: $p_T^{\text{trk}} > 1$ GeV/c

These are subtracted for UE and corrected for detector effects:

- background subtraction for jet p_T and EEC distribution
- bin-by-bin correction for detector effects


EEC measurements in more complex systems require background subtraction techniques.

1. Subtract UE energy density from the jet p_T :

$$\rho = \text{median} \left\{ \frac{p_{T,\text{jet}}^{k_T}}{A_{\text{jet}}^{k_T}} \right\} \cdot C \quad C = \frac{\sum_j A_j}{A_{\text{acc}}}$$

2. Correct the EEC distribution for combinatorial background:


- **signal-signal** (what we want!)
- **signal-background**
- **background-background**



We use the perpendicular cone to estimate the latter two contributions.

These subtraction steps are also performed for the pp baseline.

EECs are modified in p-Pb in the lowest jet p_T bin.



- Significant difference between EECs in p-Pb compared to pp!

- jet structure appears to be altered only in the lowest jet p_T range

- Initial state effect?

- some models lead to a qualitatively similar effect




- Final state effect?

- modification is qualitatively consistent with ALICE measurement* of HM/MB z_{ch} in pp

* arXiv:2311.13322


The transition region in p-Pb resembles pp.

- Universality of the EEC peak position across jet p_T and collision system.
- EEC peak height for 20-40 GeV/c jets is slightly lower than for other jets.



ALI-PREL-579739

Varying the track cut changes the EEC behavior at large R_L .



Strong sensitivity to track p_T cut in low p_T jets!

- non-perturbative effects increase for lower jet p_T


Track cut modifies the enhancement in ratio

- but not the small- R_L suppression

Suggests that the origin of the effect lies in softer interactions at small x !

Neither CGC or nPDF models fully capture the enhancement and suppression seen in data.


- CGC predictions* from Haoyu Liu et. al., over a range of saturation scale values



* models provided by Haoyu Liu, Ian Moult, and Xiaohui Liu

Neither CGC or nPDF models fully capture the enhancement and suppression seen in data.


- CGC predictions* from Haoyu Liu et. al., over a range of saturation scale values
- PYTHIA results use:
 - nPDF: EPPS21nlo_CT18Anlo
 - PDF: CT14nlo
- nPDFs are within $\sim 1\sigma$ at small R_L – but these are very large uncertainties



* models provided by Haoyu Liu, Ian Moutl, and Xiaohui Liu

Higher-twist formalism calculations come closer to reproducing the data.


- HT calculations* of final-state interactions from Yu Fu et al. show a stronger effect is possible
 - R_{pPb} (nuclear modification of nPDF) is chosen to be 0.85 (for $x=0.01$)
 - \hat{q} is $0.02 \text{ GeV}^2/\text{fm}$ (BDMPS 1997)
 - L is varied (R_{pb} is 5.5 fm)
- Simulation (e.g. JETSCAPE) required for a more realistic estimate!



* models provided by Yu Fu, Berndt Mueller, and Chaturanga Sirimanna

$\log(R_L)^2$ shape of the p-Pb/pp ratio – why?

- HT calculations* of final-state interactions from Yu Fu et al. show a stronger effect is possible
 - R_{pPb} (nuclear modification of nPDF) is chosen to be 0.85 (for $x=0.01$)
 - \hat{q} is 0.02 GeV²/fm (BDMPS 1997)
 - L is varied (R_{Pb} is 5.5 fm)
- The ratio appears to follow a $\log(R_L)^2$ scaling. Why: change in q/g ratio?



* models provided by Yu Fu, Berndt Mueller, and Chaturanga Sirimanna

Summary and outlook

- Universality of EEC shape and turnover in pp
- EECs are altered in HF jets
 - dramatic reduction in amplitude – clear flavor effect
- EECs are modified in p-Pb
 - modification is not explained by purely initial-state models

More EECs from ALICE:

Preeti Dhankher

D⁰-jet EEC measurement

poster session tonight

Beatrice Liang-Gilman

D⁰-jet EECs in Monte Carlo

poster session tonight

Anjali Nambrath

p-Pb EEC measurement

poster session tonight

Ananya Rai


E3C/E2C measurement

talk today – 12:10pm


Backup

D⁰ reconstruction steps



D⁰ candidates were reconstructed from daughter tracks using topological and particle identification selections (D⁰ → K⁻ + π⁺, and charge conjugates).




D⁰-tagged charged jets were created using the anti-k_T algorithm (R=0.4) for each candidate.



Invariant mass analysis was performed to remove combinatorial K⁻π⁺ pairs surviving the D⁰ selections.



Corrected the EECs for the efficiency of D⁰-tagged jet reconstruction and removed the contribution from beauty decays.




Corrected the EECs for detector effects with a bin-by-bin correction method.

Perp. cone for combinatorial EEC background

- Particles from jet: $\text{sig} + \text{bkg}$
- Particles from perp cone: bkg'
- Pairs in the combined cone:

$$\begin{aligned} (\text{sig} + \text{bkg} + \text{bkg}')(\text{sig} + \text{bkg} + \text{bkg}') &= \text{sig}^* \text{sig} + 2\text{sig}^* \text{bkg} \\ &+ \text{bkg}^* \text{bkg} + \boxed{2\text{sig}^* \text{bkg}' + 2\text{bkg}^* \text{bkg}'} + \boxed{\text{bkg}'^* \text{bkg}'} \\ &\quad \text{jet-perp} \qquad \text{perp-perp} \end{aligned}$$




Perp. cone for combinatorial EEC background


- Particles from jet: **sig** + **bkg**
- Particles from perp cone: **bkg'**
- Pairs in the combined cone:

$$\begin{aligned}
 (\text{sig} + \text{bkg} + \text{bkg}')(\text{sig} + \text{bkg} + \text{bkg}') &= \text{sig}^* \text{sig} + 2\text{sig}^* \text{bkg} \\
 + \text{bkg}^* \text{bkg} + &\boxed{2\text{sig}^* \text{bkg}' + 2\text{bkg}^* \text{bkg}'} + \boxed{\text{bkg}'^* \text{bkg}'} \\
 &\quad \text{jet-perp} \qquad \qquad \text{perp-perp}
 \end{aligned}$$


- Sig-bkg pairs: **jet-perp** — 2 **perp-perp**
- Bkg-bkg pairs: **perp-perp**
- **Total background: jet-perp — perp-perp**




p-Pb and pp comparison



ALI-PREL-581875




ALI-PREL-581880



ALI-PREL-581885

Quark-jet and gluon-jet EECs



Quark-jet and gluon-jet and D^0 -tagged jet EECs

