

# Method of semi-inclusive jet mass measurement in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV with STAR

Tuesday 24 September 2024 18:10 (20 minutes)

Jet quenching phenomenon serves as a crucial signature of the Quark-Gluon Plasma, observed when hard-scattered partons interact with the hot, dense QCD medium created in high-energy heavy-ion collisions. In central heavy-ion collisions, however, distinguishing jets produced by hard scattering from those originating from combinatorial background is largely limited, especially for jets with low transverse momenta ( $p_{T,jet}$ ). To address this challenge, methods for measurements of semi-inclusive recoil jets with respect to a trigger particle have been devised, leading to measurements of jet yields to the unprecedentedly low  $p_{T,jet}$  range. In particular, the STAR Collaboration has combined this semi-inclusive recoil jets measurement with a mixed-event technique as a data-driven method for the correction of uncorrelated background effects. We aim to extend the scope of the semi-inclusive approach into measurements of jet mass ( $M_{jet}$ ), and develop a 2-dimensional correction framework as a function of ( $p_{T,jet}$ ,  $M_{jet}$ ).

In this poster, we discuss the method of semi-inclusive jet mass measurements, and provide the closure test result based on simulation. Jets from PYTHIA events are embedded into  $\sqrt{s_{NN}} = 200$  GeV Au+Au collision background obtained from a thermal model. Correction procedures, including the subtraction of combinatorial jet contributions via a mixed-event technique and 2-dimensional unfolding, are tested.

## Category

Experiment

## Collaboration

STAR

**Author:** KANG, jeongmyung (Sejong University)

**Presenter:** KANG, jeongmyung (Sejong University)

**Session Classification:** Poster Session

**Track Classification:** 1. Jets modification and medium response