Contribution ID: 301

Type: Poster

Measurements of inclusive J/ ψ and ψ (2S) production at midrapidity in pp collisions at \sqrt{s} = 13.6 TeV with ALICE

Tuesday 24 September 2024 18:10 (20 minutes)

Quarkonium production in high-energy proton-proton (pp) collisions is an important tool for studying perturbative and non-perturbative aspects of quantum chromodynamics. Charmonia are bound states of charm and anti-charm quarks and their production process can be factorized into two stages: the heavy quark production and the formation of the bound state. The former happens within initial hard parton-parton scatterings with large momentum transfers, and can be well described by perturbative QCD. The second one, which involves long distances and soft momentum scales, is a typical non-perturbative process. Measurements of J/ ψ and ψ (2S) cross section in pp collisions are crucial for studying charmonium production mechanisms and testing different QCD-based model calculations. They can also provide a reference for investigating the quark-gluon plasma formed in nucleus-nucleus collisions and the cold nuclear matter effects in proton-nucleus collisions at $\sqrt{s} = 13.6$ TeV. The analysis is based on the data collected in 2022 by the upgraded ALICE detector during LHC Run 3, which offers significantly higher statistics compared to Run 1 and 2. The $p_{\rm T}$ -differential production of inclusive J/ ψ cross section, as well as the ψ (2S)-to-J/ ψ ratio, will be reported. Results will be shown along with similar measurements at forward rapidity and compared with model calculations.

Category

Experiment

Collaboration

ALICE

Author: ZHANG, Yuan (University of Science and Technology of China (CN))Presenter: ZHANG, Yuan (University of Science and Technology of China (CN))Session Classification: Poster Session

Track Classification: 3. Heavy quarks and quarkonia