

Higgs & Beyond



John Ellis

KING'S
College
LONDON

Why the Higgs boson?
What can the Higgs boson tell us?
Looking beyond it

Remember the lectures by Anna Ku

1957

The BCS Theory of Superconductivity

PHYSICAL REVIEW

VOLUME 108, NUMBER 5

DECEMBER 1, 1957

Theory of Superconductivity*

J. BARDEEN, L. N. COOPER,[†] AND J. R. SCHRIEFFER[‡]
Department of Physics, University of Illinois, Urbana, Illinois

(Received July 8, 1957)

A theory of superconductivity is presented, based on the fact that the interaction between electrons resulting from virtual exchange of phonons is attractive when the energy difference between the electrons states involved is less than the phonon energy, $\hbar\omega$. It is favorable to form a superconducting phase when this attractive interaction dominates the repulsive screened Coulomb interaction. The normal phase is described by the Bloch individual-particle model. The ground state of a superconductor, formed from a linear combination of normal state configurations in which electrons are virtually excited in pairs of opposite spin and momentum, is lower in energy than the normal state by amount proportional to an average $(\hbar\omega)^2$, consistent with the isotope effect. A mutually orthogonal set of excited states in

one-to-one correspondence with those of the normal phase is obtained by specifying occupation of certain Bloch states and by using the rest to form a linear combination of virtual pair configurations. The theory yields a second-order phase transition and a Meissner effect in the form suggested by Pippard. Calculated values of specific heats and penetration depths and their temperature variation are in good agreement with experiment. There is an energy gap for individual-particle excitations which decreases from about $3.5kT_c$ at $T=0^\circ\text{K}$ to zero at T_c . Tables of matrix elements of single-particle operators between the excited-state superconducting wave functions, useful for perturbation expansions and calculations of transition probabilities, are given.

Condensate of electron pairs of due to phonon interactions
 Lowest-energy state has charge density: breaks/hides $U(1)_{em}$

1959/62

Nambu, Anderson & “Spontaneous Breaking” of Gauge Symmetry

“Spontaneous symmetry breaking” = hidden symmetry
Gauge-invariant mass generation by plasmons in non-relativistic theory

PHYSICAL REVIEW

VOLUME 117, NUMBER 3

FEBRUARY 1, 1960

Quasi-Particles and Gauge Invariance in the Theory of Superconductivity*

YOICHIRO NAMBU

The Enrico Fermi Institute for Nuclear Studies and the Department of Physics, The University of Chicago, Chicago, Illinois

(Received July 23, 1959)

Ideas and techniques known in quantum electrodynamics have been applied to the Bardeen-Cooper-Schrieffer theory of superconductivity. In an approximation which corresponds to a generalization of the Hartree-Fock fields, one can write down an integral equation defining the self-energy of an electron in an electron gas with phonon and Coulomb interaction. The form of the equation implies the existence of a particular solution which does not follow from perturbation theory, and which leads to the energy gap equation and the quasi-particle picture analogous to Bogoliubov's.

The gauge invariance, to the first order in the external electro-

magnetic field, can be maintained in the quasi-particle picture by taking into account a certain class of corrections to the charge-current operator due to the phonon and Coulomb interaction. In fact, generalized forms of the Ward identity are obtained between certain vertex parts and the self-energy. The Meissner effect calculation is thus rendered strictly gauge invariant, but essentially keeping the BCS result unaltered for transverse fields.

It is shown also that the integral equation for vertex parts allows homogeneous solutions which describe collective excitations of quasi-particle pairs, and the nature and effects of such collective states are discussed.

PHYSICAL REVIEW

VOLUME 130, NUMBER 1

1 APRIL 1963

Plasmons, Gauge Invariance, and Mass

P. W. ANDERSON

Bell Telephone Laboratories, Murray Hill, New Jersey

(Received 8 November 1962)

Schwinger has pointed out that the Yang-Mills vector boson implied by associating a generalized gauge transformation with a conservation law (of baryonic charge, for instance) does not necessarily have zero mass, if a certain criterion on the vacuum fluctuations of the generalized current is satisfied. We show that the theory of plasma oscillations is a simple nonrelativistic example exhibiting all of the features of Schwinger's idea. It is also shown that Schwinger's criterion that the vector field $m \neq 0$ implies that the matter spectrum before including the Yang-Mills interaction contains $m=0$, but that the example of superconductivity illustrates that the physical spectrum need not. Some comments on the relationship between these ideas and the zero-mass difficulty in theories with broken symmetries are given.

1964

The Founders

Tom Kibble



Gerry Guralnik



Carl Hagen




François Englert



Robert Brout



Peter Higgs



1964

The (GN)AEBHGHKMP Mechanism

BROKEN SYMMETRY AND THE MASS OF GAUGE VECTOR MESONS*

F. Englert and R. Brout

Faculté des Sciences, Université Libre de Bruxelles, Bruxelles, Belgium

(Received 26 June 1964)

BROKEN SYMMETRIES, MASSLESS PARTICLES AND GAUGE FIELDS

P. W. HIGGS

Tait Institute of Mathematical Physics, University of Edinburgh, Scotland

Received 27 July 1964

VOLUME 13, NUMBER 16

PHYSICAL REVIEW LETTERS

4

BROKEN SYMMETRIES AND THE MASSES OF GAUGE VECTOR MESONS

Peter W. Higgs

Tait Institute of Mathematical Physics, University of Edinburgh, Scotland

(Received 31 August 1964)

The only one who mentioned a massive scalar boson

GLOBAL CONSERVATION LAWS AND MASSLESS PARTICLES*

G. S. Guralnik,† C. R. Hagen,‡ and T. W. B. Kibble
Department of Physics, Imperial College, London, England

(Received 12 October 1964)

SPONTANEOUS BREAKDOWN OF STRONG INTERACTION SYMMETRY AND THE ABSENCE OF MASSLESS PARTICLES


A. A. MIGDAL and A. M. PERELMAN

Submitted to JETP editor November 30, 1965; resubmitted February 16, 1966

J. Experimental and Theoretical Physics (USSR) 51, 125-146 (1966)

The occurrence of massless particles in the presence of spontaneous symmetry breakdown is discussed. By summing all Feynman diagrams, one obtains for the difference of the mass


Nambu, **EB, H, GHK** & Higgs



Spontaneous symmetry breaking: massless Nambu-Goldstone boson 'eaten' by massless gauge boson

Accompanied by massive scalar particle

Hungry for Higgs



(FLIP TANEDO / QUANTUM DIARIES)

Steps Towards the Higgs Boson

CAN ONE EVADE THE GOLDSTONE THEOREM?

P.W. ANDERSON POINTED OUT THAT IN A SUPERCONDUCTOR THE GOLDSTONE MODE BECOMES A MASSIVE "PLASMON" MODE DUE TO ITS ELECTROMAGNETIC INTERACTION, AND THAT THIS MODE IS JUST THE LONGITUDINAL PARTNER OF TRANSVERSELY POLARIZED ELECTROMAGNETIC MODES, WHICH ARE ALSO MASSIVE. (MEISSNER EFFECT!)

ANDERSON CONTINUED, "THE GOLDSTONE ZERO-MASS DIFFICULTY IS NOT A SERIOUS ONE, BECAUSE WE CAN PROBABLY CANCEL IT OFF AGAINST AN EQUAL YANG-MILLS ZERO-MASS PROBLEM"

- BUT (a) HE DIDN'T DISCUSS THE THEOREM
 (b) HE DIDN'T DISCUSS ANY RELATIVISTIC MODEL

1964 HOW TO EVADE GOLDSTONE'S THEOREM

G-SW PROOF INVOLVES COMMUTATOR

$$[\hat{\phi}, \hat{\phi}_0] = \hat{\phi}_0 \quad (1)$$

$$\hat{\phi} = \int d^3x \hat{j}_0(x, t) \quad (\text{GENERATOR})$$

AND $\partial_\mu \hat{j}^\mu = 0 \quad (2)$ (INVARIANCE OF $\hat{\phi}$)

MANIFEST LORENTZ INVARIANCE \Rightarrow

4D FOURIER TRANSFORM OF

$$\langle i [\hat{j}_\mu(x), \hat{\phi}_0(y)] \rangle_0$$

HAS FORM $k_\mu (\text{sign } k_0) \rho(k^2)$ [$\rho = 0$ FOR SPACELIKE k]

(2) $\rightarrow k^2 \rho(k^2) = 0 \Rightarrow \rho = C \delta(k^2)$

(1) $\rightarrow C = 2\pi \langle \hat{\phi}_0 \rangle_0 \neq 0$ (ASYMMETRIC VACUUM)

MARCH 1964

A. KLEIN & B.W. LEE

FOR (e.g.) SUPERCONDUCTOR, F.T. HAS MORE GENERAL FORM

$$k_\mu \rho_1(k^2, n \cdot k) + n_\mu \rho_2(k^2, n \cdot k)$$

WHERE $n_\mu (= (1, 0, 0, 0))$ SPECIFIES REST FRAME OF IONIC BACKGROUND.

PERHAPS THIS COULD HAPPEN IN TRULY RELATIVISTIC CASE?

JUNE 1964 W. GILBERT No!

JULY 1964 P.W.H. YES! —

BUT ONLY IF GAUGE FIELD A_μ IS COUPLED TO THE CURRENT

1964 ACCIDENTAL BIRTH OF A BOSON

Th. 16 July Phys. Rev. Letters (22 June), containing Gilbert's paper reaches Edinburgh
 F. 24 July Broken Symmetries, Massless Particles and Gauge Fields (P.W.H.) sent to Physics Letters editor at CERN
ACCEPTED

F. 31 July Broken Symmetries and the masses of Gauge Bosons (P.W.H.) sent to Physics Letters editor at CERN.
REJECTED

August Paper resubmitted (inter alia)
 "It is worth noting that an essential feature of this type of theory is the prediction of incomplete multiplets of scalar and vector bosons"


31 August Revised paper received by Physical Review Letters.
ACCEPTED

Referee (Nambu) draws to attention of PWH the paper by Englert & R. Brout, Broken Symmetry and the Mass of Gauge Vector Mesons (received by Phys. Rev. Letters 22 June, published 31 August)

The Nambu-Goldstone Mechanism

- Postulated effective scalar potential:

$$V[\phi] = -\mu^2 \phi^\dagger \phi + \lambda (\phi^\dagger \phi)^2$$



- Minimum energy at non-zero value:

$$\phi_0 = \langle 0 | \phi | 0 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ +v \end{pmatrix} \quad v = \sqrt{\frac{-\mu^2}{\lambda}}$$

- Components of scalar field: $\phi(x) = \frac{1}{\sqrt{2}}(v + \sigma(x))e^{i\pi(x)}$

- π massless, σ massive:

$$m_H^2 = 2\mu^2 = 2\lambda v$$

Abelian EBH Mechanism

- Lagrangian

$$\mathcal{L} = (D_\mu \phi)^\dagger (D^\mu \phi) - V(|\phi|) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu}, \quad D_\mu = \partial_\mu - ieA_\mu$$

- Gauge transformation

$$\phi'(x) = e^{i\alpha(x)} \phi(x) = e^{i\alpha(x)} e^{i\theta(x)} \eta(x)$$

$$A'_\mu(x) = A_\mu(x) + \frac{1}{e} \partial_\mu \alpha(x)$$

- Choose $\alpha(x) = -\theta(x)$: $\phi'(x) = \eta(x)$

- Rewrite Lagrangian: $\mathcal{L} = |(\partial - ieA'_\mu)\eta|^2 - V(\eta) - \frac{1}{4} F'_{\mu\nu} F'^{\mu\nu}$

$$\mathcal{L} = |(\partial_\mu - ieA'_\mu)(v + \frac{1}{\sqrt{2}}H)|^2 - \frac{1}{4} F'_{\mu\nu} F'^{\mu\nu} - V$$

$$= \underbrace{-\frac{1}{4} F'_{\mu\nu} F'^{\mu\nu} + v^2 e^2 A'_\mu A'^\mu}_{\text{massive A-field, } m_A \sim ev} + \underbrace{\frac{1}{2} [(\partial_\mu H)^2 - m_H^2 H^2]}_{\text{neutral scalar, } m_H \neq 0} + \dots$$

massive A-field, $m_A \sim ev$

neutral scalar, $m_H \neq 0$

Think of a Snowfield



Skier moves fast:

Like particle without mass
e.g., photon = particle of light



Snowshoer sinks into snow,
moves slower:



Like particle with mass
e.g., electron

**The LHC discovered
the snowflake:
The Higgs Boson**

Hiker sinks deep,
moves very slowly:
Particle with large mass



Weinberg: A Model of Leptons

- Electroweak sector of the Standard Model
- SU(2) x U(1)
- Mixing of Z, photon
- Neutral currents
- Higgs-lepton couplings
- No quarks

2 citations before 1971

and

$$\varphi_1 = (\varphi^0 + \varphi^{0\dagger} - 2\lambda)/\sqrt{2} \quad \varphi_2 = (\varphi^0 - \varphi^{0\dagger})/i\sqrt{2}. \quad (5)$$

The condition that φ_1 have zero vacuum expectation value to all orders of perturbation theory tells us that $\lambda^2 \cong M_1^2/2h$, and therefore the field φ_1 has mass M_1 while φ_2 and φ^- have mass zero. But we can easily see that the Goldstone bosons represented by φ_2 and φ^- have no physical coupling. The Lagrangian is gauge invariant, so we can perform a combined isospin and hypercharge gauge transformation which eliminates φ^- and φ_2 everywhere⁵ without changing anything else. We will see that G_e is very small, and in any case M_1 might be very large,⁷ so the φ_1 couplings will also be disregarded in the following.

The effect of all this is just to replace φ everywhere by its vacuum expectation value

$$\langle \varphi \rangle = \lambda \begin{pmatrix} 1 \\ 0 \end{pmatrix}. \quad (6)$$

The first four terms in \mathcal{L} remain intact, while the rest of the Lagrangian becomes

$$-\frac{1}{8}\lambda^2 g^2 [(A_\mu^1)^2 + (A_\mu^2)^2] - \frac{1}{8}\lambda^2 (gA_\mu^3 + g'B_\mu)^2 - \lambda G_e \bar{e}e. \quad (7)$$

We see immediately that the electron mass is λG_e . The charged spin-1 field is

$$W_\mu = 2^{-1/2}(A_\mu^1 + iA_\mu^2) \quad (8)$$

and has mass

$$M_W = \frac{1}{2}\lambda g. \quad (9)$$

The neutral spin-1 fields of definite mass are

$$Z_\mu = (g^2 + g'^2)^{-1/2}(gA_\mu^3 + g'B_\mu), \quad (10)$$

$$A_\mu = (g^2 + g'^2)^{-1/2}(-g'A_\mu^3 + gB_\mu). \quad (11)$$

Their masses are

$$M_Z = \frac{1}{2}\lambda(g^2 + g'^2)^{1/2}, \quad (12)$$

$$M_A = 0, \quad (13)$$

so A_μ is to be identified as the photon field. The interaction between leptons and spin-1 mesons is

$$\frac{ig}{2\sqrt{2}} \bar{e} \gamma^\mu (1 + \gamma_5) \nu W_\mu + \text{H.c.} + \frac{igg'}{(g^2 + g'^2)^{1/2}} \bar{e} \gamma^\mu e A_\mu + \frac{i(g^2 + g'^2)^{1/2}}{4} \left[\left(\frac{3g'^2 - g^2}{g'^2 + g^2} \right) \bar{e} \gamma^\mu e - \bar{\nu} \gamma^\mu \nu \gamma_5 e + \nu \gamma^\mu (1 + \gamma_5) \nu \right] Z_\mu. \quad (14)$$

We see that the rationalized electric charge is

$$e = gg' / (g^2 + g'^2)^{1/2} \quad (15)$$

and, assuming that W_μ couples as usual to hadrons and muons, the usual coupling constant of weak interactions is given by

$$G_W/\sqrt{2} = g^2/8M_W^2 = 1/2\lambda^2. \quad (16)$$

Note that then the e - φ coupling constant is

$$G_e = M_e/\lambda = 2^{1/4} M_e G_W^{1/2} = 2.07 \times 10^{-6}.$$

The coupling of φ_1 to muons is stronger by a factor M_μ/M_e , but still very weak. Note also that (14) gives g and g' larger than e , so

by this model have to do with the couplings of the neutral intermediate meson Z_μ . If Z_μ does not couple to hadrons then the best place to look for effects of Z_μ is in electron-neutron scattering. Applying a Fierz transformation to the W -exchange terms, the total effective e - ν interaction is

$$\frac{G_W}{\sqrt{2}} \bar{\nu} \gamma_\mu (1 + \gamma_5) \nu \left\{ \frac{(3g^2 - g'^2)}{2(g^2 + g'^2)} \bar{e} \gamma^\mu e + \frac{3}{2} \bar{e} \gamma^\mu \nu \gamma_5 e \right\}.$$

If $g \gg e$ then $g \gg g'$, and this is just the usual e - ν scattering matrix element times an extra factor $\frac{3}{2}$. If $g \approx e$ then $g \ll g'$, and the vector

“Whatever the final laws of nature may be, there is no reason to suppose that they are designed to make physicists happy.”

Summary of the Standard Model

- Particles and $SU(3) \times SU(2) \times U(1)$ quantum numbers:

L_L	$\begin{pmatrix} \nu_e \\ e^- \end{pmatrix}_L, \begin{pmatrix} \nu_\mu \\ \mu^- \end{pmatrix}_L, \begin{pmatrix} \nu_\tau \\ \tau^- \end{pmatrix}_L$	$(1, 2, -1)$
E_R	e_R^-, μ_R^-, τ_R^-	$(1, 1, -2)$
Q_L	$\begin{pmatrix} u \\ d \end{pmatrix}_L, \begin{pmatrix} c \\ s \end{pmatrix}_L, \begin{pmatrix} t \\ b \end{pmatrix}_L$	$(3, 2, +1/3)$
U_R	u_R, c_R, t_R	$(3, 1, +4/3)$
D_R	d_R, s_R, b_R	$(3, 1, -2/3)$

- Lagrangian:

$\mathcal{L} =$	$-\frac{1}{4} F_{\mu\nu}^a F^{a\ \mu\nu}$	gauge interactions	Tested < 0.1% before LHC
	$+ i\bar{\psi} \not{D}\psi + h.c.$	matter fermions	
	$+ \psi_i y_{ij} \psi_j \phi + h.c.$	Yukawa interactions	Testing now in progress
	$+ D_\mu \phi ^2 - V(\phi)$	Higgs potential	

Parameters of the Standard Model

- Gauge sector:
 - 3 gauge couplings: g_3, g_2, g'
 - 1 strong CP-violating phase

Unification?

- Yukawa interactions:
 - 3 charged-lepton masses
 - 6 quark masses
 - 4 CKM angles and phase

Flavour?

- Higgs sector:
 - 2 parameters: μ, λ

Mass?

- **Total: 19 parameters**

The Standard Model Lagrangian

$$\mathcal{L}_{SM} = \mathcal{L}_m + \mathcal{L}_g + \mathcal{L}_h + \mathcal{L}_y \quad ,$$

$$\mathcal{L}_m = \bar{Q}_L i \gamma^\mu D_\mu^L Q_L + \bar{q}_R i \gamma^\mu D_\mu^R q_R + \bar{L}_L i \gamma^\mu D_\mu^L L_L + \bar{l}_R i \gamma^\mu D_\mu^R l_R$$

$$\mathcal{L}_G = -\frac{1}{4} B_{\mu\nu} B^{\mu\nu} - \frac{1}{4} W_{\mu\nu}^a W^{a\mu\nu} \quad \checkmark \text{ Experiment: accuracy } < \%$$

$$\mathcal{L}_H = (D_\mu^L \phi)^\dagger (D^{L\mu} \phi) - V(\phi)$$

$$\mathcal{L}_Y = y_d \bar{Q}_L \phi q_R^d + y_u \bar{Q}_L \phi^c q_R^u + y_L \bar{L}_L \phi l_R +$$

No direct evidence
until July 4, 2012

$$D_\mu^L = \partial_\mu - ig W_\mu^a T^a - iY g' B_\mu \quad , \quad D_\mu^R = \partial_\mu - iY g' B_\mu$$

$$V(\phi) = -\mu^2 \phi^2 + \lambda \phi^4 \quad .$$

Masses for SM Gauge Bosons

- Kinetic terms for SU(2) and U(1) gauge bosons:

$$\mathcal{L} = -\frac{1}{4} G_{\mu\nu}^i G^{i\mu\nu} - \frac{1}{4} F_{\mu\nu} F^{\mu\nu}$$

where $G_{\mu\nu}^i \equiv \partial_\mu W_\nu^i - \partial_\nu W_\mu^i + ig\epsilon_{ijk} W_\mu^j W_\nu^k$ $F_{\mu\nu} \equiv \partial_\mu W_\nu^i - \partial_\nu W_\mu^i$

- Kinetic term for Higgs field:

$$\mathcal{L}_\phi = -|D_\mu \phi|^2 \quad D_\mu \equiv \partial_\mu - i g \sigma_i W_\mu^i - i g' Y B_\mu$$


- Expanding around vacuum: $\phi = \langle 0|\phi|0 \rangle + \hat{\phi}$

$$\mathcal{L}_\phi \ni -\frac{g^2 v^2}{2} W_\mu^+ W^{\mu-} - \frac{g'^2 v^2}{2} B_\mu B^\mu + g g' v^2 B_\mu W^{\mu 3} - g^2 \frac{v^2}{2} W_\mu^3 W^{\mu 3}$$

- Boson masses:

$$m_{W^\pm} = \frac{gv}{2} \quad Z_\mu = \frac{gW_\mu^3 - g'B_\mu}{\sqrt{g^2 + g'^2}} : m_Z = \frac{1}{2}\sqrt{g^2 + g'^2}v ; \quad A_\mu = \frac{g'W_\mu^3 + gB_\mu}{\sqrt{g^2 + g'^2}} : m_A = 0$$

Higgs Boson Couplings



$$\Gamma(H \rightarrow f \bar{f}) = N_c \frac{G_F M_H}{4\pi\sqrt{2}} m_f^2, \quad N_C = 3 (1) \text{ for quarks (leptons)}$$

Weinberg 1967

$$\Gamma(H \rightarrow VV) = \frac{G_F M_H^3}{8\pi\sqrt{2}} F(r) \left(\frac{1}{2} \right)_Z, \quad r = \frac{M_V}{M_H}$$

Higgs 1966

A Phenomenological Profile of the Higgs Boson

- First attempt at systematic survey

A PHENOMENOLOGICAL PROFILE OF THE HIGGS BOSON

John ELLIS, Mary K. GAILLARD * and D.V. NANOPOULOS **
CERN, Geneva

Received 7 November 1975

A discussion is given of the production, decay and observability of the scalar Higgs boson H expected in gauge theories of the weak and electromagnetic interactions such as the Weinberg-Salam model. After reviewing previous experimental limits on the mass of

We should perhaps finish with an apology and a caution. We apologize to experimentalists for having no idea what is the mass of the Higgs boson, unlike the case with charm [3,4] and for not being sure of its couplings to other particles, except that they are probably all very small. For these reasons we do not want to encourage big experimental searches for the Higgs boson, but we do feel that people performing experiments vulnerable to the Higgs boson should know how it may turn up.

2011

Status of the Standard Model before the LHC

- Perfect agreement with all *confirmed* accelerator data
- Consistency with precision electroweak data (LEP et al) *only if there is a Higgs boson*
- Agreement seems to require *a relatively light Higgs boson* weighing $< \sim 180 \text{ GeV}$
- Raises many unanswered questions:
mass? flavour? unification?

Where are the top and Higgs?

Estimating Masses with Electroweak Data

- High-precision electroweak measurements are sensitive to quantum corrections

$$m_W^2 \sin^2 \theta_W = m_Z^2 \cos^2 \theta_W \sin^2 \theta_W = \frac{\pi\alpha}{\sqrt{2}G_F}(1 + \Delta r)$$

Veltman

- Sensitivity to top mass is quadratic:

$$\frac{3G_F}{8\pi^2\sqrt{2}}m_t^2$$

- Sensitivity to Higgs mass is logarithmic:


$$\frac{\sqrt{2}G_F}{16\pi^2}m_W^2\left(\frac{11}{3}\ln\frac{M_H^2}{m_Z^2} + \dots\right), M_H \gg m_W$$

- Measurements at LEP et al. gave indications first on top mass, then on Higgs mass


$$\Delta\rho = 0.0026\frac{M_t^2}{M_Z^2} - 0.0015\ln\left(\frac{M_H}{M_W}\right)$$

Precision Tests of the Standard Model

Lepton couplings




Pulls in global fit



2011

Combining Information from Previous Direct Searches and Indirect Data



$m_H = 125 \pm 10 \text{ GeV}$

Gfitter collaboration





“... we do not want to encourage big experimental searches for the Higgs boson, but ...”

EGN 1975

A la recherche
du
Higgs perdu ...

Higgs Production at the LHC




LHC Higgs Cross-Section
Working Group
(LHSWG)

Many production modes measurable if $M_H \sim 125 \text{ GeV}$

Higgs Production at the LHC


Cross sections for
Higgs mass near
125 GeV



LHC Higgs Cross-Section
Working Group
(LHXS WG)

Higgs Decay Branching Ratios

- Couplings proportional to masses (?)




- Important couplings through (quantum) loops:
 - gluon + gluon \rightarrow Higgs $\rightarrow \gamma\gamma$

Many decay modes measurable if $M_h \sim 125$ GeV

Higgs Decay Branching Ratios

Dominant decay branching ratios for $m_H \sim 125$ GeV

LHC Higgs Cross-Section Working Group (LHXS WG)




2012


The Discovery of the Higgs Boson



Mass Higgsteria




Interesting Events



Higgsdependence Day!



Scientists from around the World



MEMBER STATES

7889

Austria	117
Belgium	120
Bulgaria	96
Czech Republic	244
Denmark	67
Finland	111
France	868
Germany	1342
Greece	237
Hungary	76
Israel	65
Italy	2045
Netherlands	168
Norway	67
Poland	350
Portugal	127
Romania	134
Slovakia	124
Spain	447
Sweden	85
Switzerland	228
United Kingdom	771

OBSERVERS

2718

Japan	314
Russia	1187
USA	1217

ASSOCIATE MEMBERS

India	357	745
Lithuania	35	
Pakistan	65	
Turkey	173	
Ukraine	115	

ASSOCIATE MEMBERS IN THE PRE-STAGE TO MEMBERSHIP

118

Cyprus	26
Serbia	57
Slovenia	35

OTHERS

1872

Afghanistan	1	Bolivia	4	Egypt	31	Kazakhstan	5	Mongolia	2	Philippines	3	Thailand	22
Albania	3	Bosnia & Herzegovina	2	El Salvador	1	Kenya	3	Montenegro	11	Saint Kitts and Nevis	1	T.F.Y.R.O.M.	2
Algeria	14	Burundi	1	Estonia	15	Korea Rep.	185	Morocco	20	Saudi Arabia	2	Tunisia	5
Argentina	27	Cameroon	1	Georgia	46	Kyrgyzstan	1	Myanmar	1	Senegal	1	Uruguay	1
Armenia	19	Canada	161	Ghana	1	Latvia	2	Nepal	10	Singapore	4	Uzbekistan	4
Australia	31	Chile	20	Hong Kong	1	Lebanon	23	New Zealand	5	South Africa	56	Venezuela	10
Azerbaijan	10	China	510	Iceland	3	Luxembourg	2	Nigeria	3	Sri Lanka	6	Viet Nam	13
Bangladesh	11	Colombia	45	Indonesia	11	Madagascar	4	North Korea	1	Sudan	1	Zambia	1
Belarus	48	Croatia	41	Iran	51	Malaysia	15	Oman	3	Swaziland	1	Zimbabwe	2
Benin	1	Cuba	12	Iraq	1	Malta	9	Palestine (O.T.)	7	Syria	1		
		Ecuador	6	Ireland	16	Mauritius	1	Paraguay	2	Taiwan	51		
				Jordan	1	Mexico	82	Peru	7				




Russian naval shells reused
in the CMS experiment

LHC Measurements

Standard Model Total Production Cross Section Measurements

Status: October 2023




Higgs Measurements

July 4 2022




[CMS-HIG-19-015](#)
JHEP 07 (2021) 027





[CMS-HIG-19-001](#)
EPJC 81 (2021) 488

$m_H = 125.38 \pm 0.14$ (total) GeV




[CMS-PAS-HIG-19-005](#)


Observation independently in all 5 decay modes



[CMS-PAS-HIG-20-013](#)





[CMS-HIG-19-010](#)
Submitted to EPJC




[CMS-HIG-18-016](#)
PRL 121 (2018) 121801

It Walks and Quacks like a Higgs

- Do couplings scale \sim mass? With scale = v ?



Emerging Decay Mode: $Z \rightarrow H \gamma$



Signal strength $\mu = 2.2 \pm 0.7$ times Standard Model value

Negligible change in NLO QCD

Higher-order EW unimportant


Statistics? BSM physics?

Buccioni, Devoto, Djouadi, JE,
Quevillon, Tancredi, arXiv:2312.12384

Chen, Chen, Qiao & Zhu,
arXiv:2404.11441


QCD Corrections to $H \rightarrow Z\gamma$

NLO QCD diagrams for signal and background



NLO QCD increases cross-section by factor ~ 2

Negative interference – but blown up by factor 10 in plot




Reduces cross-section by 3%

$$\sigma_{\text{Sig}}^{\text{NLO}} = 1.207^{+20\%}_{-15\%} \text{ fb}, \quad \sigma_{\text{Int}}^{\text{NLOsv}} = -0.0344^{+12\%}_{-12\%} \text{ fb}$$


Higher-Order Higgs Couplings

- Standard Model Lagrangian contains $HHH, VVHH$ couplings in Higgs potential $V(H)$, Higgs kinetic term $|D_\mu H|^2$, respectively
- Directly related to (m_H, m_W) and VVH , respectively
- Absence/modification would destroy consistency (renormalizability) of Standard Model
- Could be modified by, e.g., higher-order terms in effective field theory, e.g., H^6 or $|H|^2|D_\mu H|^2$
- Parameterized by $\kappa_\lambda, \kappa_{2V}$, respectively
- **Measuring them is next frontier in Higgs measurements**


Search for Triple-H Coupling



Diagrams for double-Higgs production




Loop corrections to single Higgs production




Search for HHH Coupling

Limit on double-Higgs production




Limits on triple-Higgs coupling


From double-Higgs search




Combined limit



Evidence for VVHH Coupling





Evidence for VVHH Coupling




5 - σ exclusion of $\kappa_{2V} = 0$ if other Higgs couplings have Standard Model values

Evidence for VVHH Coupling



$\kappa_{2V} = 1.02 \pm 0.23$ if other Higgs couplings have Standard Model values

Prospects for Future Higgs Measurements



Strengths relative to SM predictions

R.K. Ellis et al (European Strategy), arXiv:1910.11775